

Lac Ste. Anne FIN Summary

2018

Background

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

Fall Index Netting (FIN)

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

How is this information used?

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is long-term sustainability, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status.

For more information, please see Alberta’s FIN and FSI websites,

- <https://www.alberta.ca/fall-index-netting.aspx>
- <https://www.alberta.ca/fish-sustainability-index.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Northern Pike using the standardized Fall Index Net (FIN) method. Note: Thresholds align with species management frameworks.

Mature Walleyes/net	Mature Pike/net	Risk to Sustainability
>29.0	>21.8	Very Low
20.3-29.0	15.3-21.8	Low
14.5-20.2	10.9-15.2	Moderate
5.8-14.4	4.4-10.8	High
<5.8	<4.4	Very High

Results of the 2018 FIN at Lac Ste. Anne

Lac Ste. Anne (5659 ha) is located 58 km west from the city of Edmonton. From September 19 to 21, 2018, 12 nets captured 105 Lake Whitefish, 3 Longnose Suckers, 105 Northern Pike, 319 Walleyes, 23 White Suckers and 36 Yellow Perch.

Walleye

The mean catch rate of Walleyes was 26.6/ net-night. The catch rates of mature (Figure 1) and immature Walleyes were 22.1/ net-night and 4.3/ net-night, respectively. The corresponding FSI score for the current mature density of Walleyes was assessed at **low risk**.

The length distribution indicates somewhat unstable yet abundant recruitment, abundant 380 to 460 mm fish, and very abundant fish larger than 460 mm (Figure 2).




Figure 1 - The FIN catch rate of mature Walleyes from Lac Ste. Anne, 2018. Dashed line is the mean catch rate (22.1 fish/ net-night), with individual net data as hollow circles (n=12 nets).




Figure 2 – FIN sample of showing size of Walleyes from Lac Ste. Anne, 2018. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.

The 2018 FIN sample represented approximately 0.4% of the estimated mature Walleye population size.

Northern Pike

The mean catch rate of mature Northern Pike was 8.8/ net-night (Figure 3). The corresponding FSI score for the current mature density of Northern Pike was assessed at **high risk**.

The length distribution indicates weak recruitment, a moderate abundances of fish from 450 to 650 mm and no pike larger than 650 mm (Figure 4).

The 2018 FIN sample represented approximately 0.1% of the estimated mature Northern Pike population size.

Summary

Since the 2016 assessment, the corresponding FSI status of the Walleye population in Lac Ste. Anne has decreased from **very low** to **low** risk. The Walleye stock provides opportunities for carefully managed harvests with Special Harvest Licenses.




Figure 3 - The FIN catch rate of mature Northern Pike from Lac Ste. Anne, 2018. Dashed line is the mean catch rate (8.8 fish/ net-night), with individual net data as hollow circles (n=12 nets).




Figure 4 – FIN sample showing size of Northern Pike from Lac Ste. Anne, 2018. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

Since 2014, the FIN assessments have indicated the Northern Pike population is at a FSI status of **high risk**. To ensure the recovery and long-term sustainability of this population, conservation-based management is necessary.

Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.