

Background

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

Fall Index Netting (FIN)

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

How is this information used?

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is **long-term sustainability**, shown by the red lines on the graphs below.

Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information please see Alberta’s FIN and FSI websites,

- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fall-index-netting/default.aspx>
- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fish-sustainability-index/default.aspx>

*Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Pike using the standardized Fall Index Net (FIN) method. **Note:** Thresholds align with species management frameworks.*

Mature Walleyes / net	Mature Pike / net	Risk to Sustainability
>29.0	>21.8	Very Low
20.3-29.0	15.3-21.8	Low
14.5-20.2	10.9-15.2	Moderate
5.8-14.4	4.4-10.8	High
<5.8	<4.4	Very High

Results of the 2013 FIN at Fawcett Lake

Fawcett Lake (3445 ha) is located approximately 240 km north of the city of Edmonton. From September 29 to October 2, 2013, fourteen FIN nets captured 80 Cisco, 7 Lake Whitefish, 78 Northern Pike, 271 Walleye, 3 Longnose Suckers, 46 White Suckers, and 109 Yellow Perch.

Walleye

The mean catch rate of Walleyes was 19.4/net-night. The catch rates of mature (Figure 1) and immature Walleye were 11.7/net-night and 7.3/net-night, respectively. The corresponding FSI score for the current mature density of Walleye was assessed at **high risk**.

The length distribution shows moderate recruitment, and a moderate abundance of Walleye larger than 360 mm (Figure 2).

The 2013 FIN sample represented approximately 0.8% of the estimated mature Walleye population.




Figure 1 - The FIN catch rate of mature Walleyes from Fawcett Lake, 2013. Dashed line is the mean catch rate (11.7 fish/net-night), with net data as hollow circles (n=14 nets).




Figure 3 - The FIN catch rate of Northern Pike from Fawcett Lake, 2013. Dashed line is the mean catch rate (5.3/net-night), with net data as hollow circles (n=14 nets).




Figure 2 – FIN sample of showing size of Walleyes from Fawcett Lake, 2013. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.




Figure 4 – FIN sample showing size of Northern Pike from Fawcett Lake, 2013. The red line indicates the average length distribution of Pike from 6 Alberta lakes supporting long-term sustainable harvests of Pike.

Northern Pike

The mean catch rate of mature Northern Pike was 5.3/net-night (Figure 3). The corresponding FSI score for the current mature density of Northern Pike was assessed at **high risk**.

The length distribution shows low and unstable recruitment, and low densities of Northern Pike larger than 400 mm (Figure 4). The low density and truncation of larger Pike indicates a growth overfished population.

The 2013 FIN sample represented approximately 0.2 % of the estimated mature Northern Pike population.

Summary

Since the FINs on Fawcett Lake in 2006, 2011 and 2013 the adult abundance has been low, but stable with a corresponding FSI status of **high risk**. Continued

conservation-focused management is necessary to support the long-term sustainability of this fishery. Since the FIN on Fawcett Lake in 2006 the corresponding FSI status of mature Northern Pike has remained at **high risk**.

Dependant on the management objective, strict conservation-focused management is necessary to recover the long-term sustainability of this fishery.

Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.