

**Background**

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

**Fall Index Netting (FIN)**

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

**How is this information used?**

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is **long-term sustainability**, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to

determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information please see Alberta’s FIN and FSI websites,

- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fall-index-netting/default.aspx>
- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fish-sustainability-index/default.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Northern Pike using the standardized Fall Index Net (FIN) method. **Note:** Thresholds align with species management frameworks.

Mature Walleyes/net	Mature Pike/net	Risk to Sustainability
>29.0	>21.8	Very Low
20.3-29.0	15.3-21.8	Low
14.5-20.2	10.9-15.2	Moderate
5.8-14.4	4.4-10.8	High
<5.8	<4.4	Very High

**Results of the 2017 FIN at Lake Newell**

Lake Newell (12431 ha) is located approximately 210 km southeast from the city of Calgary. From September 25-26, 2017, six gill nets captured 33 Lake Whitefish, 8 Northern Pike, 85 Walleyes, 2 Spottail Shiners, and 4 Yellow Perch.

**Walleye**

The mean catch rate of Walleyes was 14.2/net-night. The catch rates of mature (Figure 1) and immature Walleyes were 10.3/net-night and 3.7/net-night, respectively. The corresponding FSI score for the current mature density of Walleyes was assessed at **high risk**.

The length distribution shows strong, though intermittent recruitment, very low abundances of Walleyes 220 to 460 mm, and abundant fish over 500 mm. (Figure 2). Lake Newell is an irrigation reservoir that experiences fluctuating water levels which limit fish habitats. This likely contributes to unstable recruitment.

The 2017 FIN sample represented approximately 0.1% of the estimated mature Walleye population size.

**Northern Pike**

The mean catch rate of mature Northern Pike was 1.3/net-




Figure 1 - The FIN catch rate of mature Walleyes from Lake Newell, 2017. Dashed line is the mean likelihood catch rate (10.3 fish/net-night), with individual net data as hollow circles (n=6 nets).




Figure 3 - The FIN catch rate of mature Northern Pike from Lake Newell, 2017. Dashed line is the mean likelihood catch rate (1.3 fish/net-night), with individual net data as hollow circles (n=6 nets).




Figure 2 – FIN sample of showing size of Walleyes from Lake Newell, 2017. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.




Figure 4 – FIN sample showing size of Northern Pike from Lake Newell, 2017. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

night (Figure 3). The corresponding FSI score for the mature density of Northern Pike was assessed at **very high risk**.

The length distribution shows no recent recruitment, no pike <580 mm, and low abundances across remaining size classes (Figure 4). This fishery is entirely supported by a few weak adult year classes. While reservoir levels are generally stable in Lake Newell, water levels can decrease significantly during drought years. This can negatively influence fish habitat and recruitment.

The 2017 FIN sample represented approximately 0.02% of the estimated mature Northern Pike population size.

### Summary

The abundance of mature Walleyes in Lake Newell has

increased from an FSI status of **very high risk** in 2005 to **high risk** from 2006 to 2017. Conservation-focused management likely remains necessary to ensure the long-term sustainability of the fishery.

Since 2006, Lake Newell has remained at an FSI status of **very high risk** for mature Northern Pike. Stringent conservation efforts are necessary to maintain this population.

### Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.