

# Routine Childhood Immunization Schedule

Effective December 17, 2018

(Refer to [biological pages](#) for specific vaccine indications and information)

Vaccine /Age	Birth	2 mos	4 mos	6 mos	12 mos	18 mos	4 yrs	Gr 6	Gr 9
<a href="#">DTaP-IPV-Hib</a> Diphtheria, Tetanus, Acellular Pertussis, Polio, <i>Haemophilus influenzae</i> type b		1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose		4 <sup>th</sup> dose			
<a href="#">DTaP-IPV-Hib-HB</a> Diphtheria, Tetanus, Acellular Pertussis, Hepatitis B, Polio, <i>Haemophilus influenzae</i> type b		1 <sup>st</sup> dose Infants born on or after March 1, 2018, Infants born to Hep B infected mothers, and Infants born to families from endemic countries.	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose*	*3 <sup>rd</sup> dose should not be administered before 24 weeks (6 mos) of age.				
<a href="#">PneuC-13</a> Pneumococcal Conjugate 13		1 <sup>st</sup> dose	2 <sup>nd</sup> dose		3 <sup>rd</sup> dose				
<a href="#">Rotavirus</a>		1 <sup>st</sup> dose	2 <sup>nd</sup> dose	3 <sup>rd</sup> dose					
<a href="#">MenconC</a> Meningococcal Conjugate C			1 <sup>st</sup> dose		2 <sup>nd</sup> dose				
<a href="#">MMR-Var</a> Measles, Mumps, Rubella, Varicella					1 <sup>st</sup> dose Minimum age 12 mos.		2 <sup>nd</sup> dose		
<a href="#">dTAp-IPV</a> Diphtheria, Tetanus, Acellular Pertussis, Polio							Booster		
<a href="#">HBV</a> Hepatitis B	1 <sup>st</sup> dose Infants born to Hep B infected mothers							Three Doses 0, 1, and 6 mos. (if not previously received in childhood)	
<a href="#">HPV-9</a> Human Papillomavirus -Healthy individuals								Two Doses 0 and 6 mos.	
-Immunocompromised and those 15 yrs and older								Three Doses 0, 2, and 6 mos.	
<a href="#">MenC-ACYW</a> Meningococcal Conjugate ACYW-135									One dose
<a href="#">dTAp</a> Diphtheria, Tetanus, Acellular Pertussis									Booster
<a href="#">Td</a> Tetanus, Diphtheria									
<a href="#">Pneumo-P</a> Pneumococcal Polysaccharide 23									
<a href="#">Influenza</a>					Annually starting at 6 months of age. Children under 9 years of age with no previous influenza immunization require 2 doses at least 4 weeks apart.				

- This Immunization Schedule is published by the Chief Medical Officer of Health in accordance with s. 18.5 of the [Public Health Act](#).
- A health practitioner who performs an immunization listed in this Immunization Schedule **must** comply in the following manner:
  - Vaccines **must** be provided to clients/patients no sooner than at the ages set out in this Immunization Schedule. To the greatest extent possible, doses should be provided at the ages set out.
  - Vaccines **must** be provided to clients/patients no sooner than at the intervals set out in this Immunization Schedule. The intervals represent the optimal amount of time that **must** pass before a subsequent dose of vaccine is administered. To the greatest extent possible, doses should be provided at the intervals set out.
  - Despite (A) and (B), a health practitioner who performs immunizations may provide vaccines earlier or later than the ages required in (A) and the intervals required in (B) if:
    - any such variation is necessary:
      - to reduce a risk of acquiring a disease, or
      - because it is medically indicated for the client/patient, or
      - because of a vaccine shortage, and
    - The variation is approved in advance by a Medical Officer of Health.

Refer individual to public health for any variation in schedule.

## Routine Childhood Immunization Schedule