

### Section 1: Summary

Influenza activity continues to increase across the province after higher than usual activity this summer. The activity continues to be higher than the average seasonal curve from the past few years. The predominant subtype so far this season is Influenza A (H3N2).

### Section 2: Outbreaks

Three influenza outbreaks were declared in week 43, one in Calgary Zone, one in Edmonton Zone and one in Central Zone. There have been 8 outbreaks reported to date.


	Outbreaks	
	Current Week	YTD
North	0	0
Edmonton	1	1
Central	1	1
Calgary	1	5
South	0	1
Unknown	0	0
Alberta	3	8

### Section 3: Laboratory-Confirmed Influenza Surveillance & Peak Prediction

To date in the 2017/18 season, the Provincial Laboratory for Public Health (ProvLab) has reported 300 lab-confirmed influenza cases: 257 influenza A, 31 influenza B and 12 where subtype is unavailable. The predominant subtype is influenza A (H3N2) with 74% of all isolates. While more than half of the lab-confirmed cases have been in Calgary Zone, cases have been reported in all zones.

Peak prediction for influenza will commence once there is enough laboratory data to support it. The median peak for an Influenza A (H3N2)-predominant season is week 52. Please see data notes for additional details on the seasonal threshold and peak prediction.


2017/18 Lab-Confirmed Influenza Cases by Serotype and Week, Compared to Seasonal Average



## Section 4: Hospitalizations


There have been 90 hospitalized influenza cases to date. The predominant subtype in hospitalized patients is Influenza A (H3N2).

### Hospitalized Influenza Cases in 2017/18, by Serotype and Week



## Section 5: Antiviral Prescriptions

The number of antivirals dispensed by community pharmacists provides an indication of the amount of influenza circulating in the community. There were 388 prescriptions dispensed in week 43. There have been 1,222 prescriptions dispensed to date.




## Section 6: Antiviral Resistance

As of Sept. 1, 2017, 45 influenza isolates had been characterized by the National Medical Laboratory (NML). All isolates were sensitive to zanamivir and oseltamivir.

## Section 7: Influenza-Like Illness – TARRANT

Sentinel physicians report cases of ILI seen in their practices. Sentinel physicians saw 10 cases of influenza-like illness in week 43 (0.2% of all patients seen). There were 11 cases of ILI (0.2%) seen by physicians in week 42.

	Sentinel Doctors (#)	Sentinel Recorders This Week (#)	Patients Seen (#)	ILI Cases (#)	Patients with ILI (%)
<b>Calgary Zone</b>	61	34	2,468	5	0.2
<b>Central Zone</b>	13	9	449	1	0.2
<b>Edmonton Zone</b>	10	8	1,149	2	0.2
<b>Northern Zone</b>	6	2	70	0	0.0
<b>South Zone</b>	12	9	1,181	2	0.2
<b>Alberta</b>	102	62	5,315	10	0.2



For more information, please go to: [www.tarrantviralwatch.ca](http://www.tarrantviralwatch.ca)

## Section 8: Acknowledgments

Alberta Health would like to thank Alberta Health Services (AHS), First Nations Inuit Health Branch (FNIHB), the Provincial Laboratory for Public Health (ProvLab), the National Microbiology Laboratory (NML), and TARRANT sentinel physician system for their partnership in influenza surveillance in Alberta.

## Section 9: Canadian and International Influenza Activity

---

The following links provide access to other websites related to influenza and ILI.

- Alberta Health (Influenza Reports) <http://www.health.alberta.ca/professionals/influenza-evidence.html>
- Alberta Health (Influenza Forms) <http://www.health.alberta.ca/professionals/notifiable-diseases-guide.html>
- Alberta Health Services <http://www.albertahealthservices.ca/2891.asp>
- Canada – FluWATCH [www.phac-aspc.gc.ca/fluwatch/index-eng.php](http://www.phac-aspc.gc.ca/fluwatch/index-eng.php)
- World Health Organization [www.who.int/topics/influenza/en/](http://www.who.int/topics/influenza/en/)
- USA – CDC [www.cdc.gov/flu/weekly/](http://www.cdc.gov/flu/weekly/)

## Section 10: Data Notes

---

This report utilized data from the Provincial Laboratory for Public Health (ProvLab), Alberta's influenza like illness (ILI) sentinel physician system (TARRANT), Supplemental Enhanced Service Event (SESE) physician claims data, the Pharmacy Information Network (PIN), as well as outbreak reports and hospitalized case report forms from Alberta Health's Communicable Disease Reporting System (CDRS).

### Defining Baselines and Thresholds for Lab-Confirmed Influenza Surveillance

An important function of influenza surveillance is to determine whether the timing or magnitude of the influenza season is unusual compared to previous seasons. The World Health Organization recommends comparing current-season influenza activity to the average epidemic curve, and to create two thresholds – one to determine when the influenza season has begun (seasonal threshold) and to determine at what point current influenza activity would be unusually high (alert threshold)<sup>1</sup>.

The average epidemic curve for lab-confirmed influenza surveillance was estimated using data collected between the 2010/11 and 2014/15 seasons. Lab-confirmed influenza became routinely reported in 2009; however, the 2009/10 pandemic season was excluded. The peaks of each season were aligned, and the average number of cases reported per aligned week was calculated as well as the 90 per cent confidence limit. The 90 per cent confidence limit acts as the Alert Threshold. If the number of cases reported in a week passes this threshold it is considered to be unusually high. Until the current season has peaked, the best estimate of the peak is utilized to align the current season to the average epidemic curve. The first estimate utilized was the median of the previous five seasons, week<sup>2</sup>.

The seasonal threshold was estimated as the average number of cases reported per week, in weeks considered outside of the influenza season (i.e., pre-season weeks). The start of the influenza epidemic was defined as the week in which the percent positivity of influenza A laboratory tests at ProvLab were  $\geq 10$  per cent of all respiratory laboratory tests ordered that week<sup>2</sup>. The end of the influenza epidemic was defined as the week in which the per cent positivity of influenza B was  $< 10$  per cent of all respiratory laboratory tests ordered that week<sup>2</sup>. All weeks in between these time periods were considered part of the influenza epidemic. Per cent positivity is an accepted method of determining the influenza season<sup>1</sup>, however visual inspection was also utilized to ensure face validity.

### Predicting the Peak of the Epidemic Curve

The peak of the influenza curve was predicted using a deterministic mathematical model that considered transmission across by three age groups (0–18 years, 19–64 years, and  $\geq 65$  years old). The data used to estimate the parameters in the model is obtained from PSI, CDRS, physician claims data, PIN, and the population registry. The model is simulated weekly using up-to-date laboratory confirmed data to predict the peak for influenza. As more laboratory confirmed counts become available the point estimate for the peak may vary and the width of the confidence interval will decrease.

Note that the current model assumes that influenza activity is similar across zones, which is not the case this season and is the reason why the confidence intervals around the mean are wide.

---

<sup>1</sup> World Health Organization 2013 "Global Epidemiological Surveillance Standards for Influenza". Geneva. [http://www.who.int/influenza/resources/documents/influenza\\_surveillance\\_manual/en/](http://www.who.int/influenza/resources/documents/influenza_surveillance_manual/en/)

<sup>2</sup> Provincial Laboratory for Public Health (ProvLab) Weekly Respiratory Summary