



# **Alberta tier 1 soil and groundwater remediation guidelines**



<b>Title:</b>	<b>Alberta Tier 1 Soil and Groundwater Remediation Guidelines</b>
<b>Number:</b>	<b>AEP, Land Policy, 2022, No. 4</b>
<b>Program Name:</b>	<b>Land Policy</b>
<b>Effective Date:</b>	<b>January 1, 2023</b>
<b>This document was updated on:</b>	
<b>ISBN No.</b>	<b>ISBN: 978-1-4601-5501-1</b>
<b>Disclaimer:</b>	

Original signed by: \_\_\_\_\_

Date: August 24, 2022

Scott Milligan, Executive Director

Land Policy Branch

Environment and Parks

This publication is issued under the Open Government Licence – Alberta  
(<http://open.alberta.ca/licence>).

This publication is available online at <https://open.alberta.ca/publications/1926-6243>

Alberta Tier 1 Soil and Groundwater Remediation Guidelines | Environment and Parks  
© 2022 Government of Alberta | August 26, 2022 | ISBN 978-1-4601-5501-1

# Contents

- 1. Introduction .....6
  - 1.1 General.....6
  - 1.2 Relation to Alberta’s Framework for the Management of Contaminated Sites.....6
  - 1.3 Relationship to Other Guideline Documents.....6
  - 1.4 Ongoing Commitment.....7
- 2. Alberta General Policy for Use of Tier 1 .....8
  - 2.1 Principles of Contaminant Management in Alberta .....8
  - 2.2 Soil Protection in Alberta .....9
  - 2.3 Groundwater Protection in Alberta..... 11
- 3. Alberta Framework for the Management of Contaminated Sites..... 14
  - 3.1 Implementation Framework and Management Levels ..... 14
  - 3.2 Land Use Definitions..... 14
  - 3.3 Groundwater and Surface Water Use Definitions ..... 15
  - 3.4 Conditions and Restrictions Associated with Tier 2 and Exposure Control..... 15
- 4. Scientific Basis for Developing Tier 1 Remediation Guidelines ..... 17
  - 4.1 Risk-Based Guidelines ..... 17
  - 4.2 Protection of Human Health..... 17
  - 4.3 Ecological Protection ..... 18
  - 4.4 Calculation of Guidelines ..... 19
  - 4.5 Determination of Overall Guideline ..... 19
  - 4.6 Integration of New and Existing Guidelines ..... 19
- 5. User Guidance for Tier 1 Management..... 20
  - 5.1 Information Requirements for Tier 1 (Tier 1 Assessment) ..... 20
  - 5.2 Identification of Appropriate Tier 1 Guidelines (Tier 1 Evaluation) ..... 25
  - 5.3 Tier 1 Decision and Management..... 28
- 6. References..... 29

### **List of Tables**

Table 1	Summary of Alberta Tier 1 Soil Remediation Guidelines
Table 2	Summary of Alberta Tier 1 Groundwater Remediation Guidelines
Table 3	Summary of Alberta Tier 1 Subsoil Remediation Guidelines (BTEX and PHCs only)
Table 4	Alberta Tier 1 Salt Remediation Guidelines

### **List of Figures**

Figure 1	Example of 30 m Buffer Zone Adjacent to More Sensitive Land Use
Figure 2	Implementation Framework for Tier 1, Tier 2 and Tier 3 Guidelines
Figure 3	Expanded Flow Diagram – Tier 1

### **List of Appendices**

Appendix A	Soil Remediation Guidelines – All Exposure Pathways
Appendix B	Groundwater Remediation Guidelines – All Water Uses
Appendix C	Protocols for Calculating Alberta Tier 1 Soil and Groundwater Quality Guidelines

# 1. Introduction

## 1.1 General

This document focuses on Tier 1 Soil and Groundwater Remediation Guidelines (Tier 1), the process used to develop them, and the manner in which they should be applied. Tier 1 guidelines are generic, risk-based remediation guidelines for contaminated site management in Alberta. Generic guidelines are designed using relatively conservative assumptions to be applicable at most sites in Alberta. Risk-based guidelines combine information on contaminant characteristics with site information and human and ecological receptor characteristics to identify contaminant levels that pose minimal risk.

The Tier 1 guidelines are simple tabular values. Some site information is needed to ensure that site conditions are adequately represented by the assumptions used to develop the Tier 1 guidelines. This document provides numerical Tier 1 guidelines, with a detailed explanation of the underlying rationale and requirements for applying them to contaminated sites.

Those sites that may be more sensitive (see Section 5) than conditions assumed in Tier 1 are more appropriately dealt with under Tier 2. When a site has characteristics that make it more sensitive than the Tier 1 assumptions, the resulting Tier 2 guidelines may be more restrictive than Tier 1 values.

When adverse effects are evident, contaminants must be managed to alleviate adverse effects, regardless of whether a site meets Tier 1 or 2 remediation guidelines.

This document supports the Government of Alberta's *Contaminated Sites Policy Framework* (ESRD, 2014c) and acts as a companion document to the *Alberta Tier 2 Soil and Groundwater Remediation Guidelines* (Tier 2) (ESRD, 2007 as amended), *Alberta Exposure Control Guide* (AEP, 2016b), and *Alberta Environmental Site Assessment Standard* (AEP, 2016a).

## 1.2 Relation to Alberta's Framework for the Management of Contaminated Sites

Alberta's framework for contaminated sites management is described in the *Contaminated Sites Policy Framework* (ESRD, 2014c).

This document further describes the principles for management of substance releases, focusing on the Alberta Tier 1 option for management of contaminated sites. While it focuses on the Alberta Tier 1 guidelines, it provides additional support for Alberta Tier 2 and Exposure Control options. Proponents will need to ensure that protection levels and model inputs are consistent with Alberta Tier 1 guidelines when using other options provided in the framework.

All submissions for site closure based the application of Tier 1 guidelines or using Tier 1 guidelines as part of a risk assessment or environmental site assessment must adhere to this document.

## 1.3 Relationship to Other Guideline Documents

The soil and groundwater guidelines in this document supersede the following guideline documents:

- *Alberta Tier 1 Criteria for Contaminated Soil Assessment and Remediation* (AEP, 1994)
- *Alberta Soil and Water Quality Guidelines for Hydrocarbons at Upstream Oil and Gas Sites* (AENV, 2001a); and
- *Risk Management Guidelines for Petroleum Storage Tank Sites* (AENV, 2001b).

As a component of the *Contaminated Sites Policy Framework* (ESRD, 2014c), this document, together with the following documents, forms the basis for contaminated site management in Alberta:

- *Alberta Tier 2 Soil and Groundwater Remediation Guidelines* (ESRD, 2007 as amended);
- *Alberta Environmental Site Assessment Standard* (AEP, 2016a);
- *Alberta Exposure Control Guide* (AEP, 2016b);

The following documents contain additional information needed for the application of Tier 1 guidelines and should be used in conjunction with this document:

- *Alberta Environmental Site Assessment Standard* (AEP, 2016a);
- *Guidance Manual for Environmental Site Characterization in Support of Environmental and Human Health Risk Characterization, Volumes 1 to 4* (CCME, 2016);

- *Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils* (AENV, 2011);
- *Directive for Monitoring the Impact of Sulphur Dust on Soils* (AEP, 2015)
- *Environmental Quality Guidelines for Alberta Surface Waters* (ESRD, 2014a and updates);
- *Salt Contamination Assessment and Remediation Guidelines* (AENV, 2001c);
- *Soil Remediation Guidelines for Barite: Environmental Health and Human Health* (AENV, 2009);
- *Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine* (AENV, 2010);
- *Soil Remediation Guidelines for Boron: Environmental and Human Health* (AEP, 2015);
- *Assessing Drilling Waste Disposal Areas: Compliance Options for Reclamation Certification* (AER, 2014a)
- *Canadian Environmental Quality Guidelines* (CCME, 1999 and updates);
- *Reference Method for the Canada-Wide Standard for Petroleum Hydrocarbons in Soil - Tier 1 Method* (CCME, 2001); and
- *Canadian Guidelines for the Management of Naturally Occurring Radioactive Materials (NORM)* (Health Canada, 2013).

The following documents provide the protocols and the primary technical basis for the risk-based guidelines calculated in this volume:

- For all compounds except petroleum hydrocarbons: *A Protocol for the Derivation of Environmental and Human Health Soil Quality Guidelines* (CCME, 2006);
- For petroleum hydrocarbons: *Canada-Wide Standards for Petroleum Hydrocarbons (PHCs) in Soil. Scientific Rationale* (CCME, 2008);
- *A Protocol for the Derivation of Groundwater Quality Guidelines for Use at Contaminated Sites* (CCME, 2015); and
- *A Protocol for the Derivation of Soil Vapour Quality Guidelines for Protection of Human Exposures Via Inhalation of Vapours* (CCME, 2014).

## 1.4 Ongoing Commitment

Tier 1 is a living document. Alberta Environment and Parks is committed to updating the guidelines as new information and guidelines for new substances become available.

## **2. Alberta General Policy for Use of Tier 1**

### **2.1 Principles of Contaminant Management in Alberta**

A contaminant is a substance that is present in an environmental medium in excess of natural background concentration (CCME, 2006). Three key elements of Alberta's framework for the management of contaminated sites (ESRD, 2014c) are: source control, contamination delineation, and contaminant management, including remediation.

#### **2.1.1 Source Control**

A source of contamination is anything that adds contaminant mass to the environment. Source control is a necessary action in support of pollution prevention, a key outcome of Alberta's policy on contaminated site management. If there is evidence of soil or groundwater contamination, the source, if it is still present, must be identified. Sources must be removed or controlled as soon as practicable.

Contaminants can be introduced into the environment in a number of ways. Leaking pipelines and storage tanks are common sources of contamination. Contaminated soil and groundwater may be a source of contamination to other areas of a site if the contaminants are mobile. Removal or management of these sources is a necessary part of contaminated site management. Soil or groundwater with naturally elevated substance concentrations may become a source of contamination if it is redistributed and causes the receiving soil or water to exceed Tier 1 or 2 remediation guidelines. This can be avoided by proper characterization and handling of soil and groundwater prior to redistribution. Failure to control sources allows contaminants to spread, increasing risk and remediation costs, and potentially limiting future land use if remediation to Tier 1 or 2 guidelines is not possible.

Where complete source removal is not feasible, the source must be removed to the extent possible and treatment, control, and/or management measures must be implemented to address the residual source. Treatment measures will assist in the ongoing reduction of source contaminant concentrations. Source control measures must prevent the contaminant from spreading to adjacent areas, causing the soil or groundwater there to exceed Tier 1 or 2 guidelines.

If source control measures are required, they must operate until the source meets Tier 1 or Tier 2 remediation guidelines. Source control must be supported by a monitoring program that demonstrates its efficacy.

Where source control rather than source remediation is implemented, a site is considered to be managed through an Exposure Control approach rather than a Tier 1 or Tier 2 approach.

#### **2.1.2 Delineation**

When soil or groundwater is found to contain contaminants in excess of Tier 1 or Tier 2 using the pathway exclusion approach, a delineation program must be implemented (Fig. 3). The delineation program must identify both the horizontal and vertical extent of contaminant concentrations exceeding the Tier 1 guidelines or the Tier 2 guidelines derived by the pathway exclusion approach (see Section 5 below and Table 5 in ESRD (2007 and updates)). Adequate delineation provides information needed to support appropriate decisions about contaminant remediation and management. Delineation programs must be extensive enough in both horizontal and vertical directions to allow all applicable exposure pathways and receptors to be properly assessed. Where soil contamination has reached groundwater or where a groundwater contaminant plume is shallow enough to impact surface receptors, delineation must include an investigation of contaminant concentrations in both soil and groundwater. Delineation is complete when measured concentrations are less than Tier 1 or 2 remediation guidelines.

Complete delineation must be accomplished prior to undertaking remediation. The only exception is for relatively simple sites where contaminants are removed by excavation and compliance with Tier 1 is shown by post-excavation sampling. When confirmatory samples fail to comply with Tier 1 guidelines after excavation is complete, full delineation of the remaining contamination must be undertaken and used to develop further remediation actions or risk management programs.

#### **2.1.3 Contaminant Management**

When the volume of soil or groundwater containing contaminant concentrations that exceed Tier 1 or Tier 2 using the pathway exclusion approach is completely delineated, a plan must be developed to remediate or otherwise manage the contaminants in a manner that is consistent with the framework outlined in ESRD (2014c). Using dilution to reduce contaminant concentrations is not an acceptable form of management, unless authorized by the appropriate regulatory authority under an operating approval, code of practice, or directive.



### 2.1.4 Management of Contaminants in Subsoil

In general, Tier 1 soil remediation guidelines (Table 1) apply to all soil regardless of depth. However, subsoil guidelines are provided for petroleum hydrocarbons (Table 3) and salinity (Table 4). These guidelines may be used at Tier 1 under the following circumstances:

#### Salinity

Topsoil guidelines for electrical conductivity and sodium adsorption ratio must be applied to the L, F, H, O, and A horizons (Soil Classification Working Group, 1998) or equivalent surficial material where these horizons are not present. Subsoil guidelines may be applied below the A horizon or equivalent in lieu of topsoil guidelines. Further information is available in the *Salt Contamination Assessment and Remediation Guidelines* (AENV, 2001c).

#### Petroleum Hydrocarbons

Soil guidelines in Table 1 for petroleum hydrocarbon fractions F1 to F4, benzene, toluene, ethylbenzene, and xylenes must be applied to a minimum depth of 1.5 metres. Subsoil guidelines in Table 3 may be used as follows:

- below 3 metres in depth at any site (in this case, surface soil guidelines must be applied to a depth of 3 metres),
- below 1.5 metres in depth within a 5 metre setback from an oilfield wellhead (see Directive 079, *Surface Development in Proximity to Abandoned Wells* (AER, 2014b)), or
- below 1.5 m at remote forested sites in the Green Area with fine textured soil irrespective of the distance to a wellhead. In order to use subsoil guidelines below 1.5 m outside of a 5 metre radius from the wellhead, sites must meet the requirements in the *Subsoil Petroleum Hydrocarbon Guidelines for Remote Forested Sites in the Green Area* (ESRD, 2014b).

The Tier 1 approach allows the exclusion of the ecological direct contact pathway for petroleum hydrocarbon fractions F1 to F4 for any land use below a depth of 3 metres. All other exposure pathways apply. Consistent with the *Canada-Wide Standard for Petroleum Hydrocarbons in Soil* (CCME, 2008), exclusion of the ecological direct soil contact pathway is permitted below 3 metres with the application of “management limits” for petroleum hydrocarbon fractions that represent maximum concentrations that apply at any depth. Management limits for petroleum hydrocarbon fractions F1 to F4 are provided in Appendix A. For sites that meet the requirements of the *Subsoil Petroleum Hydrocarbon Guidelines for Remote Forested Sites in the Green Area* (ESRD, 2014b) management limits provided in ESRD (2014b) may be applied.

## 2.2 Soil Protection in Alberta

### 2.2.1 Using Tier 1 Soil Remediation Guidelines

The goal of the Tier 1 soil remediation guidelines is to provide numerical targets for remediation of contaminated soil. To ensure consistency with “pollution prevention”, a key outcome of Alberta’s contaminated sites management framework, the Tier 1 soil remediation guidelines are not “pollute-up-to” levels. Care must be taken to prevent soil contamination and, when a substance release occurs, prompt actions must be taken to remediate or otherwise manage the release. Sources must not be left uncontrolled until cumulative releases result in an exceedance of Tier 1 or Tier 2 soil remediation guidelines. This results only in further contamination, increased remediation costs, and potential loss of land use options. Source control is a crucial component of pollution prevention.

### 2.2.2 Background Soil Quality

For the purpose of applying Tier 1 or Tier 2, the background concentration of a substance in soil or groundwater is defined as:

- The natural concentration of that substance in the absence of any input from anthropogenic activities or sources or;
- The background concentration in the surrounding area as a result of generalized non-point anthropogenic sources.

In some situations, the background concentration of a substance can be a significant proportion of, or even exceed, the Tier 1 guidelines. In cases when the background concentration is demonstrated to be greater than Tier 1 guidelines, the remediation level shall be set to background or to guidelines developed using Tier 2 procedures.

The definition for background cannot be used to eliminate point source emissions, anthropogenic activities that cause redistribution of soil or water sources with elevated substance concentrations, or non-point anthropogenic sources that result from activities at the site in question. For example, surface soils in urban areas that have variable levels of polycyclic aromatic hydrocarbons (PAHs) as a result of generalized automobile emissions can be considered as background based on the definition; however, additional PAH contamination may result from industrial activities at the site in question and the latter cannot be considered part of the urban background.

Similarly, some sites may have elevated electrical conductivities in the surface soil or groundwater due to natural conditions at the site. While this is considered natural background, if material with elevated conductivity is brought to the surface from deeper sediments or groundwater due to anthropogenic activities, this must be assessed as a contaminant of potential concern.

Background concentrations will vary with soil parent material, soil depth, and hydrologic regime. These factors lead to spatial variations in background concentrations that may or may not be predictable. To gain a good understanding of background conditions at a site, it is necessary to take sufficient representative samples from soils with similar characteristics to the affected site, but which are taken from outside the area affected by contamination. Sample depth and landscape position, soil profile characteristics and parent material should be recorded for all samples.

### **2.2.3 Land Use**

Potential receptors and their exposure to soil contaminants are affected by land use. For example, workers on an industrial site experience a different exposure than a toddler living on a residential property. Different ecological receptors are found in a forest setting than on an urban commercial property. Tier 1 soil remediation guidelines are calculated for five types of land use: natural areas, agricultural, residential/parkland, commercial, and industrial. These land use types may not correspond exactly to the range of municipal zoning options, but by evaluating the types of receptors and exposure conditions used in calculating the guidelines for each land use scenario, it is possible to identify which land use scenario is protective for a particular municipal zoning requirement. Assessors must determine the full range of uses allowed under the applicable zoning bylaw when determining the appropriate land use for Tier 1 application. Where a municipal zoning decision incorporates more than one land use scenario, the most conservative land use must be applied. More detailed guidance on land use may be found in Section 3.2.

In some cases, a contaminated site may be adjacent to a property with a more sensitive land use. Even though contaminant concentrations may meet appropriate guidelines for the less sensitive land use, mobile substances may migrate to the adjacent property at concentrations that exceed guidelines for the more sensitive land use. This is a particular risk for the vapour inhalation pathway and the groundwater direct ecological contact pathway. When a contaminated site is bordered by a more sensitive land use, the vapour inhalation guidelines (soil and groundwater) and the groundwater direct ecological contact guideline for the more sensitive land use apply to the contaminated site anywhere within 30 m of the more sensitive property boundary (see Figure 1).

### **2.2.4 Relationship to Air and Water Quality**

Soil contamination interacts with air through volatilization and with water through dissolution and leaching to groundwater or runoff to surface water. Mobile soil contamination that is adding contaminant mass to air or water, is automatically considered a source. Therefore it must be remediated, or the contaminant release from the soil must be controlled as noted in Section 2.1.1.

For all land uses except natural areas, the Alberta Tier 1 soil remediation guidelines include soil concentrations that have been developed to protect indoor air quality. When soil contaminant concentrations exceed the soil remediation guideline protective of indoor air, then management of this exposure pathway is required.

The Alberta Tier 1 soil remediation guidelines also include soil concentrations that have been developed to protect groundwater quality. When soil contaminant concentrations exceed the soil remediation guideline protective of any groundwater-governed pathway (e.g., protection of potable groundwater for drinking water, protection of groundwater for freshwater aquatic life, or other groundwater pathways), then an investigation of groundwater quality is required. As part of this investigation, a Tier 2 approach may be considered. Tier 2 approaches could include pathway exclusion, site-specific risk assessments, or guideline adjustments based on separation distances between the zone of contamination and the seasonally high saturated zone or the distance to the water body of concern (see ESRD, 2007 and updates; Part B, Section 6).

A groundwater quality investigation to determine if groundwater has been impacted by soil contamination is also strongly recommended when contaminant concentrations in soil are close to the groundwater protection guidelines because the presence of preferential flow paths can result in contaminants reaching groundwater even when general soil conditions appear not to pose a risk. It is also possible that mobile substances have leached out of the vadose zone into groundwater if sufficient time has elapsed since the release event.

### 2.2.5 Relationship to Land Application of Organic Materials, Industrial By-Products and Pesticides

The *Industrial Release Limits Policy* (AENV, 2000) specifies that substances regulated by Alberta Environment and Parks must be managed to prevent soil contamination.

Alberta's approach to the management of wastes, industrial byproducts, composts, and other materials is based on the potential to improve soil quality. Wastes that provide no benefits to soil quality must not be applied to land in a manner that causes soil contamination. Industrial byproducts, composts, and other materials that provide a potential benefit to soil quality may be applied to land according to good agronomic or forestry practices and in accordance with any other regulatory requirements. Even when benefits can be shown, any potential contaminants in the byproduct must be managed to prevent their buildup in soil to concentrations that exceed Tier 1 guidelines.

Soil is a biologically active medium and is sometimes used as a treatment medium for soil contaminated by biodegradable substances. Land treatment of soil contaminated by gasoline, diesel fuel, jet fuel and kerosene is allowed if registered in accordance with the *Code of Practice for Land Treatment of Soil Containing Hydrocarbons* (AENV, 2008).

Tier 1 guidelines are used to evaluate chemical concentrations in soil. They can also be used to evaluate chemical concentrations resulting from the mixing of industrial by-products or organic materials into soil. They should not be used to evaluate concentrations in the by-product or organic material itself, unless the material will be placed directly on land without being mixed with soil.

Tier 1 pesticide guidelines have been developed for a limited number of exposure pathways for which sufficient information is available. These values are intended for use in the remediation of pesticide contaminated sites and not for restricting pesticide use in accordance with registered application rates.

Disposal and treatment of wastes generated by the upstream oil and gas industry are regulated by the Alberta Energy Regulatory in accordance with *Directive 50, Drilling Waste Management* (AER, 2019) and *Directive 58, Oilfield Waste Management Requirements for the Upstream Petroleum Industry* (AER, 2006). Directive 50 Equivalent Salinity Guidelines (AER, 2014a) can be used to assess and remediate drilling waste disposal areas for reclamation certification.

## 2.3 Groundwater Protection in Alberta

Groundwater and surface water are integrally connected. Therefore, groundwater cannot be managed in isolation from surface water and aquatic ecosystems. Emphasis is placed on preventing groundwater resources from becoming contaminated. However, where contamination of this public resource has resulted in an exceedance of Tier 1 or Tier 2 guidelines, it must be remediated or managed to ensure on-going protection of human health and the environment, and the restoration of beneficial uses.

This document provides guidance for managing contaminated groundwater in Alberta, and provides risk-based remediation guidelines to indicate when groundwater quality has been restored to an acceptable level.

### 2.3.1 Protection of Domestic Use Aquifers

Groundwater for domestic use is a significant current and future resource distributed over large geographic ranges in Alberta. Consequently, there is a need to protect the quality of Domestic Use Aquifers (DUAs).

The definition of a DUA is dependent on the amount of water an aquifer can produce, rather than the quality of the water in the aquifer, recognizing that technological treatment methods exist that can reduce or remove natural background substances. Furthermore, an aquifer does not have to be currently used for domestic purposes in order to be classified as a DUA, as the intent is to define and protect these aquifers for current and future use. Alberta Environment and Parks may consider any body of groundwater above the Base of Groundwater Protection<sup>1</sup> that is capable of a sufficient yield of water to be a DUA.

---

<sup>1</sup> The Base of Groundwater Protection is the depth above which groundwater is naturally non-saline, having a natural concentration of total dissolved solids that is less than or equal to 4000 milligrams per litre. Information on the Base of Groundwater Protection is available from the Alberta Geological Survey.

For the purpose of selecting and applying a groundwater guideline for human health protection by ingestion, a DUA is defined as a geologic unit (either of a single lithology or inter-bedded units) that is above the Base of Groundwater Protection having one or more of the following properties:

- A bulk hydraulic conductivity of  $1 \times 10^{-6}$  m/s or greater and sufficient thickness to support a sustained yield of 0.76 L/min or greater; or
- Is currently being used for domestic purposes; or
- Any aquifer determined by Alberta Environment to be a DUA.

While it is possible that peat deposits and muskeg may meet the definition of a DUA, based on hydraulic conductivity and unit thickness, Alberta Environment and Parks does not consider peat deposits or muskeg to be a DUA because groundwater in them is unlikely to be used as a domestic source.

The DUA drinking water pathway cannot be excluded under the Tier 1 approach. However, using Tier 2, it is possible to screen out the DUA drinking water pathway under certain circumstances, such as if an isolating geologic unit meeting specific properties is present. If a shallow large diameter well (or bored well) has been installed in a geologic unit that otherwise would not meet the definition of a DUA, the entire geological unit is not considered a DUA but the well must be treated as a point of compliance for the DUA pathway. For more information on the criteria for screening out the DUA drinking water pathway, refer to Tier 2.

### **2.3.2 Using Tier 1 Groundwater Remediation Guidelines**

The goal of the Alberta groundwater remediation guidelines is to provide numerical targets for the remediation of contaminated groundwater. To ensure consistency with “pollution prevention,” a key outcome of Alberta’s framework for contaminated sites management, the Tier 1 groundwater remediation guidelines are not “pollute-up-to” levels. Sources must not be left uncontrolled until cumulative releases result in an exceedance of Tier 1 or Tier 2 groundwater remediation guidelines. This results only in further contamination, increased remediation costs, and potential loss of water use options. Source control is a crucial component of pollution prevention.

### **2.3.3 Background Groundwater Quality**

The background concentration of a substance in groundwater is the natural concentration of that substance in a particular groundwater zone in the absence of any input from anthropogenic activities or sources. In some situations, the background concentration of some substances can be a significant proportion of, or even exceed the Tier 1 remediation guideline. Remediation of groundwater to below background conditions is not feasible and is not required. Accordingly, it is important to have a good understanding of background groundwater conditions at a site.

Background concentrations will be specific to the groundwater zone being considered, and will vary both spatially and temporally. To gain a good understanding of background conditions at a site it is necessary to consider groundwater quality data from several monitoring wells installed in the zone of interest, located up- or cross- gradient from any contaminant sources. The more time-series data that are available, the better the understanding of background conditions will be.

Care should be taken to distinguish between apparent background concentrations that are the result of diffuse anthropogenic sources, and true, natural background conditions. In comparing against background, emphasis should always be placed on ensuring that anthropogenic sources are not identified as natural background.

### **2.3.4 Relationship to Soil, Air and Surface Water**

Environmental media are interconnected. Contaminants in soil may leach into pore water or groundwater. Volatile compounds in groundwater may volatilize at the water table and can migrate through the soil into the interior space of buildings above. Soluble contaminants in groundwater can be transported laterally with the groundwater flow, and potentially enter a surface water body (creek, slough, lake, etc.) at the point of groundwater discharge.

Alberta Tier 1 groundwater remediation guidelines are developed to protect indoor air quality, plants and soil invertebrates, and water quality for a range of uses. Guidelines to protect a particular water use are calculated based on the corresponding water quality guideline (drinking water, aquatic life, irrigation, or livestock or wildlife watering).

### **2.3.5 Points of Compliance**

For the purpose of this document, a point of compliance is the spatial location in an aquifer at which a groundwater quality guideline must be achieved to protect human and ecological receptors, to protect a groundwater resource, or to meet other conditions such as industrial use or groundwater management guidelines.

At one extreme, the compliance point could be established at the point of exposure such as a drinking water well (human-ingestion) or a river (ecological-aquatic life). However, this would imply that there could be deterioration in the quality of the groundwater between the contaminant source and the receptor, which could be judged unacceptable in terms of legislative requirements and/or restriction of potential future use of a groundwater resource. At the other extreme, a precautionary approach could set the groundwater compliance point directly beneath the contaminant source. This is likely to result in a more stringent remedial target concentration and may be unnecessary, as certain exposure pathways may be irrelevant at that particular location, the contaminated groundwater may never reach a receptor, or contaminants may be significantly attenuated in groundwater prior to reaching the exposure point.

To address both conditions, some fundamental principles are used to guide decisions for setting groundwater remediation guidelines and compliance points at individual sites in Alberta.

A DUA is an important current and future groundwater resource and must be protected to the maximum extent possible. The compliance point for the human health water ingestion pathway is everywhere within a DUA.

Groundwater aquifers can be an important current or potential future agricultural groundwater resource used for livestock watering and irrigation. For livestock watering, the compliance point is everywhere within the relevant livestock watering aquifer existing below agricultural or other grazing land. For irrigation, the compliance point is everywhere within the irrigation-use aquifer, where applicable.

Ecological receptors must be protected at key exposure points. For aquatic life or wildlife receptors, the minimum point of compliance is at the point of groundwater discharge into a surface water body that is capable of supporting an aquatic ecosystem. Groundwater guidelines are calculated to achieve this. Therefore, the groundwater at all points of groundwater discharge immediately adjacent to the aquatic water body must meet the aquatic surface water quality guideline. For terrestrial receptors (plants and soil invertebrates), the point of compliance is everywhere within the shallow groundwater zone (i.e. the extent of groundwater less than 3 m below ground surface) and at the point of ground surface discharge.

## 3. Alberta Framework for the Management of Contaminated Sites

### 3.1 Implementation Framework and Management Levels

The general framework for the management of contaminated sites in Alberta has three options and is illustrated by the flowchart presented in Figure 2. A more detailed framework specific to site management under the Tier 1 approach is presented in Figure 3. A brief description of the framework is provided below; a detailed discussion of the management and technical aspects of the Tier 1 guidelines is presented in subsequent sections of the document.

Under the Alberta framework (ESRD, 2014c), three options are provided for the management of contaminated sites as the proponent proceeds from initial site assessment to regulatory closure. The three options are:

- Tier 1 - generic remediation guidelines.
- Tier 2 - site-specific remediation guidelines based on the modification of Tier 1 guidelines.
- Exposure Control - risk management through exposure barriers or administrative controls based on site-specific risk assessment.

Regardless of the option chosen, the target level of human health and ecological protection afforded by Tier 1, Tier 2, or Exposure Control is the same.

As discussed below, regulatory closure is available for sites managed to achieve Tier 1 and Tier 2 remediation guidelines. This means that no conditions are imposed on future use of the site, within a given land use. Regulatory closure is not provided for Exposure Control. For more description on the 3 options and how they relate to each other, see the *Contaminated Sites Policy Framework* (ESRD, 2014c).

### 3.2 Land Use Definitions

For the purpose of developing and implementing soil and groundwater remediation guidelines in Alberta, five generic land uses have been defined – natural areas, agricultural, residential/ parkland, commercial, and industrial. A generic land use scenario is envisioned for each category based on typical activities on these lands. The five land uses are defined for the purpose of this document only. Where allowable land uses, as defined by a given jurisdictional authority, differ from those noted here, an assessment of allowable receptors and potential exposure pathways must be made to ensure consistency with assumptions based on definitions here. Where a more sensitive receptor or exposure pathway is allowed, the more sensitive land use description must be used in selecting the appropriate Tier 1 guidelines. Assessors must determine the full range of uses allowed under the applicable zoning bylaw when determining the appropriate land use for Tier 1 application. For most sites in Alberta, one of the five generic land use scenarios should be applicable. If none of the generic land uses are applicable, a site-specific Tier 2 approach will be required. The five land uses are defined as follows:

#### **Natural Areas**

Natural areas are defined as being away from human habitation and activities, where the primary concern is the protection of ecological receptors. Accordingly, human exposure pathways are not considered, with the exception of the protection of groundwater for drinking water pathway that, based on the definition of a DUA, applies in all land uses. Much of Alberta's forested land will fall into natural areas land use. Forested lands that are specified as grazing leases represent a special case that requires an amendment to the normal exposure scenario for natural areas. On such grazing leases, the livestock soil and food ingestion and protection of groundwater for livestock pathways must be addressed in addition to the regular pathways considered under natural areas land use. Natural areas land use must not be applied to areas that may reasonably be expected to be developed, such as those near municipalities and permanent dwellings.

#### **Agricultural Lands**

On agricultural land the primary land use is growing crops or tending livestock as well as human residence. This also includes agricultural lands that provide habitat for resident and transitory wildlife and native flora. To allow unrestricted future use of the land, a farm residence is assumed to be present anywhere on agricultural land.

## **Residential/Parkland**

The primary activity on residential/parkland is residential or recreational activity. This land use includes campground areas and urban parks, but not wildlands in provincial parks, which are considered natural areas. Where urban parks are frequented by wildlife, wildlife exposure pathways should be addressed.

## **Commercial Land Use**

On commercial land, the primary activity is commercial (e.g., shopping mall) and all members of the public, including children, have unrestricted access. Commercial land use includes day-care centres, buildings for religious services, hospitals, and medical centres. Commercial land does not include operations where food is grown directly in impacted soil on the site. Such operations would fall under agricultural land use.

## **Industrial Land Use**

Industrial land is land where the primary activity is the production, manufacture or construction of goods. Public access is restricted and children are not permitted continuous access or occupancy.

### **3.3 Groundwater and Surface Water Use Definitions**

Soils are hydrologically linked to groundwater and surface water systems. One of the objectives of Alberta's soil remediation guidelines is to manage soil-to-groundwater pathways to prevent unacceptable transfer of contaminants from the soil, which may ultimately affect groundwater and surface water use. Alberta's groundwater and surface water quality guidelines are representative of allowable chemical concentrations in groundwater or surface water at the point of compliance (see Section 2.3.5).

Alberta guidelines have been developed for four generic uses of groundwater or surface water affected by groundwater discharge:

- human consumption (potable water);
- aquatic life;
- livestock and wildlife watering; and,
- irrigation

These water uses are linked to land uses through the definitions of the generic land use categories. Other water use categories, for example recreation, or variations in water use within a defined land use category, may be addressed using the Tier 2 process.

### **3.4 Conditions and Restrictions Associated with Tier 2 and Exposure Control**

The Alberta soil and groundwater remediation guidelines and implementation framework are intended to provide the same high level of protection of human health and the environment at all levels or tiers of site management. For Tier 1, this is ensured by the use of relatively conservative assumptions in the derivation of the risk-based numerical guidelines, such that the values can be applied to the large majority of sites within a land use category without condition or restriction. In other words, the normal activities associated with a particular land use are protected without the need for ongoing management of the site or of contaminants which may be present. Alberta Environment and Parks will provide regulatory closure for a site complying with the Tier 1 guidelines, unless site conditions are unsuitable for their application (see Section 5).

Management under Tier 2 guidelines delivers the same level of health and environmental protection by incorporating site-specific data into the development of appropriate remediation guidelines. Sites remediated to Tier 2 guidelines are eligible for regulatory closure.

Certain types of site-specific data or assumptions dictate the need for ongoing site management to ensure that the assumptions used to assess human and ecological risks or to develop site-specific objectives remain valid. Ongoing management of a site, or of the contaminants present, will generally invoke a land use restriction or condition that will preclude regulatory closure. Therefore site-specific adjustments or assumptions that would trigger ongoing management requirements can only be implemented under the Exposure Control option.

To avoid the need for ongoing management and, hence, possible conditions and land use restrictions, Tier 2 adjustments are limited to parameters that are measurable and stable, such as certain soil properties, geological conditions, hydrogeology and distances to natural surface water bodies.

Tier 2 assessments that involve full site-specific risk assessment using models and assumptions that may differ from those used in the calculation of the Tier 1 guidelines may be accepted provided they do not require any form of ongoing risk management. Parameters that are unique to current site use, an existing development or the location of a receptor, such as the characteristics of a site building or the distance to a water well, may change in the future thereby invalidating the site-specific assumptions. Adjustments of such parameters are not allowed at Tier 2. In some cases exposure pathways may be inoperative under a particular site use (e.g. direct human or ecological contact with contaminated soil at a commercial site that is paved or capped) or the frequency of exposure may differ from the generic assumptions. Preservation of these conditions would require ongoing management; therefore these adjustments cannot be made at Tier 2. Further guidance on parameters and assumptions eligible for adjustment at Tier 2 is provided in ESRD (2007 and updates).



## 4. Scientific Basis for Developing Tier 1 Remediation Guidelines

### 4.1 Risk-Based Guidelines

Risk-based guidelines are developed with the understanding that for risk from chemical contamination to exist, all of the following elements must be present:

- the substance must possess toxic properties;
- the substance must be present in the environmental medium of concern (soil or groundwater);
- a receptor must be present (human, livestock, crop, or ecological receptor); and,
- there must be an exposure pathway through which the substance can reach the receptor and be taken in to the receptor's body.

Risk-based guidelines are then calculated by first determining the amount (dose or concentration) of substance to which a receptor can be safely exposed. Next, for each exposure pathway, a conservative estimate is made of a concentration of the substance in soil or groundwater that will protect the receptor from exposure exceeding the safe amount. These substance concentrations are referred to as risk-based guidelines and represent remediation objectives for the protection of human and ecological health on contaminated sites.

#### 4.1.1 Steps in the Development of Tier 1 Risk-Based Guidelines

The steps in the development of the risk-based guidelines in Tables 1, 2 and 3 are as follows.

- Identification of the potential receptors and the potential exposure pathways through which contaminants can come into contact with receptors (e.g., humans or ecological receptors such as livestock, crops, and wildlife).
- Identification of appropriate protection endpoints for each exposure pathway/receptor combination.
- Calculation of a substance concentration in soil and/or groundwater (guideline) for each exposure pathway that offers a safe level of protection.
- Determination of an overall Tier 1 risk-based guideline for soil and/or groundwater by selecting the lowest of the guidelines calculated for all relevant exposure pathways.

The remainder of this section describes how these four steps were implemented in the development of the Tier 1 soil and groundwater remediation guidelines. The details of the calculations used in Step 3 are provided in Appendix C.

### 4.2 Protection of Human Health

#### 4.2.1 Human Receptors and Exposure Pathways

In establishing appropriate risk-based guidelines, the most sensitive user of a contaminated site must be considered and protected. The most sensitive receptor is normally a function of the degree of potential exposure, the exposure pathway(s) and the substance(s) of concern. For the application of Tier 1 and Tier 2 guidelines, the land use and contaminant characteristics will dictate the critical receptor. When developing Tier 1 and Tier 2 guidelines, people of all ages are assumed to be present on agricultural, residential/parkland and commercial land. At industrial sites, only employees are assumed to be routinely present, which precludes the exposure of children. Human receptors are assumed to be absent in natural areas, although underlying groundwater is considered to be a potential source of drinking water.

The following human exposure pathways are considered when developing and implementing Tier 1 and Tier 2 remediation guidelines (based on CCME, 2006). The exposure pathways are applicable to all land uses, except where noted below.

##### **Direct Contact**

Humans coming into direct contact with contaminated soil via incidental ingestion, dermal contact, or inhalation of air-borne soil particles. Applicable to all land uses except natural areas.

##### **Drinking Water**

Humans drinking from and showering or bathing in water that is sourced from groundwater. Applicable to all land uses.

## **Inhalation**

Volatile contaminants being released from soil and/or groundwater and migrating upwards into living or working spaces where humans are exposed via inhalation. Applicable to all land uses except natural areas.

## **Off-Site Surface Migration by Wind or Water Erosion**

The soil quality guideline for commercial and industrial land use may be greater than the corresponding guideline for more sensitive land uses. Wind or water transport of contaminated soil from a commercial or industrial site onto an adjacent site with a more sensitive land use could potentially result in contaminant concentrations that exceed the human direct contact soil quality guideline applicable to the more sensitive land use. The off-site migration check is calculated to ensure that the commercial or industrial guidelines set are protective of this exposure pathway.

### **4.2.2 Human Health Protection Endpoints**

The human health protection endpoint is the same at all tiers of management, and is expressed in terms of an allowable exposure level at which the likelihood of a receptor experiencing adverse health effects is essentially negligible. Specifically, the level of human exposure to a threshold chemical must not exceed the tolerable daily intake specified by Health Canada or other appropriate regulatory agency, including background exposure to the chemical. For a non-threshold carcinogenic chemical, the incremental lifetime risk must not exceed 1 in 100,000 ( $1 \times 10^{-5}$ ), the value considered by Health Canada to be essentially negligible (Health Canada, 2004).

Human health protection endpoints that have been accepted for use in Alberta are listed in Appendix C. For more information on acceptable endpoints for use in Alberta, see the *Guidance for Selecting Toxicity Reference Values for Alberta Tier 1 and Tier 2 Soil and Groundwater Remediation Guidelines* (AEP, 2017).

## **4.3 Ecological Protection**

### **4.3.1 Ecological Receptors and Exposure Pathways**

Risk-based guidelines fulfill two main goals from the ecological standpoint: protection of ecological receptors expected to be present at a site based on land use, and preservation of an appropriate level of ecological function of the site and its ecosystem components.

Ecological receptors at a typical contaminated site, within the range of generic land uses considered in the development of the Alberta guidelines, span a range of trophic levels including soil-dependent organisms (plants, including crops, and soil invertebrates) and higher order consumers (terrestrial and avian wildlife and livestock). In addition, based on the potential for groundwater underlying a site to discharge to a surface water body that is capable of supporting an aquatic ecosystem, aquatic receptors including invertebrates, fish, and aquatic plants are considered. Receptors assigned to each land use for the purpose of guideline derivation must be both ecologically relevant to a site and sufficiently sensitive to be representative of the range of receptors normally present. In addition to the protection of ecological receptors *per se*, ecologically based guidelines must be protective of other processes such as nutrient cycling and related microbial activities.

The following ecological exposure pathways are considered in the determination and implementation of Tier 1 remediation guidelines where appropriate to the defined land uses (based on CCME, 2006).

### **Direct Contact**

Plants and soil invertebrates coming into direct contact with contaminants in soil or shallow groundwater. Direct contact is applicable to all land uses. This pathway may be eliminated below 3 m for petroleum hydrocarbon fractions 1 to 4 only.

### **Nutrient and Energy Cycling**

This exposure pathway examines the microbial functioning of the soil, including carbon and nitrogen cycling. Applicable to all land uses.

### **Livestock/Wildlife Soil and Food Ingestion**

Livestock or wildlife ingesting contaminants via the incidental ingestion of soil and ingesting contaminants that have bioaccumulated from soil into fodder. Applicable to agricultural and natural area land use only.

## Aquatic Life

Aquatic life, including fish, aquatic invertebrates and aquatic plants, being exposed to contaminants when groundwater discharges to a surface water body that is capable of supporting an aquatic ecosystem. Applicable to all land uses.

## Irrigation

Crops being exposed to contaminants when groundwater is used for irrigation. Applicable to agricultural land use only.

## Livestock/Wildlife Watering

Livestock or wildlife being exposed to contaminants when groundwater is used for livestock watering, or groundwater discharges to a surface water body where wildlife may drink. Applicable to agricultural and natural area land use only.

## Off-Site Surface Migration by Wind or Water Erosion

The soil quality guideline for commercial and industrial land use may be greater than the corresponding guideline for more sensitive land uses. Wind or water transport of contaminated soil from a commercial or industrial site onto an adjacent site with a more sensitive land use could potentially result in contaminant concentrations that exceed the ecological direct contact soil quality guideline applicable to the more sensitive land use. The off-site migration check is calculated to ensure that the commercial or industrial guidelines set are protective of this exposure pathway.

## 4.4 Calculation of Guidelines

Soil remediation guidelines were calculated for each land use and soil texture using models consistent with the latest CCME (2006) protocols. In some cases, parameter values have been updated based on more recent publications or adapted for Alberta conditions and policies where appropriate.

Groundwater remediation guidelines were calculated for each water use by adapting the models from the soil remediation guideline calculations, and CCME (2015) as appropriate.

Details of all models and parameters to be used in Alberta are provided in Appendix C. Details of where parameters or models may differ from CCME are listed in the *Contaminated Sites Policy Framework* (ESRD, 2014c and updates).

## 4.5 Determination of Overall Guideline

Tier 1 guidelines are intended to be a conservative screening tool. The lowest of the guidelines (tabulated in Appendix A and B) calculated for applicable exposure pathways and the appropriate land use and soil texture is the overall Tier 1 remediation guideline (Tables 1, 2, and 3).

## 4.6 Integration of New and Existing Guidelines

New risk-based guidelines were calculated for a wide range of organic compounds in this document using the latest CCME (2006) protocols (in some cases modified based on the 2007 revisions to the PHC CWS) and adapted for Alberta conditions and policies (Tables 1, 2, and 3). Guidelines were not recalculated for inorganic compounds, except arsenic and boron; existing risk-based guidelines were retained from *Canadian Environmental Quality Guidelines* (CCME, 1999 and updates). Existing guidelines for the soil ecological contact pathway and the soil and food ingestion pathway were also retained where they existed, or were recalculated in the case of soil ingestion guidelines for certain hydrocarbons. Soil remediation guidelines based on professional judgment were retained from previous guideline documents for antimony, beryllium, cobalt, fluoride, molybdenum, silver, elemental sulphur, and tin because risk-based Alberta Tier 1 guidelines have not yet been developed for these substances. Guidelines for these substances were adopted from *the Alberta Tier 1 Criteria for Contaminated Soil Assessment and Remediation* (AEP, 1994) for natural areas, agricultural, and residential/parkland land uses. For commercial and industrial land uses, and for all uses if the substance was not included in AEP (1994), guidelines were adopted from the *Interim Canadian Environmental Quality Criteria for Contaminated Sites* (CCME, 1991). Soil remediation guidelines for electrical conductivity and sodium adsorption ratio (Table 4) were adopted from the *Salt Contamination Assessment and Remediation Guidelines* (AENV 2001c).

## 5. User Guidance for Tier 1 Management

The process for implementing the Alberta Tier 1 Soil and Groundwater Remediation Guidelines is illustrated schematically by the flow diagrams presented in Figures 2 and 3. Detailed guidance for Tier 2 is provided in a companion Tier 2 document (ESRD, 2007 and updates).

As illustrated in Figures 2 and 3, Tier 1 management is divided into three stages: assessment, evaluation, and decision/management. These three stages are discussed in the following sections.

### 5.1 Information Requirements for Tier 1 (Tier 1 Assessment)

Site assessment requirements are documented in the *Alberta Environmental Site Assessment Standard* (AEP, 2016a). A preliminary site characterization is required for all Tier 1 assessments. Site characterization must be comprehensive enough to adequately describe site conditions and address all assessment and management options within the scope of Tier 1.

The Tier 1 assessment may be more comprehensive and detailed, at the discretion of the proponent, providing additional information for potential use in developing Tier 2 guidelines. While there are advantages associated with a phased approach to site assessment, there may also be economies in combining data collection activities into a single investigation, particularly at locations where mobilization and demobilization costs are significant.

The minimum data requirements for a Tier 1 site assessment include:

#### Land Use and Sensitivity Factors

- site description;
- land use, including the full range of uses allowed under the applicable zoning bylaw;
- proximity of the site to surface water and drinking water supplies;
- actual and potential uses of groundwater;
- human receptors; and,
- ecological receptors.

#### Physical Conditions

- soil particle size;
- stratigraphy and properties of surficial materials;
- depth to groundwater; and,
- presence and types of buildings and other structures.

#### Contaminant Characteristics and Distribution

- contamination characterization; and,
- horizontal and vertical extent of contamination.

The above information is required at all sites, to a level of detail appropriate for the requirements of the evaluation and decision stages at each site. Application of the Tier 2 protocol will require additional information to support the decision at that Tier.

#### 5.1.1 Land Use and Sensitivity Factors

One of the objectives of the Tier 1 site assessment is to determine whether the site broadly fits any of the five generic land use categories. In this regard, the minimum data requirements for a Tier 1 assessment are as follows:

##### Site Description

The site description must include basic identifying data such as location and legal description, site dimensions, registered owner(s) etc., as well as a description of the physical surface expression of the site (surface topography), vegetative cover, nature of site development and, where applicable, site history.

## Land Use

In determining land use, consideration must be given not only to present land use but also to historic and potential future land uses. Land use on private land is normally prescribed by the current and/or future zoning of the land and local development trends. Land use for public land is determined by the land manager who will use resource based input and tools such as land reservations and integrated plans as guidance. Attention must be given to all uses allowed under the applicable zoning bylaws and land use decisions.

Sufficient descriptive and legal or administrative information must be obtained to determine the generic land use category (natural areas, agricultural, residential/parkland, commercial, industrial) and whether the land use may be considered typical for the assigned category. The five generic land uses are defined and described in Section 3.2.

Land uses of adjacent or nearby properties must also be documented, as the presence of more sensitive off-site land uses may drive the site management requirements. Where a contaminated site on a less sensitive land use borders, or lies within 30m, of a more sensitive land use, the vapour inhalation guidelines (soil and groundwater) and the groundwater direct ecological contact guidelines for the more sensitive land use must be applied to the contaminated site within 30 m of the more sensitive land use boundary (see Figure 1).

## Proximity of Site to Surface Water and Drinking Water Supplies

The assessment must document all existing and potential uses of groundwater and surface water, including their locations. The distance beyond which water uses are not significant or relevant depends on site-specific conditions but, as a minimum for Tier 1, water uses within 300 m should be identified. If the direction of groundwater flow has been reliably determined by site-specific groundwater monitoring, documentation of water uses can be limited to 100 m upgradient and 300 m down gradient of the site. Flood risk areas should be taken into account when considering distance to surface water receptors. For more information see the companion Tier 2 document (ESRD, 2007 and updates).

The distance of 300 m was selected as being greater than the length of the majority of groundwater plumes of dissolved petroleum hydrocarbon. This value is based on studies of hydrocarbon plumes in the upstream oil and gas industry in western Canada undertaken by the Consortium of Research on Natural Attenuation (CAPP, 2002). The value of 300 m is also consistent with a compilation of 647 petroleum hydrocarbon plumes presented in Wiedemeier *et al.* (1999), in which it was determined that 98.1% of the plumes were less than 900 feet (274 m) long.

Water uses at greater distances must be considered on a case-by-case basis when subsurface conditions are such that adverse impact may be possible. For example, plumes of conservative soluble compounds that do not interact with the soil matrix, such as chloride, may travel farther than dissolved hydrocarbons (CAPP, 2002). Municipal water supply wells located more than 300 m from a site might be expected to be at risk from a contaminant source located anywhere within the zone of capture of the wells.

The assessment must recognize that, while the locations of most natural surface water bodies are essentially fixed, seasonal water bodies may exist at other locations, and anthropogenic groundwater uses could be initiated at any location relative to the site, in areas of usable groundwater.

## Human and Ecological Receptors

In general, the presence of human receptors will be directly related to the land uses. However, sufficient information must be obtained to determine whether the land use is typical of the respective category, or whether receptors are present that would warrant a variation from the defined generic land use categories.

### 5.1.2 Physical Conditions

Certain physical conditions and parameters must be determined to permit the implementation of Tier 1 management. The minimum physical data requirements are discussed below.

#### Soil Particle Size

As a minimum, since Tier 1 guidelines are prescribed for coarse-grained and fine-grained soils, sufficient particle size information should be obtained to permit classification of the soils as either coarse or fine. Fine-grained soils are defined as having 50% or more by mass pass a 75 µm sieve; coarse-grained soils have less than 50% by mass pass a 75 µm sieve. The absence of sufficient particle size information will result in the default application of the more conservative Tier 1 guideline for

each substance. Where both fine and coarse grained strata are present, the dominant soil particle size is determined by the stratum governing horizontal and vertical migration to a receptor.

Particle size must be determined on the soil fraction that passes a 2mm sieve, unless the fraction retained on the sieve exceeds 70% (w/w) of the sample. If the fraction retained on the sieve exceeds 70% (w/w), the sample must be classified as coarse and consideration must be given to whether the site is dominated by very coarse materials as described in Section 5.1.5.

### **Soil Stratigraphy and Physical Properties**

Information on soil stratigraphy and physical properties is normally obtained by means of an intrusive subsurface investigation, although preliminary, qualitative information can often be obtained from other sources including published surficial geological information or the results of other subsurface investigations conducted in the area. In certain circumstances, it may be possible to apply the Tier 1 guidelines without a prior subsurface investigation. For example, the decision could be made to manage a spill of limited extent by removing all materials containing contaminant concentrations in excess of Tier 1 guidelines. Confirmation of the degree and extent of impact, as well as the subsurface conditions, could be obtained at the time of soil excavation. However, subsurface investigations are commonly conducted to support the application of Tier 1 guidelines, in part for the purpose of contaminant characterization and site management planning.

The Tier 1 assessment must provide an overall description of the subsurface soil conditions and their vertical and lateral variability. Of particular importance is the uniformity of the soil particle size and the presence of any depositional or structural features, such as lenses or fissures, that could influence the fate and transport of certain chemicals in the subsurface. Where there are multiple geological deposits falling into both the coarse and fine definition such that the particle size dominating transport cannot be determined, the Tier 1 guidelines must default to the most stringent option.

### **Hydrogeological Conditions**

A hydrogeological investigation will normally form part of any subsurface investigation. Although soil guidelines have been developed to protect groundwater for various uses, care must be taken in using them to assess risk to groundwater at contaminated sites. Soils are heterogeneous. Variations in contaminant concentration and soil lithology and the presence of preferential flow paths can result in contaminants reaching groundwater even when general soil conditions appear not to pose a risk. It is also possible that mobile substances have leached out of the vadose zone into groundwater if sufficient time has elapsed since the release event. Alberta Environment and Parks strongly recommends that groundwater quality be investigated when contaminant concentrations in soil are close to the groundwater protection guidelines. The cost of installing groundwater monitoring wells can be minimized if they are installed when soil boreholes are drilled.

For some substances, the soil guidelines are not clearly linked to groundwater guidelines. For instance, soil guidelines for metals do not consider the groundwater component. In these instances, a groundwater investigation is required unless it can be clearly demonstrated that the substance has not migrated to the groundwater.

Information must be collected on depth to groundwater table, groundwater flow direction, hydraulic gradient and, with the appropriate field tests, hydraulic conductivity. While all of these parameters are influential, their quantitative determination is not required for the basic implementation of Tier 1, and qualitative information may be available from other sources. However, as with soil parameters, they can facilitate the decision between managing to Tier 1 guidelines or developing Tier 2 remediation guidelines or Exposure Control options. For some substances, groundwater chemical characteristics are required for the development of a groundwater guideline. Where this applies, Table 2 references the *Environmental Quality Guidelines for Alberta Surface Waters* (ESRD, 2014a and updates) for further information on the chemical data required.

### **Buildings and Other Structures**

When volatile substances are present, the site assessment must identify presence of buildings. When a building is not present, but the municipal zoning allows one to be built, its presence must be assumed. Site assessments must also consider the presence of structures, such as utility corridors and earthen basements, that may require a Tier 2 approach.

#### **5.1.3 Contaminant Characteristics and Distribution**

Requirements for site characterization to address all aspects for regulatory requirements or to characterize contaminants that are not included in the Tier 1 guideline process are beyond the scope of this document. It is the responsibility of the proponent to ensure that sufficient site characterization is carried out to address relevant regulatory requirements and to address any potential contaminants of concern not included in the Tier 1 soil and groundwater remediation guidelines. However, for the purposes of applying the Tier 1 guideline tables, the following general guidance can be given.

## Analytical Considerations

Analytical requirements for soil and groundwater analysis are provided in *Guidance Manual for Environmental Site Characterization in Support of Environmental and Human Health Risk Assessment, Volume 4 Analytical Methods* (CCME, 2016), with the exception of the following parameters:

Soil boron, soil electrical conductivity, and soil soluble cations for the calculation of sodium adsorption ratio must be analyzed in a saturated paste extract prepared in accordance with Method 15.2.1 in *Soil Sampling and Methods of Analysis* (Carter and Gregorich, 2008).

Barite-barium in soil must be analyzed by fusion-XRF or fusion-ICP. For more information see *Soil Remediation Guidelines for Barite: Environmental and Human Health* (AENV, 2009).

Monoethanolamine and diethanolamine must be analyzed in accordance with *Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine* (AENV, 2010), or equivalent.

Elemental sulphur in soil must be analyzed using methods specified in *Directive for Monitoring the Impact of Sulphur Dust on Soils* (AEP, 2015).

For petroleum hydrocarbons, polycyclic aromatic hydrocarbons are not subtracted from Fraction 3 and naphthalene is not subtracted from Fraction 2.

Naturally occurring hydrocarbons in materials such as peat or compost may cause false petroleum hydrocarbon guideline exceedances. The *Reference Method for the Canada-Wide Standard for Petroleum Hydrocarbons in Soil* (CCME, 2001) provides options for addressing the presence of biogenic hydrocarbons. The *Biogenic Interference Calculation (BIC) Scale* (AEP, 2018) may also be used to identify false guideline exceedances. AEP (2018) provides additional guidance and requirements for using the BIC Scale.

## Site Characterization

Sufficient characterization of the contamination must be conducted for the purpose of comparing contaminant concentrations with Tier 1 guidelines. This will normally comprise the collection and laboratory analysis of sufficient soil or groundwater samples to determine representative concentrations. As part of the Tier 1 assessment, the proponent must conduct sufficient sampling to delineate the lateral and vertical extent of impacted soil and groundwater (Fig. 3). An exception is allowed where the contaminated soil is completely removed and confirmatory samples show compliance with Tier 1 guidelines. When confirmatory samples fail to comply with Tier 1 guidelines after excavation is deemed complete, full delineation of the remaining contamination must be undertaken and used to develop further remediation actions or risk management programs. Whether delineating contamination or confirming remediation success, the data collected must be sufficient to support regulatory closure.

When delineating chloride-based soil salinity for comparison to Tier 1 soil salinity guidelines (Table 4), delineation must include the lateral and vertical extent of chloride concentrations in saturated paste extracts (expressed as mg/L) that exceed the Tier 1 groundwater guideline. The area and depth defined by soil chloride delineation requirements may extend beyond the area and depth of soil that exceeds Tier 1 soil salinity guidelines. In that case, soil remediation may be based on the Tier 1 soil salinity guidelines, but consideration must be given to the potential for chloride contamination in groundwater.

### 5.1.4 Conditions Where Additional Exposure Pathways May Be Applicable

There may be conditions where certain exposure pathways are applicable that would not normally be considered under a given land use. An example of this is a grazing lease within a natural area. In the case of such grazing leases, the livestock soil and food ingestion and protection of groundwater for livestock watering pathways apply in addition to the regular exposure pathways considered for that land use. Adding additional pathways to those typically considered for a land use may be done under the Tier 1 approach, provided no additional site information is needed for the additional pathway. Note that removing pathways that may not be applicable for a given land use from the pathways specified for the Tier 1 guidelines is not permitted, and may only be done under the Tier 2 approach.

### 5.1.5 Conditions Where Tier 1 Guidelines Are Not Applicable

Tier 1 guidelines are derived using relatively conservative default parameters and assumptions corresponding to the defined generic exposure scenarios and land use categories. As such, they are intended and expected to be protective of human health and the environment in the large majority of cases. However, there may be situations in which Tier 1 guidelines are not applicable, either where conditions violate one or more assumptions essential to the validity of the modelling used in the Tier 1 derivation, or where actual exposure conditions or receptors at a site are more sensitive than in the generic exposure scenario.

Some examples of situations where Tier 1 guidelines are not applicable are highlighted below. These situations must be addressed through the Tier 2 process. The companion Tier 2 document should be consulted for further details.

#### Source of Volatile Contaminants Present Within 30 cm of a Building Foundation

Within this distance, the model used to assess vapour intrusion to buildings is not reliable, and Tier 1 guidelines are not applicable.

#### Unusual Structural Features

Building features such as earthen floors or unusually low air exchange rates are not considered in the Tier 1 model, and a Tier 2 approach is required. Where very coarse textured materials (see below) are found within utility corridors that are controlling contaminant migration, a Tier 2 approach is required.

#### Groundwater Flow to Stagnant Water Bodies

Additional consideration is required if groundwater at a site has the potential to discharge to a stagnant water body. A stagnant water body is defined as a water body without significant outflow, and where the main pathway of water loss is via evaporation. Stagnant water bodies will tend to concentrate discharging persistent groundwater contaminants through evaporation. Water bodies with no obvious or known outflow should be considered stagnant. If outflow is suspected via groundwater and no obvious surface outflow is present, a groundwater investigation will be needed to provide confirmation.

In the assessment of whether soil or groundwater contaminants are likely to have an adverse effect on a stagnant surface water body, the current concentrations of contaminants in groundwater are less important than the long-term effect on contaminant concentrations in the stagnant water body. Accordingly, when there is the potential for a contaminant in groundwater to discharge to a stagnant surface water body, a Tier 2 mass balance assessment of the likely concentrations of that contaminant in the stagnant water body over the anticipated lifetime of the groundwater discharge is required. The assessment should take into consideration, in a qualitative sense, the likelihood of other potential future contaminant releases to the stagnant surface water body. Unless the effect on contaminant concentrations in the stagnant surface water body can be shown to be insignificant, remedial action will be required.

#### Groundwater Within 10 m of a Surface Water Body

Tier 1 soil and groundwater remediation guidelines for the protection of aquatic life assume a minimum separation of 10 m between the point that the soil or groundwater concentration is measured, and the discharge point. Accordingly, Tier 1 guidelines only apply to soil or groundwater located at least 10 m from the nearest surface water body that is capable of supporting an aquatic ecosystem. Within this distance, a Tier 2 approach is required, or in the case of groundwater guidelines, the corresponding surface water freshwater aquatic life guideline may be applied directly to groundwater quality. The 10 m offset distance must take into account potential for seasonal fluctuations in the water level. This may require information on the flood risk area. For more information, see the companion Tier 2 document.

#### Very Coarse Textured Materials Enhancing Groundwater or Vapour Transport

Groundwater velocity is a function of both hydraulic conductivity and hydraulic gradient and assumed to be  $1 \times 10^{-5}$  m/s and 0.028 m/m, respectively, for Tier 1 guideline derivation. The resulting Darcy groundwater velocity is  $3 \times 10^{-7}$  m/s.

The rate of vapour transport through coarse soil is largely governed by vapour permeability which is assumed to be  $6 \times 10^{-8}$  cm<sup>2</sup> for Tier 1 guideline derivation.

If the Darcy groundwater velocity exceeds  $3 \times 10^{-7}$  m/s, or vapour permeability is greater than  $6 \times 10^{-8}$  cm<sup>2</sup> at a site, a Tier 2 approach is required. In these cases, the Tier 2 guideline must incorporate groundwater or vapour inhalation as potentially dominant exposure pathway(s).



## Fractured Bedrock

The Tier 1 remediation guidelines were developed for unconsolidated soil material, therefore the presence of bedrock may require a Tier 2 re-evaluation. However, the guidelines may be applied to contaminants in contact with bedrock if the bedrock is likely to behave conservatively as one of the two soil textures. The actual texture of the bedrock material will often be less influential on contaminant movement than the degree of bedrock weathering and fracturing. Professional judgment must be applied in determining whether coarse or fine soil guidelines are the most appropriate, given the expected contaminant mobility within the bedrock. For instance, a weathered shale material may not automatically require a Tier 2 re-evaluation but it may require use of a coarse soil guideline due to the presence of minor fractures within the matrix.

Tier 2 re-evaluation is required where flow paths in the fractured bedrock cannot reasonably be expected to behave similarly to those in an aggregated soil medium. When the fracture length exceeds approximately 2 cm, flow paths in the fractured bedrock may be different than those in an aggregated soil medium. Under such conditions, groundwater transport in fractured bedrock is outside the scope of the calculations for Tier 1 guidelines and a Tier 2 or Exposure Control approach must be taken.

## Source Length Greater than 10 m

The model used to develop Tier 1 guidelines for groundwater remediation and protection assumes a source of groundwater contamination that is 3m deep and 10m wide with a length of 10 m parallel to the direction of groundwater flow. Sources are discussed in Section 2.1.1 and include anything that adds contaminant mass to the environment. For the purposes of the groundwater transport model used to develop Tier 1 guidelines, the source length is relevant to sources that are releasing contaminants to groundwater. Source length is determined by soil concentrations that exceed soil guidelines for the protection of groundwater pathways (see Appendix A). When the length of such a source is greater than 10 m parallel to the direction of groundwater flow, Tier 1 guidelines may be applied to the site if contaminant delineation shows that the source volume is less than 300 m<sup>3</sup>. Alternatively, if source and site characteristics indicate no potential for groundwater contamination and remediation of the site to Tier 1 guidelines will result in a source length less than 10 m or volume less than 300 m<sup>3</sup>, then remediation to Tier 1 objectives may proceed. In all other cases a Tier 2 approach is required. When the contaminated soil is not in contact with the seasonally high saturated zone, a Tier 2 guideline adjustment (ESRD, 2007 and updates, Part B, Section 6) provides the assessor with an approach for estimating contaminant attenuation between the contaminated zone and the saturated zone.

## Organic Soils

When inorganic contaminants occur in organic soils, a Tier 2 approach is required. Organic soils are defined in *The Canadian System of Soil Classification* (Soil Classification Working Group, 1998). Coarse textured soil and groundwater remediation guidelines may be used for organic contaminants in organic soil.

## 5.2 Identification of Appropriate Tier 1 Guidelines (Tier 1 Evaluation)

### 5.2.1 Land Use and Water Use Conditions and Primary Soil Type

Land use and water use conditions are established on the basis of the information compiled for the Tier 1 assessment. Land use conditions are compared with the descriptions of the generic land use categories (natural area, agricultural, residential/parkland, commercial, industrial) to assign the site to the most sensitive applicable category. In some instances municipalities or public land managers may allow a range of uses in accordance with zoning bylaws and other land use decisions. These uses must be evaluated with respect to the five land uses described for Tier 1 guidelines. The most sensitive land use will determine Tier 1 guidelines.

Groundwater and surface water use conditions are assigned, if applicable, to one or more of the generic categories of potable water, surface water that is sustaining an aquatic ecosystem, water used for livestock or wildlife watering, and irrigation.

Exceptions to the above categories are discussed in the following section.

### 5.2.2 Conformity with Generic Land and Water Use Categories

The principal criterion for the application of the Tier 1 guidelines is conformity with the generic land and water use scenarios and exposure conditions assumed in the development of the Tier 1 guidelines, and discussed in Section 3. It is the responsibility of the proponent to identify and respond to any site or receptor factors that could unduly accentuate exposure or risk beyond that envisioned in the Tier 1 exposure scenarios and, where necessary, move to the Tier 2 approach.

First, the land and water use must normally be encompassed by one or more of the defined generic categories. Examples of land uses not addressed in the development of the Tier 1 guidelines include wetlands and riparian zones. Examples of water uses not directly addressed by Tier 1 include: water used in the food processing industry (although this may be represented by the potable water scenario); and surface water used for swimming or similar recreational activity. In the absence of an applicable Tier 1 standard for the actual land or water use, a Tier 2 approach must be followed.

Second, the exposure conditions associated with the identified land and water use must not be more sensitive or critical than those assumed in the determination of the Tier 1 guidelines. Examples of factors giving rise to greater sensitivity are: a greater frequency or intensity of human or ecological exposure beyond that associated with typical use of the land or water; or, variations in physical site conditions resulting in greater exposure than that assumed (discussed in Section 5.1.5). The human and ecological exposure factors and physical parameters used for the Tier 1 guidelines are presented in Appendix C. These values must be used as the basis for assessing whether site-specific factors are indicative of more frequent or intense exposure.

Ingestion of produce, milk, and meat produced on the site is not included in Tier 1 human exposure estimates because of its site-specific nature. Where a significant portion of an individual's diet is obtained from areas contaminated by substances that bioconcentrate, this exposure pathway must be evaluated with the Tier 2 approach.

### 5.2.3 Identification of Applicable Tier 1 Guidelines

Tier 1 guidelines for soil and groundwater are presented in Tables 1 through 4. Guidelines in Tables 1 through 3 are presented for coarse-grained and fine-grained soils; selection of the appropriate set of values is based on the texture of the dominant soil type as determined in the Tier 1 assessment. The dominant soil type is that which governs fate and transport via the various transport and exposure pathways. For example, a continuous layer of coarse-grained soil beneath the water table will often govern groundwater flow in the saturated zone, even though its thickness may be small in relation to the total thickness of saturated fine-grained soils. Similarly, a thin layer of fine-grained soil in the unsaturated zone may be a more significant control on the migration of vapours than a thicker layer of coarse-grained soil. Professional judgment should be exercised in establishing the dominant soil type.

In general, Alberta Tier 1 soil remediation guidelines (Table 1) apply to all soil regardless of depth. However, subsoil guidelines for petroleum hydrocarbons (Table 3) and salinity (Table 4) may be used at Tier 1 under the following circumstances:

#### Petroleum Hydrocarbons

Surface soil guidelines for petroleum hydrocarbon fractions F1 to F4, benzene, toluene, ethylbenzene, and xylenes must be applied to a minimum depth of 1.5 m. Subsurface guidelines may be used as follows:

- below 3 metres in depth at any site (in this case, surface soil guidelines must be applied to a depth of 3 metres),
- below 1.5 metres in depth within a 5 metre setback from an oilfield wellhead (see Directive 079, *Surface Development in Proximity to Abandoned Wells* (AER, 2014b)), or
- below 1.5 m at remote forested sites in the Green Area with fine textured soil irrespective of the distance to a wellhead. In order to use subsoil guidelines outside of a 5 metre radius from the wellhead, sites must meet the requirements in the *Subsoil Petroleum Hydrocarbon Guidelines for Remote Forested Sites in the Green Area* (ESRD, 2014b).

The Tier 1 approach allows the exclusion of the ecological direct contact pathway for petroleum hydrocarbon fractions F1 to F4 for any land use below a depth of 3 metres. All other exposure pathways apply. Consistent with the *Canada-Wide Standard for Petroleum Hydrocarbons in Soil* (CCME, 2008), exclusion of the ecological direct soil contact pathway is permitted below 3 metres with the application of "management limits" for petroleum hydrocarbon fractions that represent maximum concentrations that apply at any depth. Management limits for petroleum hydrocarbon fractions F1 to F4 are provided in Appendix A. For sites that meet the requirements of the *Subsoil Petroleum Hydrocarbon Guidelines for Remote Forested Sites in the Green Area* (ESRD, 2014b) management limits provided in ESRD (2014b) may be applied.

#### Salinity

Topsoil guidelines for electrical conductivity and sodium adsorption ratio must be applied to the L, F, H, O, and A horizons (Soil Classification Working Group, 1998) or equivalent surficial material where these horizons are not present. Subsoil guidelines may be applied below the A horizon or equivalent in lieu of topsoil guidelines. Further information is available in the *Salt Contamination Assessment and Remediation Guidelines* (AENV, 2001c).

For each contaminant, the lowest guideline tabulated in Appendix A and B for the established soil type and identified land use defines the governing exposure pathway/receptor and, hence, becomes the governing Tier 1 guideline. At Tier 1, all exposure

pathways for which generic guidelines have been calculated must be considered. If the governing guideline is based on a soil to groundwater or surface water pathway that is not applicable to the site, based on the prior identification of water uses, a Tier 2 approach may be possible, and the companion Tier 2 document should be consulted.

#### **5.2.4 Additional Guidance for Specific Substances**

##### **Salinity**

Electrical conductivity (EC) and sodium adsorption ratio (SAR) guidelines are found in Table 4 for topsoil and subsoil. At Tier 1, the objective for salt contaminated site remediation is to return the site to the same rating category as non-contaminated soils of the same type. To apply the guidelines, background samples are needed from uncontaminated soils that are representative of the same type, depth, and landscape position as the contaminated soil on the site. The background samples are used to establish the appropriate rating categories for the site. The range of EC and SAR values for the appropriate rating categories become the remediation objectives for the site. Further guidance is provided in the *Salt Contamination Assessment and Remediation Guidelines* (AENV, 2001c).

Additional considerations are needed for assessing and remediating salt-impacted wellsites in native grasslands, which are sensitive to disturbance. Elevated levels of salinity may be left in place where the native plant community has not been impacted by existing salinity and risks of future impacts due to salt migration are minimal. Direction for obtaining the information needed to support leaving salinity in place is found in the *Native Prairie Protocol for Reclamation Certification of Salt-Affected Wellsites* (AEP, 2019).

##### **Barite-Barium**

Guidelines are provided for both barium (non-barite) and barite-barium. The barium guidelines are intended to apply to all barium sources, and therefore assume that the source is relatively soluble. Barite, a commonly used weighting agent in drilling fluids, is relatively insoluble and therefore guidelines have been developed specifically for this form of barium. To confirm that barium measured in soil samples meets the conditions of low solubility assumed for the barite-barium guidelines, the sample must pass a  $\text{CaCl}_2$  extraction test. True total barium concentrations from samples that pass the  $\text{CaCl}_2$  extraction test may be compared to the barite-barium guidelines. True total barium must be measured by fusion-XRF or fusion-ICP methods. Further information on applying the barite-barium guidelines, including the  $\text{CaCl}_2$  test, is provided in *Soil Remediation Guidelines for Barite: Environmental and Human Health* (AENV, 2009).

##### **Elemental Sulphur**

Tier 1 soil remediation guidelines are provided for elemental sulphur. If these values are exceeded, management options include the application of calcium carbonate to control the acidity generated by the oxidation of elemental sulphur. Elemental sulphur can be managed in this way up to a total sulphur concentration of 4%. Further guidance is provided in *Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils* (AENV, 2011).

##### **Drilling Waste Disposal Areas**

Drilling waste disposal falls under the jurisdiction of the Alberta Energy Regulator. Management and disposal options are provided by *Directive 50* (AER, 2019). Alberta Environment and Parks has developed D50 Equivalent Salinity Guidelines for assessing and remediating drilling waste disposal areas for reclamation certification. Further information is provided in *Assessing Drilling Waste Disposal Areas: Compliance Options for Reclamation Certification* (AER, 2014a).

##### **Polycyclic Aromatic Hydrocarbons (PAH)**

Carcinogenic PAHs are assessed as a group for human health purposes. For soil, the Index of Additive Cancer Risk (IACR) and the Toxic Potency Equivalent (TPE) must be calculated using the formulas provided in Table 1. For groundwater, Toxic Potency Equivalent (TPE) is calculated for carcinogenic PAHs using the calculation provided in Table 2. Some individual PAHs have guidelines for ecological pathways. Where these are provided in Tables 1 and 2, the concentration of those individual PAHs must be compared to the appropriate guideline as well as comparing the entire carcinogenic PAH group to the IACR and TPE for soil or the TPE for groundwater.

Non-carcinogenic PAHs (acenaphthene, anthracene, fluoranthene, fluorene, naphthalene, phenanthrene, pyrene) have been included as components of petroleum hydrocarbon Fractions 1 to 4. When the sources of hydrocarbon contamination are crude petroleum or refined fuels, separate analysis of non-carcinogenic PAHs is not necessary, unless the contaminants are in a Domestic Use Aquifer or a surface water aquatic life receptor. Analysis of these PAHs is necessary when dealing with non-petroleum hydrocarbons such as coal tar, creosote, hydrocarbons altered by incomplete combustion (e.g., flare pits, petroleum

or petrochemical fires) or products such as solvents where individual compounds may be present at a much greater concentration than in crude products or fuels.

## 5.3 Tier 1 Decision and Management

The Tier 1 decision process is illustrated schematically by the lower part of the flow diagram presented in Figure 3. The process consists of a number of decision nodes. The decision process and criteria at each node are discussed in the following paragraphs.

### 5.3.1 Comparison of Conditions with Identified Tier 1 Guidelines

If no exceedances are found in the Tier 1 assessment, relative to the identified Tier 1 guidelines, any sources have been removed, and site conditions are consistent with application of Tier 1, the site can be considered to be in compliance with Tier 1, and no further action is necessary.

If exceedances are found, the proponent enters a decision process whereby the option of managing to Tier 1 guidelines is weighed against the option of proceeding to the Tier 2 approaches. If site conditions preclude application of Tier 1 guidelines (see Section 5.1.5) then the proponent must use the Tier 2 process.

Verification of remediation to Tier 1 or Tier 2 guidelines can achieve regulatory closure.

### 5.3.2 Assessment of Opportunity for Tier 2

If the default assumptions used in the derivation of the governing Tier 1 guidelines are conservative relative to actual site-specific conditions, the replacement of default assumptions with site-specific data for certain influential parameters may permit the development and implementation of less stringent remediation guidelines without compromising health and environmental protection goals. Furthermore, point-of-exposure measurements, if less than those predicted using the relatively conservative modeling procedures employed to derive the Tier 1 guidelines, may also permit the implementation of less stringent guidelines. In such cases, proceeding to Tier 2 would generally be advantageous, and the companion Tier 2 document should be consulted.

## 6. References

Alberta Environment and Parks references are available at:

<https://www.alberta.ca/ministry-environment-parks.aspx>.

- AENV (Alberta Environment), 2000. Industrial Release Limits Policy. Environmental Sciences Division. Alberta Environment Publication # T/569.
- AENV (Alberta Environment), 2001a. Alberta Soil and Water Quality Guidelines for Hydrocarbons at Upstream Oil and Gas Facilities. Environmental Sciences Division, September 2001.
- AENV (Alberta Environment), 2001b. Risk Management Guidelines for Petroleum Storage Tank Sites. Science and Standards Division, Pub No. T/570, October 2001.
- AENV (Alberta Environment), 2001c. Salt Contamination Assessment and Remediation Guidelines. Environmental Sciences Division, Pub No. T/606, May 2001.
- AENV (Alberta Environment), 2008. Code of Practice for Land Treatment of Soil Containing Hydrocarbons. Alberta Queen's Printer.
- AENV (Alberta Environment), 2009. Soil Remediation Guidelines for Barite: Environmental Health and Human Health. Environmental Assurance Division. Alberta Environment.
- AENV (Alberta Environment), 2010. Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine. Environmental Assurance Division. Alberta Environment.
- AENV (Alberta Environment), 2011. Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils. Science and Technology Division. Alberta Environment.
- AEP (Alberta Environmental Protection), 1994. Alberta Tier I Criteria for Contaminated Soil Assessment and Remediation, March 1994. Pub. No. T/475.
- AEP (Alberta Environment and Parks), 2015. Directive for Monitoring the Impact of Sulphur Dust on Soils. Land Policy Branch, Policy and Planning Division.
- AEP (Alberta Environment and Parks), 2016a. Alberta Environmental Site Assessment Standard.
- AEP (Alberta Environment and Parks), 2016b. Alberta Exposure Control Guide.
- AEP (Alberta Environment and Parks), 2018. BIC Scale for Delineating Petroleum Hydrocarbons in Organic Soils and Compost. AEP Land Policy 2018-1.
- AEP (Alberta Environment and Parks), 2019. Native Prairie Protocol for Reclamation Certification of Salt-Affected Wellsites. Land Policy and Programs 2019-3.
- AER (Alberta Energy Regulator), 2006. Oilfield Waste Management Requirements for the Upstream Petroleum Industry. Directive 58.
- AER (Alberta Energy Regulator), 2019. Drilling Waste Management. Directive 50.
- AER (Alberta Energy Regulator), 2014a. Assessing Drilling Waste Disposal Areas: Compliance Options for Reclamation Certification.
- AER (Alberta Energy Regulator), 2014b. Surface Development in Proximity to Abandoned Wells, Directive 079.
- CAPP (Canadian Association of Petroleum Producers), 2002. Assessment of Monitored Natural Attenuation at Upstream Oil & Gas Facilities in Alberta – Final Report. March 2002. Canadian Association of Petroleum Producers, No. 2001-0010.
- Carter, M.R. and E.G. Gregorich (eds.), 2008. Soil Sampling and Methods of Analysis, 2<sup>nd</sup> edition. Canadian Society of Soil Science. CRC Press, Boca Raton.
- CCME (Canadian Council of Ministers of the Environment), 1991. Interim Canadian Environmental Quality Criteria for Contaminated Sites. Canadian Council of Ministers of the Environment, Winnipeg. Report CCME EPC-CS34, September 1991.
- CCME (Canadian Council of Ministers of the Environment), 1999 and updates. Canadian Environmental Quality Guidelines. Canadian Council of Ministers of the Environment, Winnipeg.

- CCME (Canadian Council of Ministers of the Environment), 2001. Reference Method for the Canada-Wide Standard for Petroleum Hydrocarbons in Soil - Tier 1 Method. PN 1310. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2006. A Protocol for the Derivation of Environmental and Human Health Soil Quality Guidelines. PN 1332. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2008. Canada-Wide Standards for Petroleum Hydrocarbons (PHCs) in Soil: Scientific Rationale.
- CCME (Canadian Council of Ministers of the Environment), 2014. A Protocol for the Derivation of Soil Vapour Quality Guidelines for Protection of Human Exposures Via Inhalation of Vapours, PN 1531. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2015. A Protocol for the Derivation of Groundwater Quality Guidelines for Use at Contaminated Sites. PN 1533. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2016. Guidance Manual for Environmental Site Characterization in Support of Environmental and Human Health Risk Assessment, Volume 4 Analytical Methods. PN 1557. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- ESRD (Alberta Environment and Sustainable Resource Development), 2007 as amended. Alberta Tier 2 Soil and Groundwater Remediation Guideline. Land and Forestry Policy Branch, Policy Division.
- ESRD (Alberta Environment and Sustainable Resource Development), 2014a and updates. Environmental Quality Guidelines for Alberta Surface Waters. Water Policy Branch, Policy Division.
- ESRD (Alberta Environment and Sustainable Resource Development), 2014b. Subsoil Petroleum Hydrocarbon Guidelines for Remote Forested Sites in the Green Area. Land Policy Branch, Policy and Planning Division.
- ESRD (Alberta Environment and Sustainable Resource Development), 2014c. Contaminated Sites Policy Framework. Land and Forestry Policy Branch, Policy Division.
- Health Canada, 2004. Federal Contaminated Site Risk Assessment in Canada. Part I: Guidance on Human Health Preliminary Quantitative Risk Assessment (PQRA). Prepared by: Environmental Health Assessment Services Safe Environments Programme, September 2004.
- Health Canada, 2013. Canadian Guidelines for the Management of Naturally Occurring Radioactive Materials (NORM). Minister of Public Works and Government Services Canada, Cat. H46-1/30-2000E.
- MEMS (Millenium EMS Solutions Ltd.), 2016. Disinfection By-product Chloroform in Shallow Groundwater Within the City of Calgary. Prepared for the City of Calgary.
- Soil Classification Working Group, 1998. The Canadian System of Soil Classification, 3<sup>rd</sup> edition. Research Branch, Agriculture and Agri-Food Canada, Publication 1646. NRC Research Press, Ottawa.
- Stantec Consulting Ltd. 2012. Ecotoxicity Assessment of a Soil Sterilant – Bromacil. Prepared for Cenovus Energy. 227 pp.
- Stantec Consulting Ltd. 2008. Ecotoxicity Evaluation in Support of the Derivation of Tier 2 Values for Tebuthiuron. Prepared for EBA Engineering Consultants. 93 pp.
- Wiedemeier, T.H., H.S. Rifai, C.J. Newell, and J.T. Wilson, 1999. Natural Attenuation of Fuels and Chlorinated Solvents in the Subsurface. John Wiley and Sons, New York.

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
General and Inorganic Parameters											
pH (in 0.01M CaCl <sub>2</sub> )	6.0-8.5	6.0-8.5	6.0-8.5	6.0-8.5	6.0-8.5	6.0-8.5	6.0-8.5	6.0-8.5	6-8.5	6.0-8.5	1
Cyanide (free)	0.90	0.90	0.90	8.0	8.0	0.90	0.90	0.90	8.0	8.0	2
Fluoride	200	200	200	2,000	2,000	200	200	200	2,000	2,000	1
Sulphur (elemental)	500	500	500	500	500	500	500	500	500	500	3
Metals											
Antimony	20	20	20	40	40	20	20	20	40	40	1
Arsenic (inorganic)	17	17	17	26	26	17	17	17	26	26	
Barium (non-barite)	750	750	500	2,000	2,000	750	750	500	2,000	2,000	2
Barite-barium	10,000	10,000	10,000	15,000	140,000	10,000	10,000	10,000	15,000	140,000	4
Beryllium	5.0	5.0	5.0	8.0	8.0	5.0	5.0	5.0	8.0	8.0	1
Boron (mg/L in saturated paste extract)	3.3	3.3	3.3	5.0	5.0	3.3	3.3	3.3	5.0	5.0	15
Cadmium	3.8	1.4	10	22	22	3.8	1.4	10	22	22	2
Chromium (hexavalent)	0.40	0.40	0.40	1.4	1.4	0.40	0.40	0.40	1.4	1.4	2
Chromium (total)	64	64	64	87	87	64	64	64	87	87	2
Cobalt	20	20	20	300	300	20	20	20	300	300	1
Copper	63	63	63	91	91	63	63	63	91	91	2
Lead	70	70	140	260	600	70	70	140	260	600	2
Mercury (inorganic)	12	6.6	6.6	24	50	12	6.6	6.6	24	50	2
Molybdenum	4.0	4.0	4.0	40	40	4.0	4.0	4.0	40	40	1
Nickel	45	45	45	89	89	45	45	45	89	89	2
Selenium	1.0	1.0	1.0	2.9	2.9	1.0	1.0	1.0	2.9	2.9	2
Silver	20	20	20	40	40	20	20	20	40	40	1
Thallium	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	2
Tin	5.0	5.0	5.0	300	300	5.0	5.0	5.0	300	300	1
Uranium	33	23	23	33	300	33	23	23	33	300	
Vanadium	130	130	130	130	130	130	130	130	130	130	2
Zinc	250	250	250	410	410	250	250	250	410	410	2

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Hydrocarbons											
Benzene	0.046	0.046	0.046	0.046	0.046	0.078	0.015	0.015	0.078	0.078	5
Toluene	0.52	0.52	0.52	0.52	0.52	0.12	0.12	0.12	0.12	0.12	5
Ethylbenzene	0.073	0.073	0.073	0.073	0.073	0.14	0.14	0.14	0.14	0.14	5
Xylenes	0.99	0.99	0.99	0.99	0.99	1.9	1.9	1.9	1.9	1.9	5
Styrene	0.68	0.68	0.68	0.68	0.68	0.80	0.80	0.80	0.80	0.80	
F1	210	210	210	320	320	210	24	24	270	270	6
F2	150	150	150	260	260	150	130	130	260	260	6
F3	1,300	1,300	1,300	2,500	2,500	300	300	300	1,700	1,700	6
F4	5,600	5,600	5,600	6,600	6,600	2,800	2,800	2,800	3,300	3,300	6
Acenaphthene	0.33	0.33	0.33	0.33	0.33	0.38	0.38	0.38	0.38	0.38	
Anthracene	1.3	1.3	1.3	1.3	1.3	0.0056	0.0056	0.0056	0.0056	0.0056	
Fluoranthene	15.4	15.4	50	180	180	0.055	0.055	0.055	0.055	0.055	
Fluorene	0.40	0.40	0.40	0.40	0.40	0.34	0.34	0.34	0.34	0.34	
Naphthalene	0.014	0.014	0.014	0.014	0.014	0.017	0.017	0.017	0.017	0.017	
Phenanthrene	0.11	0.11	0.11	0.11	0.11	0.061	0.061	0.061	0.061	0.061	
Pyrene	7.7	7.7	2,100	3,200	30,000	0.15	0.15	0.15	0.15	0.15	
Carcinogenic PAHs	IACR<1.0	IACR<1.0 and TPE ≤ 5.6	IACR<1.0 and TPE ≤ 5.6	IACR<1.0 and TPE ≤ 8.5	IACR<1.0 and TPE ≤ 23	IACR<1.0	IACR<1.0 and TPE ≤ 5.6	IACR<1.0 and TPE ≤ 5.6	IACR<1.0 and TPE ≤ 8.5	IACR<1.0 and TPE ≤ 23	7, 17, 18
Benz[a]anthracene	6.2	6.2	-	-	-	6.2	6.2	-	-	-	8
Benzo[b+j]fluoranthene	6.2	6.2	-	-	-	6.2	6.2	-	-	-	8
Benzo[k]fluoranthene	6.2	6.2	-	-	-	6.2	6.2	-	-	-	8
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	-	-	-	
Benzo[a]pyrene	0.60	0.60	20	72	72	0.60	0.60	20	72	72	8
Chrysene	6.2	6.2	-	-	-	6.2	6.2	-	-	-	8
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	-	-	-	
Halogenated Aliphatics											
Vinyl chloride	0.014	0.0083	0.0083	0.014	0.014	0.020	0.00034	0.00034	0.0043	0.0043	
1,1-Dichloroethene	0.15	0.15	0.15	0.15	0.15	0.24	0.021	0.021	0.24	0.24	



TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Trichloroethene (Trichloroethylene, TCE)	0.054	0.0043	0.0043	0.030	0.030	0.081	0.00019	0.00019	0.0023	0.0023	5, 11
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.26	0.26	0.26	0.26	0.26	0.46	0.018	0.018	0.22	0.22	
1,2-Dichloroethane	0.025	0.0062	0.025	0.025	0.025	0.041	0.0027	0.0027	0.033	0.033	
Dichloromethane (Methylene chloride)	0.10	0.052	0.10	0.10	0.10	0.095	0.048	0.095	0.095	0.095	
Trichloromethane (Chloroform)	0.16	0.16	0.16	0.16	0.16	0.030	0.030	0.030	0.030	0.030	
Tetrachloromethane (Carbon tetrachloride)	0.037	0.022	0.035	0.037	0.037	0.062	0.0015	0.0015	0.018	0.018	
Dibromochloromethane	0.91	0.12	0.91	0.91	0.91	1.5	0.12	0.27	1.5	1.5	
Chlorinated Aromatics											
Chlorobenzene	0.61	0.39	0.39	0.61	0.61	1.1	0.018	0.018	0.22	0.22	12
1,2-Dichlorobenzene	0.097	0.097	0.097	0.097	0.097	0.18	0.18	0.18	0.18	0.18	12
1,4-Dichlorobenzene	0.051	0.051	0.051	0.051	0.051	0.098	0.098	0.098	0.098	0.098	
1,2,3-Trichlorobenzene	0.26	0.26	0.26	0.26	0.26	0.31	0.26	0.26	0.31	0.31	
1,2,4-Trichlorobenzene	0.78	0.78	0.78	0.78	0.78	0.93	0.23	0.23	0.93	0.93	
1,3,5-Trichlorobenzene	1.9	1.9	1.9	1.9	1.9	3.6	0.13	0.13	1.3	1.3	
1,2,3,4-Tetrachlorobenzene	0.042	0.042	0.042	0.042	0.042	0.050	0.050	0.050	0.050	0.050	
1,2,3,5-Tetrachlorobenzene	0.37	0.37	0.37	0.37	0.37	0.70	0.10	0.10	0.70	0.70	
1,2,4,5-Tetrachlorobenzene	0.19	0.19	0.19	0.19	0.19	0.37	0.052	0.052	0.37	0.37	
Pentachlorobenzene	24	22	22	24	24	5.2	5.2	5.2	5.2	5.2	
Hexachlorobenzene	1.8	0.80	1.8	1.8	1.8	3.6	0.20	0.20	2.3	2.3	
2,4-Dichlorophenol	0.0029	0.0029	0.0029	0.0029	0.0029	0.0034	0.0034	0.0034	0.0034	0.0034	
2,4,6-Trichlorophenol	0.19	0.19	0.19	0.19	0.19	0.37	0.37	0.37	0.37	0.37	
2,3,4,6-Tetrachlorophenol	0.039	0.039	0.039	0.039	0.039	0.047	0.047	0.047	0.047	0.047	
Pentachlorophenol	0.025	0.025	0.025	0.025	0.025	0.029	0.029	0.029	0.029	0.029	5
Dioxins & Furans	0.00025	0.000004	0.000004	0.000004	0.000004	0.00025	0.000004	0.000004	0.000004	0.000004	9
PCBs	1.3	0.085	0.085	0.13	1.2	1.3	0.085	0.085	0.13	1.2	5
Pesticides											
Aldicarb	0.041	0.012	0.041	0.041	0.041	0.065	0.012	0.065	0.065	0.065	12
Aldrin	0.24	0.24	0.24	0.24	0.24	0.46	0.31	0.31	0.46	0.46	

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Atrazine and metabolites	0.0088	0.0088	0.0088	0.0088	0.0088	0.010	0.010	0.010	0.010	0.010	
Azniphos-methyl (Guthion)	0.41	0.41	0.41	0.41	0.41	0.75	0.75	0.75	0.75	0.75	
Bromacil	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	0.009	13
Bromoxynil	0.044	0.044	0.044	0.044	0.044	0.052	0.052	0.052	0.052	0.052	13
Carbaryl	1.9	1.9	1.9	1.9	1.9	3.6	3.6	3.6	3.6	3.6	12
Carbofuran	0.68	0.082	0.68	0.68	0.68	1.2	0.089	1.2	1.2	1.2	12
Chlorothalonil	0.0085	0.0085	0.0085	0.0085	0.0085	0.010	0.010	0.010	0.010	0.010	
Chlorpyrifos	49	3.2	49	49	49	95	3.8	95	95	95	12
2,4-D	0.43	0.10	0.43	0.43	0.43	0.67	0.10	0.67	0.67	0.67	12
DDT	0.70	0.70	11	12	12	0.70	0.70	11	12	12	5
Diazinon	2.2	2.2	2.2	2.2	2.2	4.2	4.2	4.2	4.2	4.2	12
Dicamba	0.50	0.12	0.50	0.50	0.50	0.79	0.12	0.79	0.79	0.79	12,13
Diclofop-methyl	NGR	0.079	40	61	570	2.4	0.095	2.4	2.4	2.4	
Dieldrin	0.025	0.025	0.025	0.025	0.025	0.048	0.048	0.048	0.048	0.048	
Dimethoate	0.0058	0.0028	0.0058	0.0058	0.0058	0.0055	0.0027	0.0055	0.0055	0.0055	
Dinoseb	2.8	1.4	2.8	2.8	2.8	5.5	1.7	5.5	5.5	5.5	12
Diquat	11	11	11	11	11	21	21	21	21	21	
Diuron	1.9	1.9	1.9	1.9	1.9	3.5	3.5	3.5	3.5	3.5	
Endosulfan	0.80	0.80	0.80	0.80	0.80	0.0016	0.0016	0.0016	0.0016	0.0016	
Endrin	2.4	2.4	2.4	2.4	2.4	4.7	4.7	4.7	4.7	4.7	
Glyphosate	0.054	0.054	0.054	0.054	0.054	0.049	0.049	0.049	0.049	0.049	
Heptachlor epoxide	0.039	0.039	0.039	0.039	0.039	0.076	0.010	0.010	0.076	0.076	
Lindane	0.31	0.11	0.31	0.31	0.31	0.60	0.13	0.60	0.60	0.60	12
Linuron	0.051	0.051	0.051	0.051	0.051	0.059	0.059	0.059	0.059	0.059	13
Malathion	0.82	0.82	0.82	0.82	0.82	1.3	1.3	1.3	1.3	1.3	12
MCPA	0.42	0.026	0.42	0.42	0.42	0.66	0.025	0.66	0.66	0.66	12,13
Methoxychlor	NGR	110	110	170	800	0.32	0.32	0.32	0.32	0.32	
Metolachlor	0.048	0.048	0.048	0.048	0.048	0.055	0.055	0.055	0.055	0.055	
Metribuzin	0.024	0.012	0.024	0.024	0.024	0.028	0.014	0.028	0.028	0.028	
Paraquat (as dichloride)	1.1	1.1	1.1	1.1	1.1	2.2	2.2	2.2	2.2	2.2	
Phorate	0.075	0.075	0.075	0.075	0.075	0.14	0.14	0.14	0.14	0.14	

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Picloram	0.024	0.024	0.024	0.024	0.024	0.022	0.022	0.022	0.022	0.022	
Simazine	0.033	0.033	0.033	0.033	0.033	0.038	0.038	0.038	0.038	0.038	13
Tebuthiuron	0.046	0.046	0.046	0.60	0.60	0.046	0.046	0.046	0.60	0.60	12,13
Terbufos	0.080	0.080	0.080	0.080	0.080	0.15	0.15	0.15	0.15	0.15	
Toxaphene	3.3	3.3	3.3	3.3	3.3	6.3	4.8	4.8	6.3	6.3	
Triallate	0.0077	0.0077	0.0077	0.0077	0.0077	0.0092	0.0092	0.0092	0.0092	0.0092	
Trifluralin	0.22	0.22	0.22	0.22	0.22	0.045	0.045	0.045	0.045	0.045	
Other Organics											
Aniline	0.36	0.36	0.36	0.36	0.36	0.60	0.60	0.60	0.60	0.60	12
Di- <i>n</i> -butyl phthalate	0.54	0.54	0.54	0.54	0.54	0.65	0.65	0.65	0.65	0.65	
Dichlorobenzidine	0.60	0.60	0.60	0.60	0.60	1.2	1.2	1.2	1.2	1.2	
Diethanolamine	2.0	2.0	2.0	2.0	2.0	3.5	3.5	3.5	3.5	3.5	14
Diethylene glycol	10	10	10	10	10	15	15	15	15	15	
Diisopropanolamine	14	14	14	14	14	17	17	17	17	17	5
Ethylene glycol	60	60	60	60	60	62	62	62	62	62	5
Hexachlorobutadiene	0.026	0.026	0.026	0.026	0.026	0.031	0.0067	0.0067	0.031	0.031	
Methanol	37	37	37	37	37	11	11	11	11	11	
Methylmethacrylate	36	36	36	36	36	49	1.4	1.4	17	17	
Monoethanolamine	20	20	20	20	20	10	10	10	10	10	14
MTBE	0.044	0.044	0.044	0.044	0.044	0.062	0.046	0.046	0.062	0.062	
Nonylphenol + ethoxylates	5.7	5.7	5.7	14	14	5.7	5.7	5.7	14	14	5
Perfluooctane sulfonate (PFOS)	0.010	0.010	0.070	0.070	0.070	0.010	0.010	0.14	0.14	0.14	
Phenol	0.0028	0.0014	0.0028	0.0028	0.0028	0.0024	0.0012	0.0024	0.0024	0.0024	5, 16
Sulfolane	0.18	0.18	0.18	0.18	0.18	0.21	0.21	0.21	0.21	0.21	5
Triethylene glycol	100	100	100	100	100	150	150	150	150	150	
Radionuclides											
Uranium-238 Series (all progeny)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Uranium-238 ( <sup>238</sup> U, <sup>234</sup> Th, <sup>234</sup> mPa, <sup>234</sup> U)	10	10	10	10	10	10	10	10	10	10	10
Thorium-230	10	10	10	10	10	10	10	10	10	10	10

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Radium-226 (in equilibrium with its progeny)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Lead-210 (in equilibrium with <sup>210</sup> Bi and <sup>210</sup> Po)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Thorium-232 Series (all progeny)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Thorium-232	10	10	10	10	10	10	10	10	10	10	10
Radium-228 (in equilibrium with <sup>228</sup> Ac)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Thorium-228 (in equilibrium with its progeny)	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	10
Potassium-40	17	17	17	17	17	17	17	17	17	17	10

Notes:

Guideline values calculated for this document using latest available CCME protocols except where noted.

1. Value adopted from AEP (1994) and/or CCME (1991)
2. Value adopted from CCME (1999, as amended)
3. For more information see Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils (AENV, 2011)
4. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)
5. Ecological direct contact values from CCME (1999, as amended), other values calculated in this document
6. Ecological direct contact values from CCME (2007), other values calculated in this document
7. Carcinogenic PAH concentrations must meet the Index of Additive Cancer Risk ( IACR) <1 guideline. Individual PAH compounds must also meet guidelines for ecological receptors where specified in Table 1 with footnote 8. The IACR is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture, as follows:

Fine Soil:

$$IACR = \frac{[Benz(a)anthracene]}{6.4} + \frac{[Benzo(b + j)fluoranthene]}{3.0} + \frac{[Benzo(k)fluoranthene]}{0.64} + \frac{[Benzo(g, h, i)perylene]}{130} + \frac{[Benzo(a)pyrene]}{7.0} + \frac{[Chrysene]}{40} + \frac{[Dibenz(a, h)anthracene]}{4.4} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{51}$$

Coarse Soil:

$$IACR = \frac{[Benz(a)anthracene]}{12} + \frac{[Benzo(b + j)fluoranthene]}{5.8} + \frac{[Benzo(k)fluoranthene]}{1.2} + \frac{[Benzo(g, h, i)perylene]}{250} + \frac{[Benzo(a)pyrene]}{14} + \frac{[Chrysene]}{78} + \frac{[Dibenz(a, h)anthracene]}{8.5} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{98}$$

8. For ecological receptors only
9. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-TCDD (See CCME, 1999 and updates)

TABLE 1. ALBERTA TIER 1 SOIL REMEDIATION GUIDELINES

10. When two or more radionuclides are found, the following relationship should be satisfied:

$$\frac{[\text{Radionuclide}]_1}{\text{Guideline}_1} + \frac{[\text{Radionuclide}]_2}{\text{Guideline}_2} + \dots + \frac{[\text{Radionuclide}]_i}{\text{Guideline}_i} \leq 1$$

For more information see *Canadian Guidelines for the Management of Naturally Occurring Radioactive Materials (NORM)* (Health Canada, 2013)

11. If trichloroethene is found in soil, its degradation product vinyl chloride must also be measured and compared to guideline values
12. Guideline for protection of aquatic life is below detection limit and has not been included in the Tier 1 derivation. Groundwater monitoring is required for all land uses if the Tier 1 guideline is exceeded.
13. Guideline for protection of irrigation water is below detection limit and has not been included in the Tier 1 derivation. Groundwater monitoring is required for agricultural land use if the Tier 1 guideline is exceeded.
14. Analytical methodology specified in the Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
15. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008).
16. Toxicity assumptions for protection of aquatic life and livestock watering are based on mono- and dihydric phenols. Toxicity assumptions for other pathways are based on phenol. Soil samples may be analysed for phenolic compounds as a screening procedure. If analytical results exceed the guideline value, then background concentrations should be evaluated to determine if the phenolic compounds are naturally occurring. Speciation to determine mono- and dihydric phenols may also be undertaken and the results compared to the Tier 1 guideline.
17. Human health direct soil contact guidelines for carcinogenic PAHs are based on B[a]P Total Potency Equivalents (TPE). TPEs are calculated by multiplying the soil concentration of individual carcinogenic PAHs by a the standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (WHO/IPCS, 1998) scheme, as follows:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

18. The B[a]P Total Potency Equivalents (TPEs) calculated for specific soil samples using Potency Equivalency Factors (PEFs) should be multiplied by an Uncertainty Factor of 3 when evaluating PAH mixtures associated with creosote or coal tar-type environmental releases, prior to evaluating against the human health direct contact soil remediation guideline.

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
General and Inorganic Parameters									
pH	6.5-8.5	6.5-8.5	6.5-8.5	6.5-8.5	6.5-8.5	6.5-8.5	6.5-8.5	6.5-8.5	
Ammonia	see note 1	see note 1	see note 1	see note 1	see note 1	see note 1	see note 1	see note 1	1
Bromate	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
Chloride	120	100	120	120	120	100	120	120	
Cyanide (free)	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	
Electrical conductivity (dS/m)	-	1.0	-	-	-	1.0	-	-	
Fluoride	1.5	1.0	1.5	1.5	1.5	1.0	1.5	1.5	
Nitrate (as nitrogen)	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	
Nitrate + nitrite (as nitrogen)	-	100	-	-	-	100	-	-	
Nitrite (as nitrogen)	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	2
Sodium	200	200	200	200	200	200	200	200	
Sodium adsorption ratio	-	5.0	-	-	-	5.0	-	-	
Sulphate	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	2
Sulphide – Total (as S)	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	0.0019	8
Total Dissolved Solids (TDS)	500	500	500	500	500	500	500	500	
Metals									
Aluminum	see note 1	see note 2	see note 1	see note 1	see note 1	see note 2	see note 1	see note 1	1,2
Antimony	0.0060	0.0060	0.0060	0.0060	0.0060	0.0060	0.0060	0.0060	
Arsenic	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	
Barium	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	
Boron	1.50	1.0	1.50	1.50	1.50	1.0	1.50	1.50	
Cadmium	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	2
Chromium (trivalent)	0.0089	0.0049	0.0089	0.0089	0.0089	0.0049	0.0089	0.0089	
Chromium (hexavalent)	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Chromium (total)	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	9
Copper	0.0070	0.0070	0.0070	0.0070	0.0070	0.0070	0.0070	0.0070	

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Iron	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	
Lead	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	see note 2	2
Manganese	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	
Mercury (total)	0.0000050	0.0000050	0.0000050	0.0000050	0.0000050	0.0000050	0.0000050	0.0000050	9
Nickel	see note 1	see note 2	see note 1	see note 1	see note 1	see note 2	see note 1	see note 1	1,2
Selenium	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	
Silver	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	
Uranium	0.015	0.010	0.015	0.015	0.015	0.010	0.015	0.015	
Zinc	0.030	0.030	0.030	0.030	0.030	0.030	0.030	0.030	
Hydrocarbons									
Benzene	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	
Toluene	0.024	0.024	0.024	0.024	0.021	0.021	0.021	0.021	
Ethylbenzene	0.0016	0.0016	0.0016	0.0016	0.0016	0.0016	0.0016	0.0016	
Xylenes	0.020	0.020	0.020	0.020	0.020	0.020	0.020	0.020	
Styrene	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072	
F1	2.2	2.2	2.2	2.2	2.2	0.81	0.81	2.2	
F2	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	
Acenaphthene	0.0060	0.0060	0.0060	0.0060	0.0058	0.0058	0.0058	0.0058	
Anthracene	0.0034	0.0034	0.0034	0.0034	0.000012	0.000012	0.000012	0.000012	
Fluoranthene	0.24	0.24	0.24	0.86	0.000057	0.000057	0.000057	0.000057	
Fluorene	0.0042	0.0042	0.0042	0.0042	0.0030	0.0030	0.0030	0.0030	
Naphthalene	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Phenanthrene	0.00086	0.00086	0.00086	0.00086	0.00040	0.00040	0.00040	0.00040	
Pyrene	0.71	0.71	0.71	0.71	0.000092	0.000092	0.000092	0.000092	
Carcinogenic PAHs (as B(a)P TPE)	0.000040	0.000040	0.000040	0.000040	0.000040	0.000040	0.000040	0.000040	3
Benz[a]anthracene									
Benzo[b+j]fluoranthene									
Benzo[k]fluoranthene									
Benzo[g,h,i]perylene									

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Benzo[a]pyrene	0.0018	0.0018	0.0018	0.0066	0.0018	0.0018	0.0018	0.0066	4
Chrysene									
Dibenz[a,h]anthracene									
Indeno[1,2,3-c,d]pyrene									
Halogenated Aliphatics									
Vinyl chloride	0.0020	0.0020	0.0020	0.0020	0.0020	0.0011	0.0011	0.0020	
1,1-Dichloroethene	0.014	0.014	0.014	0.014	0.014	0.014	0.014	0.014	
Trichloroethene (Trichloroethylene, TCE)	0.0050	0.0050	0.0050	0.0050	0.0050	0.00032	0.00032	0.0038	5
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
1,2-Dichloroethane	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	0.0050	
Dichloromethane (Methylene chloride)	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.050	
Trichloromethane (Chloroform)	0.080	0.080	0.080	0.080	0.018	0.018	0.018	0.018	
Tetrachloromethane (Carbon tetrachloride)	0.0020	0.0020	0.0020	0.0020	0.0020	0.0015	0.0015	0.0020	
Dibromochloromethane	0.19	0.10	0.19	0.19	0.19	0.10	0.19	0.19	
Chlorinated Aromatics									
Chlorobenzene	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	
1,2-Dichlorobenzene	0.00070	0.00070	0.00070	0.00070	0.00070	0.00070	0.00070	0.00070	
1,4-Dichlorobenzene	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
1,2,3-Trichlorobenzene	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	0.0080	
1,2,4-Trichlorobenzene	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	
1,3,5-Trichlorobenzene	0.014	0.014	0.014	0.014	0.014	0.014	0.014	0.014	
1,2,3,4-Tetrachlorobenzene	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	
1,2,3,5-Tetrachlorobenzene	0.0038	0.0038	0.0038	0.0038	0.0038	0.0038	0.0038	0.0038	
1,2,4,5-Tetrachlorobenzene	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	
Pentachlorobenzene	0.0094	0.0094	0.0094	0.0094	0.0069	0.0069	0.0069	0.0069	
Hexachlorobenzene	0.00029	0.00029	0.00029	0.00029	0.00029	0.00029	0.00029	0.00029	
2,4-Dichlorophenol	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	
2,4,6-Trichlorophenol	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	



TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
2,3,4,6-Tetrachlorophenol	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Pentachlorophenol	0.00051	0.00051	0.00051	0.00051	0.00050	0.00050	0.00050	0.00050	
Dioxins & Furans	0.00000012	0.00000012	0.00000012	0.00000012	0.00000012	0.00000012	0.00000012	0.00000012	10
PCBs	0.000094	0.000094	0.000094	0.000094	0.000094	0.000094	0.000094	0.000094	
Pesticides									
Aldicarb	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Aldrin	0.000028	0.000028	0.000028	0.000028	0.000028	0.000028	0.000028	0.000028	
Atrazine and metabolites	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	
Azniphos-methyl (Guthion)	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	
Bromacil	0.0050	0.00020	0.0050	0.0050	0.0050	0.00020	0.0050	0.0050	
Bromoxynil	0.0050	0.00044	0.0050	0.0050	0.0050	0.00044	0.0050	0.0050	
Carbaryl	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	
Carbofuran	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	0.0018	
Chlorothalonil	0.00018	0.00018	0.00018	0.00018	0.00018	0.00018	0.00018	0.00018	
Chlorpyrifos	0.0000046	0.0000046	0.0000046	0.0000046	0.0000020	0.0000020	0.0000020	0.0000020	
2,4-D	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	0.0040	
DDT	0.0014	0.0014	0.0014	0.0014	0.0014	0.0014	0.0014	0.0014	
Diazinon	0.00017	0.00017	0.00017	0.00017	0.00017	0.00017	0.00017	0.00017	
Dicamba	0.010	0.0000080	0.010	0.010	0.010	0.0000080	0.010	0.010	
Diclofop-methyl	0.0090	0.00024	0.009	0.0090	0.0061	0.00024	0.0061	0.0061	
Dieldrin	0.000029	0.000029	0.000029	0.000029	0.000029	0.000029	0.000029	0.000029	
Dimethoate	0.0062	0.0030	0.0062	0.0062	0.0062	0.0030	0.0062	0.0062	
Dinoseb	0.000055	0.000055	0.000055	0.000055	0.000050	0.000050	0.000050	0.000050	
Diquat	0.070	0.070	0.070	0.070	0.070	0.070	0.070	0.070	
Diuron	0.15	0.15	0.15	0.15	0.15	0.15	0.15	0.15	
Endosulfan	0.0019	0.0019	0.0019	0.0019	0.0000031	0.0000031	0.0000031	0.0000031	
Endrin	0.0028	0.0028	0.0028	0.0028	0.0028	0.0028	0.0028	0.0028	
Glyphosate	0.065	0.065	0.065	0.065	0.065	0.065	0.065	0.065	
Heptachlor epoxide	0.000052	0.000052	0.000052	0.000052	0.000052	0.000052	0.000052	0.000052	

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Lindane	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	0.000010	
Linuron	0.0070	0.00011	0.0070	0.0070	0.0070	0.00011	0.0070	0.0070	
Malathion	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	0.00010	
MCPA	0.0026	0.000040	0.0026	0.0026	0.0026	0.000040	0.0026	0.0026	
Methoxychlor	0.90	0.90	0.90	0.90	0.00017	0.00017	0.00017	0.00017	
Metolachlor	0.0078	0.0078	0.0078	0.0078	0.0078	0.0078	0.0078	0.0078	
Metribuzin	0.0010	0.00050	0.0010	0.0010	0.0010	0.00050	0.0010	0.0010	
Paraquat (as dichloride)	0.010	0.010	0.010	0.010	0.010	0.010	0.010	0.010	
Phorate	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	0.0020	
Picloram	0.029	0.029	0.029	0.029	0.029	0.029	0.029	0.029	
Simazine	0.010	0.00050	0.010	0.010	0.010	0.00050	0.010	0.010	
Tebuthiuron	0.0016	0.00043	0.0016	0.0016	0.0016	0.00043	0.0016	0.0016	
Terbufos	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Toxaphene	0.00043	0.00043	0.00043	0.00043	0.00043	0.00043	0.00043	0.00043	
Triallate	0.00024	0.00024	0.00024	0.00024	0.00024	0.00024	0.00024	0.00024	
Trifluralin	0.0012	0.0012	0.0012	0.0012	0.00020	0.00020	0.00020	0.00020	
Other Organics									
Aniline	0.0022	0.0022	0.0022	0.0022	0.0022	0.0022	0.0022	0.0022	
Di- <i>n</i> -butyl phthalate	0.019	0.019	0.019	0.019	0.019	0.019	0.019	0.019	
Dichlorobenzidine	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	0.0010	
Diethanolamine	0.060	0.060	0.060	0.060	0.060	0.060	0.060	0.060	
Diethylene glycol	6.0	6.0	6.0	6.0	6.0	6.0	6.0	6.0	
Diisopropanolamine	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	
Ethylene glycol	31	31	31	31	31	31	31	31	
Hexachlorobutadiene	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	0.0013	
Methanol	19	19	19	19	19	19	19	19	
Methylmethacrylate	13	13	13	13	13	11	11	13	
Monoethanolamine	0.60	0.60	0.60	0.60	0.60	0.60	0.60	0.60	
MTBE	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Nitritotriacetic acid	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	
Nonylphenol + ethoxylates	0.0081	0.0081	0.0081	0.020	0.0081	0.0081	0.0081	0.020	
Perfluorooctane sulfonate (PFOS)	0.00060	0.00060	0.00060	0.00060	0.00060	0.00060	0.00060	0.00060	11
Perfluorooctanoic acid (PFOA)	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	0.00020	11
Phenol	0.0040	0.0020	0.0040	0.0040	0.0040	0.0020	0.0040	0.0040	6
Sulfolane	0.090	0.090	0.090	0.090	0.090	0.090	0.090	0.090	
Triethylene glycol	60	60	60	60	60	60	60	60	
Trihalomethanes - total (THMs)	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	
Naturally Occurring Radionuclides									
Beryllium-7	4,000	4,000	4,000	4,000	4,000	4,000	4,000	4,000	7
Bismuth-210	70	70	70	70	70	70	70	70	7
Lead-210	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	7
Polonium-210	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	7
Radium -224	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	7
Radium-226	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	7
Radium-228	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50	7
Thorium-228	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	7
Thorium-230	0.40	0.40	0.40	0.40	0.40	0.40	0.40	0.40	7
Thorium-232	0.10	0.10	0.10	0.10	0.10	0.10	0.10	0.10	7
Thorium-234	20	20	20	20	20	20	20	20	7
Uranium-234	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	7
Uranium-235	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	7
Uranium-238	4.0	4.0	4.0	4.0	4.0	4.0	4.0	4.0	7
Other Radionuclides									
Americium-241	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	7
Antimony-122	50	50	50	50	50	50	50	50	7
Antimony-124	40	40	40	40	40	40	40	40	7
Antimony-125	100	100	100	100	100	100	100	100	7
Barium-140	40	40	40	40	40	40	40	40	7

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Bromine-82	300	300	300	300	300	300	300	300	7
Calcium-45	200	200	200	200	200	200	200	200	7
Calcium-47	60	60	60	60	60	60	60	60	7
Carbon-14a	200	200	200	200	200	200	200	200	7
Cerium-141	100	100	100	100	100	100	100	100	7
Cerium-144	20	20	20	20	20	20	20	20	7
Cesium-131	2,000	2,000	2,000	2,000	2,000	2,000	2,000	2,000	7
Cesium-136	50	50	50	50	50	50	50	50	7
Cesium-137	10	10	10	10	10	10	10	10	7
Chromium-51	3,000	3,000	3,000	3,000	3,000	3,000	3,000	3,000	7
Cobalt-57	40	40	40	40	40	40	40	40	7
Cobalt-58	20	20	20	20	20	20	20	20	7
Cobalt-60	2.0	2.0	2.0	2.0	2.0	2.0	2.0	2.0	7
Gallium-67	500	500	500	500	500	500	500	500	7
Gold-198	90	90	90	90	90	90	90	90	7
Indium-111	400	400	400	400	400	400	400	400	7
Iodine-129	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	7
Iodine-131	6.0	6.0	6.0	6.0	6.0	6.0	6.0	6.0	7
Iron-55	300	300	300	300	300	300	300	300	7
Iron-59	40	40	40	40	40	40	40	40	7
Manganese-54	200	200	200	200	200	200	200	200	7
Mercury-197	400	400	400	400	400	400	400	400	7
Mercury-203	80	80	80	80	80	80	80	80	7
Neptunium-239	100	100	100	100	100	100	100	100	7
Niobium-95	200	200	200	200	200	200	200	200	7
Phosphorus-32	50	50	50	50	50	50	50	50	7
Plutonium-238	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	7
Plutonium-239	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	7
Plutonium-240	0.20	0.20	0.20	0.20	0.20	0.20	0.20	0.20	7
Plutonium-241	10	10	10	10	10	10	10	10	7
Rhodium-105	300	300	300	300	300	300	300	300	7

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Soil Type	Fine				Coarse				Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial/ Industrial	
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	
Rubidium-81	3,000	3,000	3,000	3,000	3,000	3,000	3,000	3,000	7
Rubidium-86	50	50	50	50	50	50	50	50	7
Ruthenium-103	100	100	100	100	100	100	100	100	7
Ruthenium-106	10	10	10	10	10	10	10	10	7
Selenium-75	70	70	70	70	70	70	70	70	7
Silver-108m	70	70	70	70	70	70	70	70	7
Silver-110m	50	50	50	50	50	50	50	50	7
Silver-111	70	70	70	70	70	70	70	70	7
Sodium-22	50	50	50	50	50	50	50	50	7
Strontium-85	300	300	300	300	300	300	300	300	7
Strontium-89	40	40	40	40	40	40	40	40	7
Strontium-90	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0	7
Sulphur-35	500	500	500	500	500	500	500	500	7
Technetium-99	200	200	200	200	200	200	200	200	7
Technetium-99m	7,000	7,000	7,000	7,000	7,000	7,000	7,000	7,000	7
Tellurium-129m	40	40	40	40	40	40	40	40	7
Tellurium-131m	40	40	40	40	40	40	40	40	7
Tellurium-132	40	40	40	40	40	40	40	40	7
Thallium-201	2,000	2,000	2,000	2,000	2,000	2,000	2,000	2,000	7
Tritium	7,000	7,000	7,000	7,000	7,000	7,000	7,000	7,000	7
Ytterbium-169	100	100	100	100	100	100	100	100	7
Yttrium-90	30	30	30	30	30	30	30	30	7
Yttrium-91	30	30	30	30	30	30	30	30	7
Zinc-65	40	40	40	40	40	40	40	40	7
Zirconium-95	100	100	100	100	100	100	100	100	7

- Notes:
1. See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014) for further guidance on aquatic life pathway.
  2. Tier 1 guideline = lowest of aquatic life guideline and all other guidelines (See Appendix B).
  3. B[a]P TPE (Total Potency Equivalents) are calculated by multiplying the groundwater concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the estimates of carcinogenic potential and are based on the World Health Organization (1999) scheme, as follows:

TABLE 2. ALBERTA TIER 1 GROUNDWATER REMEDIATION GUIDELINES

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

4. For ecological receptors only
5. If trichloroethene is found in groundwater, its degradation product vinyl chloride must also be measured and compared to guideline values
6. Toxicity assumptions for protection of aquatic life and livestock watering are based on mono- and dihydric phenols. Toxicity assumptions for other pathways are based on phenol. Groundwater samples may be analysed for phenolic compounds as a screening procedure. If analytical results exceed the guideline value, then background concentrations should be evaluated to determine if the phenolic compounds are naturally occurring. Speciation to determine mono- and dihydric phenols may also be undertaken and the results compared to the Tier 1 guideline.
7. When two or more radionuclides are found, the following relationship should be satisfied:

[Radionuclide]<sub>1</sub>

Guideline<sub>1</sub>

+

[Radionuclide]<sub>2</sub>

Guideline<sub>2</sub>

+

....

[Radionuclide]<sub>i</sub>

Guideline<sub>i</sub>

≤ 1

8. As S, but can be applied to undissociated H<sub>2</sub>S if concerns arise.

9. Total means all chemical species.

10. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-TCDD (See CCME, 1999 and updates)

11. As the toxicological effects of PFOA and perfluorooctanoyl sulfonate (PFOS) are considered to be additive, the sum of the ratios of the detected concentrations to the corresponding MACs for PFOS and PFOA should not exceed 1.

NGR = no guideline required - values for all exposure pathways that could be calculated are above compound solubility

Tier 1 Soil and Groundwater Remediation Guidelines

46

Classification: Public

TABLE 3. ALBERTA TIER 1 SUBSOIL REMEDIATION GUIDELINES (BTEX AND PHCS ONLY)

Soil Type	Fine					Coarse					Notes
Land Use	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	Natural Area	Agricultural	Residential/ Parkland	Commercial	Industrial	
Unit	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Benzene	0.046	0.046	0.046	0.046	0.046	0.078	0.021	0.021	0.078	0.078	2
Toluene	0.52	0.52	0.52	0.52	0.52	0.12	0.12	0.12	0.12	0.12	2
Ethylbenzene	0.073	0.073	0.073	0.073	0.073	0.14	0.14	0.14	0.14	0.14	2
Xylenes	0.99	0.99	0.99	0.99	0.99	1.9	1.9	1.9	1.9	1.9	2
F1	420	420	420	640	640	420	30	30	440	440	1, 3
F2	300	300	300	520	520	300	160	160	520	520	1, 3
F3	2,600	2,600	2,600	4,300	4,300	600	600	600	3,400	3,400	1, 3
F4	10,000	10,000	10,000	10,000	10,000	5,600	5,600	5,600	6,600	6,600	1, 3

Notes:

Guideline values calculated for this document using latest available CCME protocols except where noted.

- 1. Ecological direct contact pathway may be eliminated below 3 metres in depth for F1 to F4 only; the next lowest guideline value applies.
- 2. Ecological direct contact values from CCME (1999), other values calculated in this document.
- 3. All values calculated in this document.

**TABLE 4. ALBERTA TIER 1 SALT REMEDIATION GUIDELINES**

Rating Categories	Good	Fair	Poor	Unsuitable	Commercial/ Industrial
Topsoil <sup>e</sup>					
EC <sup>a</sup> (dS/m)	<2.0 <sup>c</sup>	2.0 to 4.0	4.0 to 8.0	>8.0	4.0
SAR <sup>b</sup>	<4.0	4.0 to 8.0	8.0 to 12	>12 <sup>d</sup>	12
Subsoil <sup>e</sup>					
EC <sup>a</sup> (dS/m)	<3.	3.0 to 5.0	5.0 to 10	>10	4.0
SAR <sup>b</sup>	<4.0	4.0 to 8	8.0 to 12	>12 <sup>d</sup>	12

**Notes:**

- Electrical conductivity, measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregoritch, 2008).
- Sodium Adsorption Ratio, measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregoritch, 2008).
- Material characterized by SAR of 12 to 20 may be rated as poor if texture is sandy loam or coarser and saturation % is less than 100.
- Topsoil: surface A, L, F, H, and O horizons on the control area, or the equivalent surface soil where these horizons are not present.  
Subsoil: B and C horizons and the upper portion of the parent material.






Figure 1: Example of the 30 m buffer zone for the more sensitive land use. The diagram is for illustration purposes and can be applied for any land use scenario where a more sensitive land use is adjacent to a less sensitive land use.





Figure 2: Implementation Framework for Tier 1, Tier 2 and Exposure Control Guidelines



**Figure 3: Expanded Flow Diagram – Tier 1**

**Appendix A**

**Soil Remediation Guidelines**

**All Exposure Pathways**

Appendix A consists of ten tables, comprised of surface soil and subsoil tables for each of the land uses. Each table provides the soil remediation guideline for each exposure pathway, where available. Tier 1 guidelines must be chosen from Tables 1 to 4, not from Appendix A or B. The only exception occurs when a more sensitive land use borders on, or is less than 30m from, the site of interest. Under this condition guidelines for specific exposure pathways on the more sensitive land use must be evaluated and if they are lower than the Tier 1 guideline, they must be applied to the site of interest (See Sections 3.2 and 5.1.1 and Figure 1).

The ten tables are as follows:

- Table A-1. Surface Soil Remediation Guidelines for Natural Area Land Use
- Table A-2. Surface Soil Remediation Guidelines for Agricultural Land Use
- Table A-3. Surface Soil Remediation Guidelines for Residential/Parkland Use
- Table A-4. Surface Soil Remediation Guidelines for Commercial Land Use
- Table A-5. Surface Soil Remediation Guidelines for Industrial Land Use
- Table A-6. Subsoil Remediation Guidelines for Natural Area Land Use
- Table A-7. Subsoil Remediation Guidelines for Agricultural Land Use
- Table A-8. Subsoil Remediation Guidelines for Residential/Parkland Use
- Table A-9. Subsoil Remediation Guidelines for Commercial Land Use
- Table A-10. Subsoil Remediation Guidelines for Industrial Land Use

In some cases, there is a value for the overall guideline, but no values for individual exposure pathways. This situation corresponds to non-risk-based guidelines that have been adopted from previous guideline documents, as explained in the main text.

All exposure pathways are applicable at Tier 1. However, it may be possible to exclude or modify certain pathways at Tier 2. The companion Tier 2 should be consulted for further information. The information in the tables in this appendix will assist in determining whether a Tier 2 approach for soil is likely to be useful at a given site.

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological								Other		
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit	
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
General and Inorganic Parameters															
pH (in 0.01M CaCl <sub>2</sub> )	6.0-8.5	6.0-8.5	-	-	6.0-8.5	6.0-8.5	-	-	-	-	-	-	-	-	-
Cyanide (free)	0.90	0.90	-	-	0.90	0.90	-	11	-	-	-	-	-	-	-
Fluoride	200	200	-	-	200	200	-	-	-	-	-	-	-	-	-
Sulphur (elemental) <sup>a</sup>	500	500	-	-	500	500	-	-	-	-	-	-	-	-	-
Metals															
Antimony	20	20	-	-	20	20	-	-	-	-	-	-	-	-	-
Arsenic (inorganic)	17	17	-	-	17	17	-	380	-	-	-	-	-	-	-
Barium (non-barite)	750	750	-	-	750	750	-	-	-	-	-	-	-	-	-
Barite-barium <sup>b</sup>	10,000	10,000	-	-	200,000	200,000	-	30,000	10,000	-	-	-	-	-	-
Beryllium	5.0	5.0	-	-	5.0	5.0	-	-	-	-	-	-	-	-	-
Boron (mg/L in saturated paste extract) <sup>k</sup>	3.3	3.3	65	120	3.3	3.3	-	-	-	5.0	5.0	17	17	-	-
Cadmium	3.8	3.8	-	-	10	10	54	3.8	-	-	-	-	-	-	-
Chromium (hexavalent)	0.40	0.40	-	-	0.40	0.40	-	-	-	-	-	-	-	-	-
Chromium (total)	64	64	-	-	64	64	-	-	-	-	-	-	-	-	-
Cobalt	20	20	-	-	20	20	-	-	-	-	-	-	-	-	-
Copper	63	63	-	-	63	63	350	300	-	-	-	-	-	-	-
Lead	70	70	-	-	300	300	720	70	-	-	-	-	-	-	-
Mercury (inorganic)	12	12	-	-	12	12	20	-	-	-	-	-	-	-	-
Molybdenum	4.0	4.0	-	-	4.0	4.0	-	-	-	-	-	-	-	-	-

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human		Ecological								Other	
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit		
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Nickel	45	45	-	-	45	45	170	530	-	-	-	-	-	-	-	
Selenium	1.0	1.0	-	-	1.0	1.0	-	4.5	-	-	-	-	-	-	-	
Silver	20	20	-	-	20	20	-	-	-	-	-	-	-	-	-	
Thallium	1.0	1.0	-	-	1.4	1.4	-	1.0	-	-	-	-	-	-	-	
Tin	5.0	5.0	-	-	5.0	5.0	-	-	-	-	-	-	-	-	-	
Uranium	33	33	-	-	500	500	-	33	-	-	-	-	-	-	-	
Vanadium	130	130	-	-	130	130	260	-	-	-	-	-	-	-	-	
Zinc	250	250	-	-	250	250	280	980	-	-	-	-	-	-	-	
Hydrocarbons																
Benzene	0.046	0.078	0.046	0.078	60	31	-	44	18	7.9	0.17	15	0.33	-	-	
Toluene	0.52	0.12	0.52	0.95	110	75	-	2,500	980	63,000	0.12	NGR	1,000	-	-	
Ethylbenzene	0.073	0.14	0.073	0.14	120	55	-	1,600	640	NGR	540	NGR	17,000	-	-	
Xylenes	0.99	1.9	0.99	1.9	65	95	-	6,600	2,600	NGR	41	NGR	16,000	-	-	
Styrene	0.68	0.8	110	210	-	-	-	-	-	0.68	0.8	-	-	-	-	
F1	210	210	1,100	2,200	210	210	-	27,000	11,000	30,000	1,300	30,000	30,000	800	700	
F2	150	150	1,500	2,900	150	150	-	25,000	9,800	30,000	520	30,000	30,000	1,000	1,000	
F3	1,300	300	-	-	1,300	300	-	30,000	16,000	-	-	-	-	3,500	2,500	
F4	5,600	2,800	-	-	5,600	2,800	-	21,000	8,400	-	-	-	-	10,000	10,000	
Acenaphthene	0.33	0.38	NGR	NGR	-	-	-	22	22	0.33	0.38	NGR	NGR	-	-	
Anthracene	1.3	0.0056	NGR	NGR	2.5	2.5	-	62	62	1.3	0.0056	NGR	NGR	-	-	
Fluoranthene	15	0.055	NGR	NGR	50	50	-	15	15	NGR	0.055	NGR	NGR	-	-	

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological								Other		
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit	
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Fluorene	0.40	0.34	NGR	NGR	-	-	-	15	15	0.40	0.34	NGR	NGR	-	-
Naphthalene	0.014	0.017	28	53	-	-	-	8.8	8.8	0.014	0.017	NGR	NGR	-	-
Phenanthrene	0.11	0.061	-	-	-	-	-	43	43	0.11	0.061	NGR	NGR	-	-
Pyrene	7.7	0.15	NGR	NGR	-	-	-	7.7	7.7	NGR	0.15	NGR	NGR	-	-
Carcinogenic PAHs	IACR<1.0 °	IACR<1.0 °	IACR<1.0 °	IACR<1.0 °	-	-	-	-	-	-	-	-	-	-	-
Benz[a]anthracene <sup>d</sup>	6.2	6.2	6.4	12	-	-	-	6.2	6.2	NGR	NGR	NGR	NGR	-	-
Benzo[b+j]fluoranthene <sup>d</sup>	6.2	6.2	3.0	5.8	-	-	-	6.2	6.2	-	-	NGR	NGR	-	-
Benzo[k]fluoranthene <sup>d</sup>	6.2	6.2	0.64	1.2	-	-	-	6.2	6.2	-	-	NGR	NGR	-	-
Benzo[g,h,i]perylene	-	-	130	250	-	-	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>d</sup>	0.60	0.60	7.0	14	20	20	-	0.60	0.60	NGR	NGR	NGR	NGR	-	-
Chrysene <sup>d</sup>	6.2	6.2	40	78	-	-	-	6.2	6.2	-	-	NGR	NGR	-	-
Dibenz[a,h]anthracene	-	-	4.4	8.5	-	-	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	51	98	-	-	-	-	-	-	-	-	-	-	-
Chlorinated Aliphatics															
Vinyl chloride	0.014	0.020	0.014	0.020	-	-	-	-	-	-	-	-	-	-	-
1,1-Dichloroethene	0.15	0.24	0.15	0.24	-	-	-	-	-	-	-	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.054	0.081	0.054	0.093	3.0	3.0	-	-	-	0.72	0.081	-	-	-	-
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.26	0.46	0.26	0.46	-	-	-	-	-	0.69	0.77	-	-	-	-
1,2-Dichloroethane	0.025	0.041	0.025	0.041	-	-	-	-	-	0.12	0.12	-	-	-	-
Dichloromethane (Methylene chloride)	0.10	0.095	0.21	0.32	-	-	-	-	-	0.10	0.095	-	-	-	-

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological								Other		
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit	
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Trichloromethane (Chloroform) <sup>1</sup>	0.16	0.030	0.53	0.88	-	-	-	-	-	0.16	0.030	-	-	-	-
Tetrachloromethane (Carbon tetrachloride)	0.037	0.062	0.037	0.062	-	-	-	-	-	0.059	0.062	-	-	-	-
Dibromochloromethane	0.91	1.5	0.91	1.5	-	-	-	-	-	-	-	-	-	-	-
Chlorinated Aromatics															
Chlorobenzene <sup>e</sup>	0.61	1.1	0.61	1.1	-	-	-	-	-	BDL	BDL	-	-	-	-
1,2-Dichlorobenzene <sup>e</sup>	0.097	0.18	0.097	0.18	-	-	-	-	-	BDL	BDL	-	-	-	-
1,4-Dichlorobenzene	0.051	0.098	0.051	0.098	-	-	-	-	-	0.32	0.38	-	-	-	-
1,2,3-Trichlorobenzene	0.26	0.31	1.9	3.6	-	-	-	-	-	0.26	0.31	-	-	-	-
1,2,4-Trichlorobenzene	0.78	0.93	2.0	3.9	-	-	-	-	-	0.78	0.93	-	-	-	-
1,3,5-Trichlorobenzene	1.9	3.6	1.9	3.6	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.042	0.050	3.1	5.9	-	-	-	-	-	0.042	0.050	-	-	-	-
1,2,3,5-Tetrachlorobenzene	0.37	0.70	0.37	0.70	-	-	-	-	-	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.19	0.37	0.19	0.37	-	-	-	-	-	-	-	-	-	-	-
Pentachlorobenzene	24	5.2	24	47	-	-	-	-	-	NGR	5.2	-	-	-	-
Hexachlorobenzene	1.8	3.6	1.8	3.6	-	-	-	-	-	-	-	-	-	-	-
2,4-Dichlorophenol	0.0029	0.0034	0.018	0.034	-	-	-	-	-	0.0029	0.0034	-	-	-	-
2,4,6-Trichlorophenol	0.19	0.37	0.19	0.37	-	-	-	-	-	0.42	0.50	-	-	-	-
2,3,4,6-Tetrachlorophenol	0.039	0.047	0.16	0.31	-	-	-	-	-	0.039	0.047	-	-	-	-
Pentachlorophenol	0.025	0.029	6.0	12	11	11	-	-	-	0.025	0.029	-	-	-	-
Dioxins & Furans <sup>f,g</sup>	0.00025	0.00025	-	-	-	-	-	0.00025	0.00025	-	-	-	-	-	-
PCBs	1.3	1.3	-	-	33	33	-	1.3	1.3	-	-	-	-	-	-



TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological								Other		
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit	
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Pesticides															
Aldicarb °	0.041	0.065	0.041	0.065	-	-	-	-	-	BDL	BDL	-	-	-	-
Aldrin	0.24	0.46	0.24	0.46	-	-	-	-	-	-	-	-	-	-	-
Atrazine and metabolites	0.0088	0.010	0.10	0.19	-	-	-	-	-	0.0088	0.010	-	-	-	-
Azniphos-methyl (Guthion)	0.41	0.75	0.41	0.75	-	-	-	-	-	-	-	-	-	-	-
Bromacil <sup>l</sup>	0.0090	0.0090	7.0	10	0.20	0.12	-	-	-	0.0090	0.0090	-	-	-	-
Bromoxynil	0.044	0.052	0.18	0.35	-	-	-	-	-	0.044	0.052	-	-	-	-
Carbaryl °	1.9	3.6	1.9	3.6	-	-	-	-	-	BDL	BDL	-	-	-	-
Carbofuran °	0.68	1.2	0.68	1.2	-	-	-	-	-	BDL	BDL	-	-	-	-
Chlorothalonil	0.0085	0.010	27	53	-	-	-	-	-	0.0085	0.010	-	-	-	-
Chlorpyrifos °	49	95	49	95	-	-	-	-	-	BDL	BDL	-	-	-	-
2,4-D °	0.43	0.67	0.43	0.67	-	-	-	-	-	BDL	BDL	-	-	-	-
DDT	0.70	0.70	89	170	12	12	547	0.70	0.70	-	-	-	-	-	-
Diazinon °	2.2	4.2	2.2	4.2	-	-	-	-	-	BDL	BDL	-	-	-	-
Dicamba °	0.50	0.79	0.50	0.79	-	-	-	-	-	BDL	BDL	-	-	-	-
Diclofop-methyl	NGR	2.4	NGR	NGR	-	-	-	-	-	NGR	2.4	-	-	-	-
Dieldrin	0.025	0.048	0.025	0.048	-	-	-	-	-	-	-	-	-	-	-
Dimethoate	0.0058	0.0055	0.077	0.12	-	-	-	-	-	0.0058	0.0055	-	-	-	-
Dinoseb °	2.8	5.5	2.8	5.5	-	-	-	-	-	BDL	BDL	-	-	-	-
Diquat	11	21	11	21	-	-	-	-	-	-	-	-	-	-	-
Diuron	1.9	3.5	1.9	3.5	-	-	-	-	-	-	-	-	-	-	-

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological										Other	
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit		
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Endosulfan	0.80	0.0016	99	190	-	-	-	-	-	0.80	0.0016	-	-	-	-	
Endrin	2.4	4.7	2.4	4.7	-	-	-	-	-	-	-	-	-	-	-	
Glyphosate	0.054	0.049	0.95	1.4	-	-	-	-	-	0.054	0.049	-	-	-	-	
Heptachlor epoxide	0.039	0.076	0.039	0.076	-	-	-	-	-	-	-	-	-	-	-	
Lindane °	0.31	0.60	0.31	0.60	-	-	-	-	-	BDL	BDL	-	-	-	-	
Linuron	0.051	0.059	0.56	1.1	-	-	-	-	-	0.051	0.059	-	-	-	-	
Malathion °	0.82	1.3	0.82	1.3	-	-	-	-	-	BDL	BDL	-	-	-	-	
MCPA °	0.42	0.66	0.42	0.66	-	-	-	-	-	BDL	BDL	-	-	-	-	
Methoxychlor	NGR	0.32	NGR	NGR	-	-	-	-	-	NGR	0.32	-	-	-	-	
Metolachlor	0.048	0.055	1.3	2.4	-	-	-	-	-	0.048	0.055	-	-	-	-	
Metribuzin	0.024	0.028	7.8	15	-	-	-	-	-	0.024	0.028	-	-	-	-	
Paraquat (as dichloride)	1.1	2.2	1.1	2.2	-	-	-	-	-	-	-	-	-	-	-	
Phorate	0.075	0.14	0.075	0.14	-	-	-	-	-	-	-	-	-	-	-	
Picloram	0.024	0.022	0.64	0.94	-	-	-	-	-	0.024	0.022	-	-	-	-	
Simazine	0.033	0.038	0.14	0.25	-	-	-	-	-	0.033	0.038	-	-	-	-	
Tebuthiuron °j	0.046	0.046	2.5	3.7	0.046	0.046	-	-	-	BDL	BDL	-	-	-	-	
Terbufos	0.080	0.15	0.080	0.15	-	-	-	-	-	-	-	-	-	-	-	
Toxaphene	3.3	6.3	3.3	6.3	-	-	-	-	-	-	-	-	-	-	-	
Triallate	0.0077	0.0092	16	31	-	-	-	-	-	0.0077	0.0092	-	-	-	-	
Trifluralin	0.22	0.045	NGR	NGR	-	-	-	-	-	0.22	0.045	-	-	-	-	

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human		Ecological								Other		
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit	
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Other Organics															
Aniline <sup>e</sup>	0.36	0.60	0.36	0.60	-	-	-	-	-	BDL	BDL	-	-	-	-
Di- <i>n</i> -butyl phthalate	0.54	0.65	70	130	-	-	-	-	-	0.54	0.65	-	-	-	-
Dichlorobenzidine	0.60	1.2	0.60	1.2	-	-	-	-	-	-	-	-	-	-	-
Diethanolamine <sup>h</sup>	2.0	3.5	2.0	3.5	1,000	1,000	-	-	-	500,000	45	-	-	-	-
Diethylene glycol	10	15	10	15	1,000	1,000	-	-	-	2,000	65	-	-	-	-
Diisopropanolamine	14	17	130	250	360	360	-	-	-	14	17	-	-	-	-
Ethylene glycol	60	62	60	68	1,100	1,100	1,700	-	-	89	62	-	-	-	-
Hexachlorobutadiene	0.026	0.031	0.50	0.95	-	-	-	-	-	0.026	0.031	-	-	-	-
Methanol	37	11	37	42	1,200	1,200	-	-	-	300	11	-	-	750	750
Methylmethacrylate	36	49	36	49	-	-	-	-	-	-	-	-	-	-	-
Monoethanolamine <sup>h</sup>	20	10	20	40	1,500	1,500	-	-	-	300,000	10	-	-	-	-
MTBE	0.044	0.062	0.044	0.062	-	-	-	-	-	7.1	6.1	-	-	-	-
Nonylphenol + ethoxylates	5.7	5.7	-	-	5.7	5.7	-	-	-	NGR	2,000	-	-	-	-
Perfluorooctane sulfonate (PFOS)	0.010	0.010	0.070	0.14	10	10	-	0.010	0.010	0.19	0.23	1.5	1.8	-	-
Phenol	0.0028	0.0024	1.6	2.3	20	20	-	-	-	0.0028	0.0024	-	-	-	-
Sulfolane	0.18	0.21	0.18	0.21	210	210	-	-	-	24	18	-	-	-	-
Triethylene glycol	100	150	100	150	5,000	5,000	-	-	-	10,000	200	-	-	-	-

Notes:

a. For more information see *Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils* (AENV, 2011)

TABLE A-1. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

- b. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)
- c. The Index of Additive Cancer Risk (IACR) is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture, as follows:

Fine Soil:

$$IACR = \frac{[Benz(a)anthracene]}{6.4} + \frac{[Benzo(b + j)fluoranthene]}{3.0} + \frac{[Benzo(k)fluoranthene]}{0.64} + \frac{[Benzo(g, h, i)perylene]}{130} + \frac{[Benzo(a)pyrene]}{7.0} + \frac{[Chrysene]}{40} + \frac{[Dibenz(a, h)anthracene]}{4.4} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{51}$$

Coarse Soil:

$$IACR = \frac{[Benz(a)anthracene]}{12} + \frac{[Benzo(b + j)fluoranthene]}{5.8} + \frac{[Benzo(k)fluoranthene]}{1.2} + \frac{[Benzo(g, h, i)perylene]}{250} + \frac{[Benzo(a)pyrene]}{14} + \frac{[Chrysene]}{78} + \frac{[Dibenz(a, h)anthracene]}{8.5} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{98}$$

- d. Overall guideline value for ecological receptors only.
- e. Guideline for protection of aquatic life is below detection limit, groundwater monitoring is required.
- f. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- g. Guideline values adopted directly from CCME (1999 and updates) without change.
- h. Analytical methodology specified in the *Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine* (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
- i. Eco-contact guidelines from Stantec (2012)
- j. Eco-contact guidelines from Stantec (2008)
- k. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008)
- l. Guideline for protection of aquatic life (fine soil) is based on a groundwater guideline of 0.10 g/L. See Appendix B for more information.

BDL - Below detection limit

NGR - no guideline required, calculated value >1,000,000 mg/kg; or for PAH groundwater protection, calculated value results in groundwater concentration greater than solubility

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human						Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit			
	Soil Type	Fine		Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
General and Inorganic Parameters																									
pH (in 0.01M CaCl <sub>2</sub> )	6.0-8.5	6.0-8.5	-	-	-	-	-	-	-	6.0-8.5	6.0-8.5	-	-	-	-	-	-	-	-	-	-	-	-	-	
Cyanide (free)	0.90	0.90	29	-	-	-	-	-	-	0.90	0.90	-	11	-	-	-	-	-	-	-	-	-	-	-	
Fluoride	200	200	-	-	-	-	-	-	-	200	200	-	-	-	-	-	-	-	-	-	-	-	-	-	
Sulphur (elemental) <sup>a</sup>	500	500	-	-	-	-	-	-	-	500	500	-	-	-	-	-	-	-	-	-	-	-	-	-	
Metals																									
Antimony	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-	-	-	-	-	-	-	-	-	
Arsenic (inorganic)	17	17	27	-	-	-	-	-	-	17	17	-	380	-	-	-	-	-	-	-	-	-	-	-	
Barium (non-barite)	750	750	6,800	-	-	-	-	-	-	750	750	-	-	-	-	-	-	-	-	-	-	-	-	-	
Barite-barium <sup>b</sup>	10,000	10,000	10,000	-	-	-	-	-	-	200,000	200,000	-	30,000	10,000	-	-	-	-	-	-	-	-	-	-	
Beryllium	5.0	5.0	-	-	-	-	-	-	-	5.0	5.0	-	-	-	-	-	-	-	-	-	-	-	-	-	
Boron (mg/L in saturated paste extract) <sup>m</sup>	3.3	3.3	7,500	-	-	-	-	65	120	3.3	3.3	-	-	-	5.0	5.0	17	17	17	17	3.4	3.4	-	-	
Cadmium	1.4	1.4	1.4	-	-	-	-	-	-	10	10	54	3.8	-	-	-	-	-	-	-	-	-	-	-	
Chromium (hexavalent)	0.40	0.40	-	-	-	-	-	-	-	0.40	0.40	-	-	-	-	-	-	-	-	-	-	-	-	-	
Chromium (total)	64	64	220	-	-	-	-	-	-	64	64	-	-	-	-	-	-	-	-	-	-	-	-	-	
Cobalt	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-	-	-	-	-	-	-	-	-	
Copper	63	63	1,100	-	-	-	-	-	-	63	63	350	300	-	-	-	-	-	-	-	-	-	-	-	
Lead	70	70	140	-	-	-	-	-	-	300	300	720	70	-	-	-	-	-	-	-	-	-	-	-	
Mercury (inorganic)	6.6	6.6	6.6	-	-	-	-	-	-	12	12	20	-	-	-	-	-	-	-	-	-	-	-	-	
Molybdenum	4.0	4.0	-	-	-	-	-	-	-	4.0	4.0	-	-	-	-	-	-	-	-	-	-	-	-	-	
Nickel	45	45	200	-	-	-	-	-	-	45	45	170	530	-	-	-	-	-	-	-	-	-	-	-	

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit		
	Fine	Coarse		-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Selenium	1.0	1.0	80	-	-	-	-	-	-	1.0	1.0	-	4.5	-	-	-	-	-	-	-	-	-	-	-
Silver	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-	-	-	-	-	-	-	-	
Thallium	1.0	1.0	1.0	-	-	-	-	-	-	1.4	1.4	-	1.0	-	-	-	-	-	-	-	-	-	-	
Tin	5.0	5.0	-	-	-	-	-	-	-	5.0	5.0	-	-	-	-	-	-	-	-	-	-	-	-	
Uranium	23	23	23	-	-	-	-	-	-	500	500	-	33	-	-	-	-	-	-	-	-	-	-	
Vanadium	130	130	-	-	-	-	-	-	-	130	130	260	-	-	-	-	-	-	-	-	-	-	-	
Zinc	250	250	10,000	-	-	-	-	-	-	250	250	280	980	-	-	-	-	-	-	-	-	-	-	
Hydrocarbons																								
Benzene	0.046	0.015	360	0.36	0.33	0.021	0.015	0.046	0.078	60	31	-	44	18	7.9	0.17	0.20	0.21	15	0.33	-	-	-	-
Toluene	0.52	0.12	690	1,300	1,200	80	57	0.52	0.95	110	75	-	2,500	980	63,000	0.12	26	29	NGR	1,000	-	-	-	-
Ethylbenzene	0.073	0.14	1,900	2,100	1,900	120	88	0.073	0.14	120	55	-	1,600	640	NGR	540	36	42	NGR	17,000	-	-	-	-
Xylenes	0.99	1.9	510	150	140	9.0	6.6	0.99	1.9	65	95	-	6,600	2,600	NGR	41	160	180	NGR	16,000	-	-	-	-
Styrene	0.68	0.80	12,000	2,300	2,100	130	94	110	210	-	-	-	-	-	0.68	0.80	-	-	-	-	-	-	-	
F1	210	24	14,000	710	610	30	24	1,100	2,200	210	210	-	27,000	11,000	30,000	1,300	6,600	7,400	30,000	30,000	-	-	800	700
F2	150	130	6,800	3,600	3,100	160	130	1,500	2,900	150	150	-	25,000	9,800	30,000	520	16,000	19,000	30,000	30,000	-	-	1,000	1,000
F3	1,300	300	15,000	-	-	-	-	-	-	1,300	300	-	30,000	16,000	-	-	-	-	-	-	-	-	3,500	2,500
F4	5,600	2,800	21,000	-	-	-	-	-	-	5,600	2,800	-	21,000	8,400	-	-	-	-	-	-	-	-	10,000	10,000
Acenaphthene	0.33	0.38	5,500	120,000	99,000	4,800	3,900	NGR	NGR	-	-	-	22	22	0.33	0.38	NGR	NGR	NGR	NGR	-	-	-	-
Anthracene	1.3	0.0056	27,000	NGR	NGR	780,000	670,000	NGR	NGR	2.5	2.5	-	62	62	1.3	0.0056	NGR	NGR	NGR	NGR	-	-	-	-
Fluoranthene	15.4	0.055	3,700	NGR	NGR	550,000	480,000	NGR	NGR	50	50	-	15	15	NGR	0.055	NGR	NGR	NGR	NGR	-	-	-	-
Fluorene	0.40	0.34	2,800	270,000	220,000	10,000	8,600	NGR	NGR	-	-	-	15	15	0.40	0.34	NGR	NGR	NGR	NGR	-	-	-	-
Naphthalene	0.014	0.017	2,000	260	220	13	9.6	28	53	-	-	-	8.8	8.8	0.014	0.017	NGR	NGR	NGR	NGR	-	-	-	-

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological												Other		
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit			
				Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Soil Type		Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type				-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)		(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Phenanthrene		0.11	0.061	-	-	-	-	-	-	-	-	-	43	43	0.11	0.061	NGR	NGR	NGR	NGR	-	-	-	-	
Pyrene		7.7	0.15	2,200	NGR	NGR	810,000	730,000	NGR	NGR	-	-	7.7	7.7	NGR	0.15	NGR	NGR	NGR	NGR	-	-	-	-	
Carcinogenic PAHs (as B(a)P TPE) <sup>c</sup>		IACR<1.0 <sup>e</sup> and TPE ≤ 5.6	IACR<1.0 <sup>e</sup> and TPE ≤ 5.6	5.6 <sup>d</sup>	NGR	NGR	NGR	NGR	IACR<1.0 <sup>e</sup>	IACR<1.0 <sup>e</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-		
Benz[a]anthracene <sup>f</sup>		6.2	6.2	-	-	-	-	-	6.4	12	-	-	-	6.2	6.2	NGR	NGR	NGR	NGR	NGR	NGR	-	-	-	-
Benzo[b+] fluoranthene <sup>f</sup>		6.2	6.2	-	-	-	-	-	3.0	5.8	-	-	-	6.2	6.2	-	-	NGR	NGR	NGR	NGR	-	-	-	-
Benzo[k] fluoranthene <sup>f</sup>		6.2	6.2	-	-	-	-	-	0.64	1.2	-	-	-	6.2	6.2	-	-	NGR	NGR	NGR	NGR	-	-	-	-
Benzo[g,h,i]perylene		-	-	-	-	-	-	-	130	250	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Benzo[a]pyrene <sup>f</sup>		0.60	0.60	-	-	-	-	-	7.0	14	20	20	-	0.60	0.60	NGR	NGR	NGR	NGR	NGR	NGR	-	-	-	-
Chrysene <sup>f</sup>		6.2	6.2	-	-	-	-	-	40	78	-	-	-	6.2	6.2	-	-	NGR	NGR	NGR	NGR	-	-	-	-
Dibenz[a,h]anthracene		-	-	-	-	-	-	-	4.4	8.5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Indeno[1,2,3-c,d]pyrene		-	-	-	-	-	-	-	51	98	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Chlorinated Aliphatics																									
Vinyl chloride		0.0083	0.00034	63	0.0090	0.0083	0.00049	0.00034	0.014	0.020	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
1,1-Dichloroethene		0.15	0.021	120	0.50	0.46	0.030	0.021	0.15	0.24	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Trichloroethene (Trichloroethylene, TCE)		0.0043	0.00019	37	0.0048	0.0043	0.00026	0.00019	0.054	0.093	3.0	3.0	-	-	-	0.72	0.081	0.13	0.14	-	-	-	-	-	
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)		0.26	0.018	190	0.46	0.41	0.025	0.018	0.26	0.46	-	-	-	-	-	0.69	0.77	-	-	-	-	-	-	-	
1,2-Dichloroethane		0.0062	0.0027	9,100	0.060	0.055	0.0038	0.0027	0.025	0.041	-	-	-	-	-	0.12	0.12	0.0062	0.0062	-	-	-	-	-	

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit		
				-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Soil Type		Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type				-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)		(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Dichloromethane (Methylene chloride)		0.052	0.048	340	4.8	4.4	0.28	0.19	0.21	0.32	-	-	-	-	-	0.10	0.095	0.052	0.048	-	-	-	-	-
Trichloromethane (Chloroform) <sup>n</sup>		0.16	0.030	76	3.2	3.0	0.21	0.15	0.53	0.88	-	-	-	-	-	0.16	0.030	0.16	0.17	-	-	-	-	-
Tetrachloromethane (Carbon tetrachloride)		0.022	0.0015	29	0.038	0.035	0.0021	0.0015	0.037	0.062	-	-	-	-	-	0.059	0.062	0.022	0.023	-	-	-	-	-
Dibromochloromethane		0.12	0.12	800	11	7.8	0.28	0.27	0.91	1.5	-	-	-	-	-	-	-	0.12	0.12	-	-	-	-	-
Chlorinated Aromatics																								
Chlorobenzene <sup>g</sup>		0.39	0.018	17,000	0.44	0.39	0.024	0.018	0.61	1.1	-	-	-	-	-	BDL	BDL	-	-	-	-	-	-	-
1,2-Dichlorobenzene <sup>g</sup>		0.097	0.18	17,000	260	230	14	10	0.097	0.18	-	-	-	-	-	BDL	BDL	-	-	-	-	-	-	-
1,4-Dichlorobenzene		0.051	0.098	4,400	10	9.1	0.56	0.42	0.051	0.098	-	-	-	-	-	0.32	0.38	-	-	-	-	-	-	-
1,2,3-Trichlorobenzene		0.26	0.26	51	8.8	6.8	0.30	0.26	1.9	3.6	-	-	-	-	-	0.26	0.31	-	-	-	-	-	-	-
1,2,4-Trichlorobenzene		0.78	0.23	40	7.6	6.0	0.26	0.23	2.0	3.9	-	-	-	-	-	0.78	0.93	-	-	-	-	-	-	-
1,3,5-Trichlorobenzene		1.9	0.13	47	4.1	3.2	0.14	0.13	1.9	3.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene		0.042	0.050	75	27	20	0.88	0.84	3.1	5.9	-	-	-	-	-	0.042	0.050	-	-	-	-	-	-	-
1,2,3,5-Tetrachlorobenzene		0.37	0.10	16	3.3	2.5	0.10	0.10	0.37	0.70	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene		0.19	0.052	4.6	1.7	1.3	0.054	0.052	0.19	0.37	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pentachlorobenzene		22	5.2	22	160	140	7.9	6.1	24	47	-	-	-	-	-	NGR	5.2	-	-	-	-	-	-	-
Hexachlorobenzene		0.80	0.20	3.3	5.4	4.7	0.26	0.20	1.8	3.6	-	-	-	-	-	-	-	0.80	0.97	-	-	-	-	-
2,4-Dichlorophenol		0.0029	0.0034	4,000	170,000	140,000	6,300	5,400	0.018	0.034	-	-	-	-	-	0.0029	0.0034	-	-	-	-	-	-	-
2,4,6-Trichlorophenol		0.19	0.37	260	11,000	8,200	360	320	0.19	0.37	-	-	-	-	-	0.42	0.50	-	-	-	-	-	-	-
2,3,4,6-Tetrachlorophenol		0.039	0.047	400	15,000	11,000	480	460	0.16	0.31	-	-	-	-	-	0.039	0.047	-	-	-	-	-	-	-
Pentachlorophenol		0.025	0.029	130	NGR	NGR	110,000	83,000	6.0	12	11	11	-	-	-	0.025	0.029	-	-	-	-	-	-	-
Dioxins & Furans <sup>h,i</sup>		0.0000040	0.0000040	0.0000040	-	-	-	-	-	-	-	-	-	0.00025	0.00025	-	-	-	-	-	-	-	-	-



TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human							Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit			
	Soil Type	Fine		Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
PCBs	0.085	0.085	0.085	-	-	-	-	-	-	33	33	-	1.3	1.3	-	-	-	-	-	-	-	-	-	-	
Pesticides																									
Aldicarb <sup>g</sup>	0.012	0.012	22	-	-	-	-	0.041	0.065	-	-	-	-	-	BDL	BDL	0.012	0.012	-	-	0.079	0.078	-	-	
Aldrin	0.24	0.31	0.31	44	30	1.4	1.5	0.24	0.46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Atrazine and metabolites	0.0088	0.010	11	-	-	-	-	0.10	0.19	-	-	-	-	-	0.0088	0.010	0.025	0.028	-	-	0.049	0.057	-	-	
Azinphos-methyl (Guthion)	0.41	0.75	55	-	-	-	-	0.41	0.75	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Bromacil <sup>g,k</sup>	0.0090	0.0090	2,000	-	-	-	-	7.0	10	0.20	0.12	-	-	-	0.0090	0.0090	2.0	2.0	-	-	BDL	BDL	-	-	
Bromoxynil <sup>g</sup>	0.044	0.052	11	-	-	-	-	0.18	0.35	-	-	-	-	-	0.044	0.052	0.097	0.11	-	-	BDL	BDL	-	-	
Carbaryl <sup>g</sup>	1.9	3.6	220	-	-	-	-	1.9	3.6	-	-	-	-	-	BDL	BDL	5.7	6.5	-	-	-	-	-	-	
Carbofuran <sup>g</sup>	0.082	0.089	220	-	-	-	-	0.68	1.2	-	-	-	-	-	BDL	BDL	0.082	0.089	-	-	-	-	-	-	
Chlorothalonil	0.0085	0.010	330	-	-	-	-	27	53	-	-	-	-	-	0.0085	0.010	7.9	9.5	-	-	0.43	0.52	-	-	
Chlorpyrifos <sup>g</sup>	3.2	3.8	220	-	-	-	-	49	95	-	-	-	-	-	BDL	BDL	3.2	3.8	-	-	-	-	-	-	
2,4-D <sup>g</sup>	0.10	0.10	400	-	-	-	-	0.43	0.67	-	-	-	-	-	BDL	BDL	0.10	0.10	-	-	-	-	-	-	
DDT	0.70	0.70	11	87,000	59,000	2,700	3,000	89	170	12	12	547	0.70	0.70	-	-	1500	1800	-	-	-	-	-	-	
Diazinon <sup>g</sup>	2.2	4.2	44	-	-	-	-	2.2	4.2	-	-	-	-	-	BDL	BDL	-	-	-	-	-	-	-	-	
Dicamba <sup>g</sup>	0.12	0.12	280	-	-	-	-	0.50	0.79	-	-	-	-	-	BDL	BDL	0.12	0.12	-	-	BDL	BDL	-	-	
Diclofop-methyl	0.079	0.095	40	-	-	-	-	NGR	NGR	-	-	-	-	-	NGR	2.4	3.0	3.6	-	-	0.079	0.095	-	-	
Dieldrin	0.025	0.048	0.33	21	14	0.67	0.74	0.025	0.048	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Dimethoate	0.0028	0.0027	80	-	-	-	-	0.077	0.12	-	-	-	-	-	0.0058	0.0055	0.0028	0.0027	-	-	-	-	-	-	
Dinoseb <sup>g</sup>	1.4	1.7	22	-	-	-	-	2.8	5.5	-	-	-	-	-	BDL	BDL	10	12	-	-	1.4	1.7	-	-	
Diquat	11	21	180	-	-	-	-	11	21	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Diuron	1.9	3.5	350	-	-	-	-	1.9	3.5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor		Overall Guideline		Human							Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit	
				-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse					Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Soil Type		Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type				-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)		(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Endosulfan		0.80	0.0016	240	-	-	-	-	99	190	-	-	-	-	0.80	0.0016	-	-	-	-	-	-	-	-
Endrin		2.4	4.7	6.7	-	-	-	-	2.4	4.7	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Glyphosate		0.054	0.049	670	-	-	-	-	0.95	1.4	-	-	-	-	0.054	0.049	0.23	0.21	-	-	-	-	-	-
Heptachlor epoxide		0.039	0.010	0.29	0.31	0.21	0.010	0.012	0.039	0.076	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Lindane <sup>g</sup>		0.11	0.13	6.7	-	-	-	-	0.31	0.6	-	-	-	-	BDL	BDL	0.11	0.13	-	-	-	-	-	-
Linuron <sup>g</sup>		0.051	0.059	44	-	-	-	-	0.56	1.1	-	-	-	-	0.051	0.059	-	-	-	-	BDL	BDL	-	-
Malathion <sup>g</sup>		0.82	1.3	440	-	-	-	-	0.82	1.3	-	-	-	-	BDL	BDL	-	-	-	-	-	-	-	-
MCPA <sup>g</sup>		0.026	0.025	460	-	-	-	-	0.42	0.66	-	-	-	-	BDL	BDL	0.026	0.025	-	-	BDL	BDL	-	-
Methoxychlor		110	0.32	110	-	-	-	-	NGR	NGR	-	-	-	-	NGR	0.32	-	-	-	-	-	-	-	-
Metolachlor		0.048	0.055	110	-	-	-	-	1.3	2.4	-	-	-	-	0.048	0.055	0.30	0.35	-	-	0.17	0.20	-	-
Metribuzin		0.012	0.014	180	-	-	-	-	7.8	15	-	-	-	-	0.024	0.028	1.9	2.2	-	-	0.012	0.014	-	-
Paraquat (as dichloride)		1.1	2.2	22	-	-	-	-	1.1	2.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Phorate		0.075	0.14	4.4	-	-	-	-	0.075	0.14	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Picloram		0.024	0.022	440	-	-	-	-	0.64	0.94	-	-	-	-	0.024	0.022	0.15	0.14	-	-	-	-	-	-
Simazine <sup>g</sup>		0.033	0.038	29	-	-	-	-	0.14	0.25	-	-	-	-	0.033	0.038	0.033	0.038	-	-	BDL	BDL	-	-
Tebuthiuron <sup>g,j</sup>		0.046	0.046	1600	-	-	-	-	2.5	3.7	0.046	0.046	-	-	BDL	BDL	0.12	0.11	-	-	BDL	BDL	-	-
Terbufos		0.080	0.15	1.1	-	-	-	-	0.080	0.15	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Toxaphene		3.3	4.8	4.8	4,600	3,100	150	170	3.3	6.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Triallate		0.0077	0.0092	520	-	-	-	-	16	31	-	-	-	-	0.0077	0.0092	7.4	8.8	-	-	-	-	-	-
Trifluralin		0.22	0.045	190	-	-	-	-	NGR	NGR	-	-	-	-	0.22	0.045	8.4	10	-	-	-	-	-	-
Other Organics																								
Aniline <sup>g</sup>		0.36	0.60	280	23	20	1.1	0.82	0.36	0.60	-	-	-	-	BDL	BDL	-	-	-	-	-	-	-	-

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological												Other	
Pathway			Direct Soil Contact	Vapour Inhalation				Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life	Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit		
				-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Soil Type		Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse	-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type				-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)		(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Di- <i>n</i> -butyl phthalate		0.54	0.65	1,300	220,000	180,000	9,200	7,400	70	130	-	-	-	-	0.54	0.65	-	-	-	-	-	-	-	-
Dichlorobenzidine		0.60	1.2	12	NGR	NGR	NGR	NGR	0.60	1.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Diethanolamine <sup>j</sup>		2.0	3.5	150	-	-	-	-	2.0	3.5	1,000	1,000	-	-	500,000	45	-	-	-	-	-	-	-	-
Diethylene glycol		10	15	15,000	-	-	-	-	10	15	1,000	1,000	-	-	2,000	65	-	-	-	-	-	-	-	-
Diisopropanolamine		14	17	26,000	-	-	-	-	130	250	360	360	-	-	14	17	-	-	-	-	29	34	-	-
Ethylene glycol		60	62	130,000	NGR	NGR	120,000	86,000	60	68	1,100	1,100	1,700	-	89	62	-	-	-	-	-	-	-	-
Hexachlorobutadiene		0.026	0.0067	390	0.18	0.16	0.0087	0.0067	0.5	0.95	-	-	-	-	0.026	0.031	-	-	-	-	-	-	-	-
Methanol		37	11	16,000	34,000	33,000	2,100	1,400	37	42	1,200	1,200	-	-	300	11	-	-	-	-	-	-	750	750
Methylmethacrylate		36	1.4	56,000	45	40	1.9	1.4	36	49	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Monoethanolamine <sup>j</sup>		20	10	1,500	-	-	-	-	20	40	1,500	1,500	-	-	300,000	10	-	-	-	-	-	-	-	-
MTBE		0.044	0.046	400	1.2	1.1	0.065	0.046	0.044	0.062	-	-	-	-	7.1	6.1	-	-	-	-	-	-	-	-
Nonylphenol + ethoxylates		5.7	5.7	-	-	-	-	-	-	-	5.7	5.7	-	-	NGR	2,000	-	-	-	-	-	-	-	-
Perfluorooctane sulfonate (PFOS)		0.010	0.010	2.1	-	-	-	-	0.070	0.14	10	10	-	0.010	0.010	0.19	0.23	1.7	2.0	1.5	1.8	-	-	-
Phenol		0.0014	0.0012	2,400	14,000	13,000	660	480	1.6	2.3	20	20	-	-	0.0028	0.0024	0.0014	0.0012	-	-	-	-	-	-
Sulfolane		0.18	0.21	350	-	-	-	-	0.18	0.21	210	210	-	-	24	18	-	-	-	-	0.39	0.28	-	-
Triethylene glycol		100	150	150,000	-	-	-	-	100	150	5,000	5,000	-	-	10,000	200	-	-	-	-	-	-	-	-

Notes:

- a. For more information see Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils (AENV, 2011)
- b. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)
- c. Human health direct soil contact guidelines for carcinogenic PAHs are based on B[a]P Total Potency Equivalents (TPE). TPEs are calculated by multiplying the soil concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (WHO/IPCS, 1998) scheme, as follows:

TABLE A-2. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see sections on Land Use and Land Use and Sensitivity Factors). Tier 1 soil guidelines are found in Table 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Carcinogenic PAH	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

- d. The B[a]P Total Potency Equivalents (TPEs) calculated for specific soil samples using Potency Equivalency Factors (PEFs) should be multiplied by an Uncertainty Factor of 3 when evaluating PAH mixtures associated with creosote or coal tar-type environmental releases, prior to evaluating against the human health direct contact soil remediation guideline.
- e. The Index of Additive Cancer Risk (IACR) is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture, as follows:

Fine Soil:

$$\text{IACR} = \frac{[\text{Benz(a)anthracene}]}{6.4} + \frac{[\text{Benzo(b + j)fluoranthene}]}{3.0} + \frac{[\text{Benzo(k)fluoranthene}]}{0.64} + \frac{[\text{Benzo(g, h, i)perylene}]}{130} + \frac{[\text{Benzo(a)pyrene}]}{7.0} + \frac{[\text{Chrysene}]}{40} + \frac{[\text{Dibenz(a, h)anthracene}]}{4.4} + \frac{[\text{Indeno(1,2,3 - c, d)pyrene}]}{51}$$

Coarse Soil:

$$\text{IACR} = \frac{[\text{Benz(a)anthracene}]}{12} + \frac{[\text{Benzo(b + j)fluoranthene}]}{5.8} + \frac{[\text{Benzo(k)fluoranthene}]}{1.2} + \frac{[\text{Benzo(g, h, i)perylene}]}{250} + \frac{[\text{Benzo(a)pyrene}]}{14} + \frac{[\text{Chrysene}]}{78} + \frac{[\text{Dibenz(a, h)anthracene}]}{8.5} + \frac{[\text{Indeno(1,2,3 - c, d)pyrene}]}{98}$$

- f. Overall guideline value for ecological receptors only.
- g. Guideline for protection of aquatic life or irrigation water is below detection limit, groundwater monitoring is required.
- h. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- i. Guideline values adopted directly from CCME (1999 and updates) without change.
- j. Analytical methodology specified in the Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
- k. Eco-contact guidelines from Stantec (2012)
- l. Eco-contact guidelines from Stantec (2008)
- m. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008)
- n. Guideline for protection of aquatic life (fine soil) is based on a groundwater guideline of 0.10 g/L. See Appendix B for more information

BDL - Below detection limit

NGR - no guideline required, calculated value >1,000,000 mg/kg; or for PAH groundwater protection, calculated value results in groundwater concentration greater than solubility.

**TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS**  
**This table must not be used for Tier 1 assessment and remediation**, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological				Other	
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
General and Inorganic Parameters																
pH (in 0.01M CaCl <sub>2</sub> )	6.0-8.5	6.0-8.5	-	-	-	-	-	-	-	6.0-8.5	6.0-8.5	-	-	-	-	-
Cyanide (free)	0.90	0.90	29	-	-	-	-	-	-	0.90	0.90	-	-	-	-	-
Fluoride	200	200	-	-	-	-	-	-	-	200	200	-	-	-	-	-
Sulphur (elemental) <sup>a</sup>	500	500	-	-	-	-	-	-	-	500	500	-	-	-	-	-
Metals																
Antimony	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-
Arsenic (inorganic)	17	17	27	-	-	-	-	-	-	17	17	-	-	-	-	-
Barium (non-barite)	500	500	6,800	-	-	-	-	-	-	500	500	-	-	-	-	-
Barite-barium <sup>b</sup>	10,000	10,000	10,000	-	-	-	-	-	-	200,000	200,000	-	-	-	-	-
Beryllium	5.0	5.0	-	-	-	-	-	-	-	5.0	5.0	-	-	-	-	-
Boron (mg/L in saturated paste extract) <sup>m</sup>	3.3	3.3	7,500	-	-	-	-	65	120	3.3	3.3	-	5.0	5.0	-	-
Cadmium	10	10	14	-	-	-	-	-	-	10	10	54	-	-	-	-
Chromium (hexavalent)	0.40	0.40	-	-	-	-	-	-	-	0.40	0.40	-	-	-	-	-
Chromium (total)	64	64	220	-	-	-	-	-	-	64	64	-	-	-	-	-
Cobalt	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-
Copper	63	63	1,100	-	-	-	-	-	-	63	63	350	-	-	-	-
Lead	140	140	140	-	-	-	-	-	-	300	300	720	-	-	-	-
Mercury (inorganic)	6.6	6.6	6.6	-	-	-	-	-	-	12	12	20	-	-	-	-

TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other	
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit		
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse	
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Molybdenum	4.0	4.0	-	-	-	-	-	-	-	4.0	4.0	-	-	-	-	-	
Nickel	45	45	200	-	-	-	-	-	-	45	45	170	-	-	-	-	
Selenium	1.0	1.0	80	-	-	-	-	-	-	1.0	1.0	-	-	-	-	-	
Silver	20	20	-	-	-	-	-	-	-	20	20	-	-	-	-	-	
Thallium	1.0	1.0	1.0	-	-	-	-	-	-	1.4	1.4	-	-	-	-	-	
Tin	5.0	5.0	-	-	-	-	-	-	-	5.0	5.0	-	-	-	-	-	
Uranium	23	23	23	-	-	-	-	-	-	500	500	-	-	-	-	-	
Vanadium	130	130	-	-	-	-	-	-	-	130	130	260	-	-	-	-	
Zinc	250	250	10,000	-	-	-	-	-	-	250	250	280	-	-	-	-	
Hydrocarbons																	
Benzene	0.046	0.015	360	0.36	0.33	0.021	0.015	0.046	0.078	60	31	-	7.9	0.17	-	-	
Toluene	0.52	0.12	690	1,300	1,200	80	57	0.52	0.95	110	75	-	63,000	0.12	-	-	
Ethylbenzene	0.073	0.14	1,900	2,100	1,900	120	88	0.073	0.14	120	55	-	NGR	540	-	-	
Xylenes	0.99	1.9	510	150	140	9.0	6.6	0.99	1.9	65	95	-	NGR	41	-	-	
Styrene	0.68	0.80	12,000	2,300	2,100	130	94	110	210	-	-	-	0.68	0.80	-	-	
F1	210	24	14,000	710	610	30	24	1,100	2,200	210	210	-	30,000	1,300	800	700	
F2	150	130	6,800	3,600	3,100	160	130	1,500	2,900	150	150	-	30,000	520	1,000	1,000	
F3	1,300	300	15,000	-	-	-	-	-	-	1,300	300	-	-	-	3,500	2,500	
F4	5,600	2,800	21,000	-	-	-	-	-	-	5,600	2,800	-	-	-	10,000	10,000	
Acenaphthene	0.33	0.38	5,500	120,000	99,000	4,800	3,900	NGR	NGR	-	-	-	0.33	0.38	-	-	
Anthracene	1.3	0.0056	27,000	NGR	NGR	780,000	670,000	NGR	NGR	2.5	2.5	-	1.3	0.0056	-	-	

TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human							Ecological				Other		
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Fluoranthene	50	0.055	3,700	NGR	NGR	550,000	480,000	NGR	NGR	50	50	-	NGR	0.055	-	-
Fluorene	0.40	0.34	2,800	270,000	220,000	10,000	8,600	NGR	NGR	-	-	-	0.40	0.34	-	-
Naphthalene	0.014	0.017	2,000	260	220	13	9.6	28	53	-	-	-	0.014	0.017	-	-
Phenanthrene	0.11	0.061	-	-	-	-	-	-	-	-	-	-	0.11	0.061	-	-
Pyrene	2,100	0.15	2,200	NGR	NGR	810,000	730,000	NGR	NGR	-	-	-	NGR	0.15	-	-
Carcinogenic PAHs (as B(a)P TPE) <sup>c</sup>	IACR<1.0 <sup>e</sup> and TPE ≤ 5.6	IACR<1.0 <sup>e</sup> and TPE ≤ 5.6	5.6 <sup>d</sup>	NGR	NGR	NGR	NGR	IACR<1.0 <sup>e</sup>	IACR<1.0 <sup>e</sup>	-	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	-	-	6.4	12	-	-	-	NGR	NGR	-	-
Benzo[b+j]fluoranthene	-	-	-	-	-	-	-	3.0	5.8	-	-	-	-	-	-	-
Benzo[k]fluoranthene	-	-	-	-	-	-	-	0.64	1.2	-	-	-	-	-	-	-
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	130	250	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>f</sup>	20	20	-	-	-	-	-	7.0	14	20	20	-	NGR	NGR	-	-
Chrysene	-	-	-	-	-	-	-	40	78	-	-	-	-	-	-	-
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	4.4	8.5	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	51	98	-	-	-	-	-	-	-
Chlorinated Aliphatics																
Vinyl chloride	0.0083	0.00034	63	0.0090	0.0083	0.00049	0.00034	0.014	0.020	-	-	-	-	-	-	-
1,1-Dichloroethene	0.15	0.021	120	0.50	0.46	0.030	0.021	0.15	0.24	-	-	-	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.0043	0.00019	37	0.0048	0.0043	0.00026	0.00019	0.054	0.093	3.0	3.0	-	0.72	0.081	-	-
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.26	0.018	190	0.46	0.41	0.025	0.018	0.26	0.46	-	-	-	0.69	0.77	-	-
1,2-Dichloroethane	0.025	0.0027	9,100	0.060	0.055	0.0038	0.0027	0.025	0.041	-	-	-	0.12	0.12	-	-

**TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS**  
**This table must not be used for Tier 1 assessment and remediation,** required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human							Ecological				Other		
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Dichloromethane (Methylene chloride)	0.10	0.095	340	4.8	4.4	0.28	0.19	0.21	0.32	-	-	-	0.10	0.095	-	-
Trichloromethane (Chloroform) <sup>n</sup>	0.16	0.030	76	3.2	3.0	0.21	0.15	0.53	0.88	-	-	-	0.16	0.030	-	-
Tetrachloromethane (Carbon tetrachloride)	0.035	0.0015	29	0.038	0.035	0.0021	0.0015	0.037	0.062	-	-	-	0.059	0.062	-	-
Dibromochloromethane	0.91	0.27	800	11	7.8	0.28	0.27	0.91	1.5	-	-	-	-	-	-	-
Chlorinated Aromatics																
Chlorobenzene <sup>g</sup>	0.39	0.018	17,000	0.44	0.39	0.024	0.018	0.61	1.1	-	-	-	BDL	BDL	-	-
1,2-Dichlorobenzene <sup>g</sup>	0.097	0.18	17,000	260	230	14	10	0.097	0.18	-	-	-	BDL	BDL	-	-
1,4-Dichlorobenzene	0.051	0.098	4,400	10	9.1	0.56	0.42	0.051	0.098	-	-	-	0.32	0.38	-	-
1,2,3-Trichlorobenzene	0.26	0.26	51	8.8	6.8	0.30	0.26	1.9	3.6	-	-	-	0.26	0.31	-	-
1,2,4-Trichlorobenzene	0.78	0.23	40	7.6	6.0	0.26	0.23	2.0	3.9	-	-	-	0.78	0.93	-	-
1,3,5-Trichlorobenzene	1.9	0.13	47	4.1	3.2	0.14	0.13	1.9	3.6	-	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.042	0.050	75	27	20	0.88	0.84	3.1	5.9	-	-	-	0.042	0.050	-	-
1,2,3,5-Tetrachlorobenzene	0.37	0.10	16	3.3	2.5	0.10	0.10	0.37	0.70	-	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.19	0.052	4.6	1.7	1.3	0.054	0.052	0.19	0.37	-	-	-	-	-	-	-
Pentachlorobenzene	22	5.2	22	160	140	7.9	6.1	24	47	-	-	-	NGR	5.2	-	-
Hexachlorobenzene	1.8	0.20	3.3	5.4	4.7	0.26	0.20	1.8	3.6	-	-	-	-	-	-	-
2,4-Dichlorophenol	0.0029	0.0034	4,000	170,000	140,000	6,300	5,400	0.018	0.034	-	-	-	0.0029	0.0034	-	-
2,4,6-Trichlorophenol	0.19	0.37	260	11,000	8,200	360	320	0.19	0.37	-	-	-	0.42	0.50	-	-
2,3,4,6-Tetrachlorophenol	0.039	0.047	400	15,000	11,000	480	460	0.16	0.31	-	-	-	0.039	0.047	-	-
Pentachlorophenol	0.025	0.029	130	NGR	NGR	110,000	83,000	6.0	12	11	11	-	0.025	0.029	-	-
Dioxins & Furans <sup>h,i</sup>	0.0000040	0.0000040	0.0000040	-	-	-	-	-	-	-	-	-	-	-	-	-



**TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS**  
**This table must not be used for Tier 1 assessment and remediation,** required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human							Ecological				Other		
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
PCBs	0.085	0.085	0.085	-	-	-	-	-	-	33	33	-	-	-	-	-
Pesticides																
Aldicarb <sup>g</sup>	0.041	0.065	22	-	-	-	-	0.041	0.065	-	-	-	BDL	BDL	-	-
Aldrin	0.24	0.31	0.31	44	30	1.4	1.5	0.24	0.46	-	-	-	-	-	-	-
Atrazine and metabolites	0.0088	0.010	11	-	-	-	-	0.10	0.19	-	-	-	0.0088	0.010	-	-
Azniphos-methyl (Guthion)	0.41	0.75	55	-	-	-	-	0.41	0.75	-	-	-	-	-	-	-
Bromacil <sup>k</sup>	0.0090	0.0090	2,000	-	-	-	-	7.0	10	0.20	0.12	-	0.0090	0.0090	-	-
Bromoxynil	0.044	0.052	11	-	-	-	-	0.18	0.35	-	-	-	0.044	0.052	-	-
Carbaryl <sup>g</sup>	1.9	3.6	220	-	-	-	-	1.9	3.6	-	-	-	BDL	BDL	-	-
Carbofuran <sup>g</sup>	0.68	1.2	220	-	-	-	-	0.68	1.2	-	-	-	BDL	BDL	-	-
Chlorothalonil	0.0085	0.010	330	-	-	-	-	27	53	-	-	-	0.0085	0.010	-	-
Chlorpyrifos <sup>g</sup>	49	95	220	-	-	-	-	49	95	-	-	-	BDL	BDL	-	-
2,4-D <sup>g</sup>	0.43	0.67	400	-	-	-	-	0.43	0.67	-	-	-	BDL	BDL	-	-
DDT	11	11	11	87,000	59,000	2,700	3,000	89	170	12	12	547	-	-	-	-
Diazinon <sup>g</sup>	2.2	4.2	44	-	-	-	-	2.2	4.2	-	-	-	BDL	BDL	-	-
Dicamba <sup>g</sup>	0.50	0.79	280	-	-	-	-	0.50	0.79	-	-	-	BDL	BDL	-	-
Diclofop-methyl	40	2.4	40	-	-	-	-	NGR	NGR	-	-	-	NGR	2.4	-	-
Dieldrin	0.025	0.048	0.33	21	14	0.67	0.74	0.025	0.048	-	-	-	-	-	-	-
Dimethoate	0.0058	0.0055	80	-	-	-	-	0.077	0.12	-	-	-	0.0058	0.0055	-	-
Dinoseb <sup>g</sup>	2.8	5.5	22	-	-	-	-	2.8	5.5	-	-	-	BDL	BDL	-	-
Diquat	11	21	180	-	-	-	-	11	21	-	-	-	-	-	-	-

**TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS**  
**This table must not be used for Tier 1 assessment and remediation**, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human							Ecological				Other		
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Diuron	1.9	3.5	350	-	-	-	-	1.9	3.5	-	-	-	-	-	-	-
Endosulfan	0.80	0.0016	240	-	-	-	-	99	190	-	-	-	0.80	0.0016	-	-
Endrin	2.4	4.7	6.7	-	-	-	-	2.4	4.7	-	-	-	-	-	-	-
Glyphosate	0.054	0.049	670	-	-	-	-	0.95	1.4	-	-	-	0.054	0.049	-	-
Heptachlor epoxide	0.039	0.010	0.29	0.31	0.21	0.010	0.012	0.039	0.076	-	-	-	-	-	-	-
Lindane <sup>g</sup>	0.31	0.60	6.7	-	-	-	-	0.31	0.60	-	-	-	BDL	BDL	-	-
Linuron	0.051	0.059	44	-	-	-	-	0.56	1.1	-	-	-	0.051	0.059	-	-
Malathion <sup>g</sup>	0.82	1.3	440	-	-	-	-	0.82	1.3	-	-	-	BDL	BDL	-	-
MCPA <sup>g</sup>	0.42	0.66	460	-	-	-	-	0.42	0.66	-	-	-	BDL	BDL	-	-
Methoxychlor	110	0.32	110	-	-	-	-	NGR	NGR	-	-	-	NGR	0.32	-	-
Metolachlor	0.048	0.055	110	-	-	-	-	1.3	2.4	-	-	-	0.048	0.055	-	-
Metribuzin	0.024	0.028	180	-	-	-	-	7.8	15	-	-	-	0.024	0.028	-	-
Paraquat (as dichloride)	1.1	2.2	22	-	-	-	-	1.1	2.2	-	-	-	-	-	-	-
Phorate	0.075	0.14	4.4	-	-	-	-	0.075	0.14	-	-	-	-	-	-	-
Picloram	0.024	0.022	440	-	-	-	-	0.64	0.94	-	-	-	0.024	0.022	-	-
Simazine	0.033	0.038	29	-	-	-	-	0.14	0.25	-	-	-	0.033	0.038	-	-
Tebuthiuron <sup>g,l</sup>	0.046	0.046	1,600	-	-	-	-	2.5	3.7	0.046	0.046	-	BDL	BDL	-	-
Terbufos	0.080	0.15	1.1	-	-	-	-	0.080	0.15	-	-	-	-	-	-	-
Toxaphene	3.3	4.8	4.8	4,600	3,100	150	170	3.3	6.3	-	-	-	-	-	-	-
Triallate	0.0077	0.0092	520	-	-	-	-	16	31	-	-	-	0.0077	0.0092	-	-
Trifluralin	0.22	0.045	190	-	-	-	-	NGR	NGR	-	-	-	0.22	0.045	-	-

TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological				Other	
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Other Organics																
Aniline <sup>9</sup>	0.36	0.60	280	23	20	1.1	0.82	0.36	0.60	-	-	-	BDL	BDL	-	-
Di- <i>n</i> -butyl phthalate	0.54	0.65	1,300	220,000	180,000	9,200	7,400	70	130	-	-	-	0.54	0.65	-	-
Dichlorobenzidine	0.60	1.2	12	NGR	NGR	NGR	NGR	0.6	1.2	-	-	-	-	-	-	-
Diethanolamine <sup>i</sup>	2.0	3.5	150	-	-	-	-	2.0	3.5	1,000	1,000	-	500,000	45	-	-
Diethylene glycol	10	15	15,000	-	-	-	-	10	15	1,000	1,000	-	2,000	65	-	-
Diisopropanolamine	14	17	26,000	-	-	-	-	130	250	360	360	-	14	17	-	-
Ethylene glycol	60	62	130,000	NGR	NGR	120,000	86,000	60	68	1,100	1,100	1,700	89	62	-	-
Hexachlorobutadiene	0.026	0.0067	390	0.18	0.16	0.0087	0.0067	0.5	0.95	-	-	-	0.026	0.031	-	-
Methanol	37	11	16,000	34,000	33,000	2,100	1,400	37	42	1,200	1,200	-	300	11	750	750
Methylmethacrylate	36	1.4	56,000	45	40	1.9	1.4	36	49	-	-	-	-	-	-	-
Monoethanolamine <sup>i</sup>	20	10	1,500	-	-	-	-	20	40	1,500	1,500	-	300,000	10	-	-
MTBE	0.044	0.046	400	1.2	1.1	0.065	0.046	0.044	0.062	-	-	-	7.1	6.1	-	-
Nonylphenol + ethoxylates	5.7	5.7	-	-	-	-	-	-	-	5.7	5.7	-	NGR	2,000	-	-
Perfluorooctane sulfonate (PFOS)	0.070	0.14	2.1	-	-	-	-	0.070	0.14	10	10	-	0.19	0.23	-	-
Phenol	0.0028	0.0024	2,400	14,000	13,000	660	480	1.6	2.3	20	20	-	0.0028	0.0024	-	-
Sulfolane	0.18	0.21	350	-	-	-	-	0.18	0.21	210	210	-	24	18	-	-
Triethylene glycol	100	150	150,000	-	-	-	-	100	150	5,000	5,000	-	10,000	200	-	-

Notes:

- a. For more information see Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils (AENV, 2011)
- b. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)

TABLE A-3. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

- c. Human health direct soil contact guidelines for carcinogenic PAHs are based on B[a]P Total Potency Equivalents (TPE). TPEs are calculated by multiplying the soil concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (WHO/IPCS, 1998) scheme, as follows:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

- d. The B[a]P Total Potency Equivalents (TPEs) calculated for specific soil samples using Potency Equivalency Factors (PEFs) should be multiplied by an Uncertainty Factor of 3 when evaluating PAH mixtures associated with creosote or coal tar-type environmental releases, prior to evaluating against the human health direct contact soil remediation guideline.
- e. The Index of Additive Cancer Risk (IACR) is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture. For example, the IACR for coarse textured soil would be calculated as follows:

Fine Soil:

$$IACR = \frac{[Benz(a)anthracene]}{6.4} + \frac{[Benzo(b + j)fluoranthene]}{3.0} + \frac{[Benzo(k)fluoranthene]}{0.64} + \frac{[Benzo(g, h, i)perylene]}{130} + \frac{[Benzo(a)pyrene]}{7.0} + \frac{[Chrysene]}{40} + \frac{[Dibenz(a, h)anthracene]}{4.4} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{51}$$

Coarse Soil:

$$IACR = \frac{[Benz(a)anthracene]}{12} + \frac{[Benzo(b + j)fluoranthene]}{5.8} + \frac{[Benzo(k)fluoranthene]}{1.2} + \frac{[Benzo(g, h, i)perylene]}{250} + \frac{[Benzo(a)pyrene]}{14} + \frac{[Chrysene]}{78} + \frac{[Dibenz(a, h)anthracene]}{8.5} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{98}$$

- f. Overall guideline value for ecological receptors only.
- g. Guideline for protection of aquatic life is below detection limit, groundwater monitoring is required.
- h. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- i. Guideline values adopted directly from CCME (1999 and updates) without change.
- j. Analytical methodology specified in the Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
- k. Eco-contact guidelines from Stantec (2012)
- l. Eco-contact guidelines from Stantec (2008)
- m. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008)
- n. Guideline for protection of aquatic life (fine soil) is based on a groundwater guideline of 0.10 g/L. See Appendix B for more information

BDL - Below detection limit

NGR - no guideline required, calculated value >1,000,000 mg/kg; or for PAH groundwater protection, calculated value results in groundwater concentration greater than solubility

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human					Ecological					Other	
Pathway			Direct Soil Contact	Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
General and Inorganic Parameters																
pH (in 0.01M CaCl <sub>2</sub> )	6.0-8.5	6.0-8.5	-	-	-	-	-	-	6.0-8.5	6.0-8.5	-	-	-	-	-	-
Cyanide (free)	8.0	8.0	110	-	-	-	-	-	8.0	8.0	-	-	-	-	-	-
Fluoride	2,000	2,000	-	-	-	-	-	-	2,000	2,000	-	-	-	-	-	-
Sulphur (elemental) <sup>a</sup>	500	500	-	-	-	-	-	-	500	500	-	-	-	-	-	-
Metals																
Antimony	40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-
Arsenic (inorganic)	26	26	35	-	-	-	-	-	26	26	-	-	-	-	-	-
Barium (non-barite)	2,000	2,000	10,000	-	-	-	-	96,000	2,000	2,000	-	-	-	-	-	-
Barite-barium <sup>b</sup>	15,000	15,000	15,000	-	-	-	-	140,000	200,000	200,000	-	-	-	140,000	-	-
Beryllium	8.0	8.0	-	-	-	-	-	-	8.0	8.0	-	-	-	-	-	-
Boron (mg/L in saturated paste extract) <sup>m</sup>	5.0	5.0	11,000	-	-	65	120	110,000	7.9	7.9	-	5.0	5.0	46	-	-
Cadmium	22	22	49	-	-	-	-	-	22	22	200	-	-	-	-	-
Chromium (hexavalent)	1.4	1.4	-	-	-	-	-	-	1.4	1.4	-	-	-	-	-	-
Chromium (total)	87	87	630	-	-	-	-	-	87	87	-	-	-	-	-	-
Cobalt	300	300	-	-	-	-	-	-	300	300	-	-	-	-	-	-
Copper	91	91	4,000	-	-	-	-	-	91	91	350	-	-	-	-	-
Lead	260	260	260	-	-	-	-	-	600	600	830	-	-	-	-	-
Mercury (inorganic)	24	24	24	-	-	-	-	-	50	50	52	-	-	-	-	-

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other	
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse	
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Molybdenum	40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-	
Nickel	89	89	310	-	-	-	-	2,500	89	89	240	-	-	290	-	-	
Selenium	2.9	2.9	130	-	-	-	-	1,100	2.9	2.9	-	-	-	5.0	-	-	
Silver	40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-	
Thallium	1.0	1.0	1.0	-	-	-	-	-	3.6	3.6	-	-	-	-	-	-	
Tin	300	300	-	-	-	-	-	-	300	300	-	-	-	-	-	-	
Uranium	33	33	33	-	-	-	-	-	2000	2000	-	-	-	-	-	-	
Vanadium	130	130	-	-	-	-	-	-	130	130	260	-	-	-	-	-	
Zinc	410	410	16,000	-	-	-	-	140,000	450	450	410	-	-	2,900	-	-	
Hydrocarbons																	
Benzene	0.046	0.078	550	2.2	0.19	0.046	0.078	5,100	310	180	-	7.9	0.17	440	-	-	
Toluene	0.52	0.12	1,100	7,900	700	0.52	0.95	9,900	330	250	-	63,000	0.12	1,100	-	-	
Ethylbenzene	0.073	0.14	2,900	13,000	1,100	0.073	0.14	27,000	430	300	-	NGR	540	790	-	-	
Xylenes	0.99	1.9	780	950	80	0.99	1.9	7,300	230	350	-	NGR	41	930	-	-	
Styrene	0.68	0.80	18,000	14,000	1,100	110	210	170,000	-	-	-	0.68	0.80	-	-	-	
F1	320	270	22,000	4,500	270	1,100	2,200	30,000	320	320	-	30,000	1,300	3,000	800	700	
F2	260	260	10,000	23,000	1,500	1,500	2,900	30,000	260	260	-	30,000	520	2,100	1,000	1,000	
F3	2,500	1,700	23,000	-	-	-	-	30,000	2,500	1,700	-	-	-	4,300	5,000	3,500	
F4	6,600	3,300	30,000	-	-	-	-	30,000	6,600	3,300	-	-	-	30,000	10,000	10,000	
Acenaphthene	0.33	0.38	8,300	770,000	43,000	NGR	NGR	79,000	-	-	-	0.33	0.38	-	-	-	

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human				Ecological					Other			
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Anthracene			1.3	0.0056	41,000	NGR	NGR	NGR	NGR	390,000	32	32	-	1.3	0.0056	36	-	-
Fluoranthene			180	0.055	5,600	NGR	NGR	NGR	NGR	53,000	180	180	-	NGR	0.055	720	-	-
Fluorene			0.40	0.34	4,300	NGR	91,000	NGR	NGR	40,000	-	-	-	0.40	0.34	-	-	-
Naphthalene			0.014	0.017	3,000	1,600	110	28	53	29,000	-	-	-	0.014	0.017	-	-	-
Phenanthrene			0.11	0.061	-	-	-	-	-	-	-	-	-	0.11	0.061	-	-	-
Pyrene			3,200	0.15	3,300	NGR	NGR	NGR	NGR	31,000	-	-	-	NGR	0.15	-	-	-
Carcinogenic PAHs (as B(a)P TPE) <sup>c</sup>			IACR<1.0 <sup>e</sup> and TPE ≤ 8.5	IACR<1.0 <sup>e</sup> and TPE ≤ 8.5	8.5 <sup>d</sup>	NGR	NGR	IACR<1.0 <sup>e</sup>	IACR<1.0 <sup>e</sup>	80	-	-	-	-	-	-	-	-
Benz[a]anthracene			-	-	-	-	-	6.4	12	-	-	-	-	NGR	NGR	-	-	-
Benzo[b+]]fluoranthene			-	-	-	-	-	3.0	5.8	-	-	-	-	-	-	-	-	-
Benzo[k]fluoranthene			-	-	-	-	-	0.64	1.2	-	-	-	-	-	-	-	-	-
Benzo[g,h,i]perylene			-	-	-	-	-	130	250	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>f</sup>			72	72	-	-	-	7.0	14	-	72	72	-	NGR	NGR	290	-	-
Chrysene			-	-	-	-	-	40	78	-	-	-	-	-	-	-	-	-
Dibenz[a,h]anthracene			-	-	-	-	-	4.4	8.5	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene			-	-	-	-	-	51	98	-	-	-	-	-	-	-	-	-
Chlorinated Aliphatics																		
Vinyl chloride			0.014	0.0043	95	0.055	0.0043	0.014	0.020	900	-	-	-	-	-	-	-	-
1,1-Dichloroethene			0.15	0.24	180	3.1	0.27	0.15	0.24	1,700	-	-	-	-	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)			0.030	0.0023	56	0.030	0.0023	0.054	0.093	530	50	50	-	0.72	0.081	43	-	-

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse	
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.26	0.22	290	2.9	0.22	0.26	0.46	2,700	-	-	-	0.69	0.77	-	-	-	
1,2-Dichloroethane	0.025	0.033	14,000	0.37	0.033	0.025	0.041	130,000	-	-	-	0.12	0.12	-	-	-	
Dichloromethane (Methylene chloride)	0.10	0.095	510	30	2.4	0.21	0.32	4,900	-	-	-	0.10	0.095	-	-	-	
Trichloromethane (Chloroform) <sup>n</sup>	0.16	0.030	110	20	1.8	0.53	0.88	1,100	-	-	-	0.16	0.030	-	-	-	
Tetrachloromethane (Carbon tetrachloride)	0.037	0.018	43	0.24	0.018	0.037	0.062	410	-	-	-	0.059	0.062	-	-	-	
Dibromochloromethane	0.91	1.5	1,200	76	2.5	0.91	1.5	11,000	-	-	-	-	-	-	-	-	
Chlorinated Aromatics																	
Chlorobenzene <sup>g</sup>	0.61	0.22	26,000	2.7	0.22	0.61	1.1	240,000	-	-	-	BDL	BDL	-	-	-	
1,2-Dichlorobenzene <sup>g</sup>	0.097	0.18	26,000	1,700	130	0.097	0.18	240,000	-	-	-	BDL	BDL	-	-	-	
1,4-Dichlorobenzene	0.051	0.098	6,600	65	5.0	0.051	0.098	63,000	-	-	-	0.32	0.38	-	-	-	
1,2,3-Trichlorobenzene	0.26	0.31	77	58	2.7	1.9	3.6	730	-	-	-	0.26	0.31	-	-	-	
1,2,4-Trichlorobenzene	0.78	0.93	61	51	2.4	2.0	3.9	570	-	-	-	0.78	0.93	-	-	-	
1,3,5-Trichlorobenzene	1.9	1.3	72	27	1.3	1.9	3.6	670	-	-	-	-	-	-	-	-	
1,2,3,4-Tetrachlorobenzene	0.042	0.050	110	190	7.9	3.1	5.9	1,100	-	-	-	0.042	0.050	-	-	-	
1,2,3,5-Tetrachlorobenzene	0.37	0.70	25	23	1.0	0.37	0.70	230	-	-	-	-	-	-	-	-	
1,2,4,5-Tetrachlorobenzene	0.19	0.37	7.0	12	0.49	0.19	0.37	66	-	-	-	-	-	-	-	-	
Pentachlorobenzene	24	5.2	34	1000	70	24	47	320	-	-	-	NGR	5.2	-	-	-	
Hexachlorobenzene	1.8	2.3	5.0	34	2.3	1.8	3.6	47	-	-	-	-	-	-	-	-	
2,4-Dichlorophenol	0.0029	0.0034	6,100	NGR	57,000	0.018	0.034	57,000	-	-	-	0.0029	0.0034	-	-	-	



TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human						Ecological						Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration		Direct Soil Contact		Nutrient/ Energy Cycling Check		Protection of Freshwater Aquatic Life		Off-Site Migration		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse				
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-				
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)				
2,4,6-Trichlorophenol	0.19	0.37	400	71,000	3,300	0.19	0.37	3,700	-	-	-	0.42	0.50	-	-	-				
2,3,4,6-Tetrachlorophenol	0.039	0.047	610	110,000	4,400	0.16	0.31	5,700	-	-	-	0.039	0.047	-	-	-				
Pentachlorophenol	0.025	0.029	200	NGR	950,000	6.0	12	1,900	28	28	-	0.025	0.029	160	-	-				
Dioxins & Furans <sup>h,i</sup>	0.0000040	0.0000040	0.0000040	-	-	-	-	0.0000040	-	-	-	-	-	-	-	-				
PCBs	0.13	0.13	0.13	-	-	-	-	1.2	33	33	-	-	-	470	-	-				
Pesticides																				
Aldicarb <sup>g</sup>	0.041	0.065	34	-	-	0.041	0.065	320	-	-	-	BDL	BDL	-	-	-				
Aldrin	0.24	0.46	0.47	330	13	0.24	0.46	4.4	-	-	-	-	-	-	-	-				
Atrazine and metabolites	0.0088	0.010	17	-	-	0.10	0.19	160	-	-	-	0.0088	0.010	-	-	-				
Azniphos-methyl (Guthion)	0.41	0.75	84	-	-	0.41	0.75	790	-	-	-	-	-	-	-	-				
Bromacil <sup>k</sup>	0.0090	0.0090	3,500	-	-	7.0	10	30,000	0.49	0.20	-	0.0090	0.0090	1.7	-	-				
Bromoxynil	0.044	0.052	17	-	-	0.18	0.35	160	-	-	-	0.044	0.052	-	-	-				
Carbaryl <sup>g</sup>	1.9	3.6	340	-	-	1.9	3.6	3,200	-	-	-	BDL	BDL	-	-	-				
Carbofuran <sup>g</sup>	0.68	1.2	340	-	-	0.68	1.2	3,200	-	-	-	BDL	BDL	-	-	-				
Chlorothalonil	0.0085	0.010	500	-	-	27	53	4,800	-	-	-	0.0085	0.010	-	-	-				
Chlorpyrifos <sup>g</sup>	49	95	340	-	-	49	95	3,200	-	-	-	BDL	BDL	-	-	-				
2,4-D <sup>g</sup>	0.43	0.67	610	-	-	0.43	0.67	5,700	-	-	-	BDL	BDL	-	-	-				
DDT	12	12	17	660,000	25,000	89	170	160	12	12	547	-	-	170	-	-				
Diazinon <sup>g</sup>	2.2	4.2	67	-	-	2.2	4.2	630	-	-	-	BDL	BDL	-	-	-				
Dicamba <sup>g</sup>	0.50	0.79	420	-	-	0.50	0.79	4,000	-	-	-	BDL	BDL	-	-	-				

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human						Ecological						Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration		Direct Soil Contact		Nutrient/ Energy Cycling Check		Protection of Freshwater Aquatic Life		Off-Site Migration		Management Limit	
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse		
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-		
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)		
Diclofop-methyl			61	2.4	61	-	-	NGR	NGR	570	-	-	-	NGR	2.4	-	-	-		
Dieldrin			0.025	0.048	0.50	160	6.1	0.025	0.048	4.7	-	-	-	-	-	-	-	-		
Dimethoate			0.0058	0.0055	120	-	-	0.077	0.12	1,100	-	-	-	0.0058	0.0055	-	-	-		
Dinoseb <sup>9</sup>			2.8	5.5	34	-	-	2.8	5.5	320	-	-	-	BDL	BDL	-	-	-		
Diquat			11	21	270	-	-	11	21	2,500	-	-	-	-	-	-	-	-		
Diuron			1.9	3.5	520	-	-	1.9	3.5	4,900	-	-	-	-	-	-	-	-		
Endosulfan			0.80	0.0016	370	-	-	99	190	3,400	-	-	-	0.80	0.0016	-	-	-		
Endrin			2.4	4.7	10	-	-	2.4	4.7	96	-	-	-	-	-	-	-	-		
Glyphosate			0.054	0.049	1,000	-	-	0.95	1.4	9,500	-	-	-	0.054	0.049	-	-	-		
Heptachlor epoxide			0.039	0.076	0.44	2.4	0.094	0.039	0.076	4.1	-	-	-	-	-	-	-	-		
Lindane <sup>9</sup>			0.31	0.60	10	-	-	0.31	0.60	95	-	-	-	BDL	BDL	-	-	-		
Linuron			0.051	0.059	67	-	-	0.56	1.1	630	-	-	-	0.051	0.059	-	-	-		
Malathion <sup>9</sup>			0.82	1.3	670	-	-	0.82	1.3	6,300	-	-	-	BDL	BDL	-	-	-		
MCPA <sup>9</sup>			0.42	0.66	690	-	-	0.42	0.66	160	-	-	-	BDL	BDL	-	-	-		
Methoxychlor			170	0.32	170	-	-	NGR	NGR	1,600	-	-	-	NGR	0.32	-	-	-		
Metolachlor			0.048	0.055	170	-	-	1.3	2.4	1,600	-	-	-	0.048	0.055	-	-	-		
Metribuzin			0.024	0.028	280	-	-	7.8	15	2,600	-	-	-	0.024	0.028	-	-	-		
Paraquat (as dichloride)			1.1	2.2	34	-	-	1.1	2.2	320	-	-	-	-	-	-	-	-		
Phorate			0.075	0.14	6.7	-	-	0.075	0.14	63	-	-	-	-	-	-	-	-		
Picloram			0.024	0.022	670	-	-	0.64	0.94	6,300	-	-	-	0.024	0.022	-	-	-		

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other	
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse	
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Simazine	0.033	0.038	44	-	-	0.14	0.25	410	-	-	-	0.033	0.038	-	-	-	
Tebuthiuron <sup>g,l</sup>	0.60	0.60	2,400	-	-	2.5	3.7	22,000	0.60	0.60	-	BDL	BDL	0.66	-	-	
Terbufos	0.080	0.15	1.7	-	-	0.080	0.15	16	-	-	-	-	-	-	-	-	
Toxaphene	3.3	6.3	7.3	36,000	1,400	3.3	6.3	69	-	-	-	-	-	-	-	-	
Triallate	0.0077	0.0092	790	-	-	16	31	7,400	-	-	-	0.0077	0.0092	-	-	-	
Trifluralin	0.22	0.045	290	-	-	NGR	NGR	2,700	-	-	-	0.22	0.045	-	-	-	
Other Organics																	
Aniline <sup>g</sup>	0.36	0.60	430	140	9.8	0.36	0.60	4,000	-	-	-	BDL	BDL	-	-	-	
Di- <i>n</i> -butyl phthalate	0.54	0.65	1,900	NGR	82,000	70	130	19,000	-	-	-	0.54	0.65	-	-	-	
Dichlorobenzidine	0.60	1.2	18	NGR	NGR	0.60	1.2	170	-	-	-	-	-	-	-	-	
Diethanolamine <sup>l</sup>	2.0	3.5	200	-	-	2.0	3.5	2,000	2,000	2,000	-	500,000	45	15,000	-	-	
Diethylene glycol	10	15	20,000	-	-	10	15	200,000	1,500	1,500	-	2,000	65	15,000	-	-	
Diisopropanolamine	14	17	39,000	-	-	130	250	370,000	750	750	-	14	17	5,100	-	-	
Ethylene glycol	60	62	200,000	NGR	NGR	60	68	NGR	1,800	1,800	2,000	89	62	16,000	-	-	
Hexachlorobutadiene	0.026	0.031	590	1.2	0.078	0.50	0.95	5,600	-	-	-	0.026	0.031	-	-	-	
Methanol	37	11	24,000	210,000	18,000	37	42	-	1,600	1,600	-	300	11	-	750	750	
Methylmethacrylate	36	17	85,000	280	17	36	49	800,000	-	-	-	-	-	-	-	-	
Monoethanolamine <sup>l</sup>	20	10	2,000	-	-	20	40	20,000	1,500	1,500	-	300,000	10	20,000	-	-	
MTBE	0.044	0.062	610	7.4	0.57	0.044	0.062	5,700	-	-	-	7.1	6.1	-	-	-	
Nonylphenol + ethoxylates	14	14	-	-	-	-	-	-	14	14	-	NGR	2,000	82	-	-	

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other				
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration		Direct Soil Contact		Nutrient/ Energy Cycling Check		Protection of Freshwater Aquatic Life		Off-Site Migration		Management Limit	
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse		
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-		
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)		
Perfluorooctane sulfonate (PFOS)			0.070	0.14	3.2	-	-	0.070	0.14	30	60	60	-	0.19	0.23	140	-	-		
Phenol			0.0028	0.0024	3,700	90,000	5,800	1.6	2.3	34,000	130	130	-	0.0028	0.0024	290	-	-		
Sulfolane			0.18	0.21	540	-	-	0.18	0.21	5,000	430	430	-	24	18	3,000	-	-		
Triethylene glycol			100	150	200,000	-	-	100	150	NGR	7,000	7,000	-	10,000	200	70,000	-	-		

- Notes:
- a. For more information see Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils (AENV, 2011)
  - b. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)
  - c. Human health direct soil contact guidelines for carcinogenic PAHs are based on B[a]P Total Potency Equivalents (TPE). TPEs are calculated by multiplying the soil concentration of individual carcinogenic PAHs by a the standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (WHO/IPCS, 1998) scheme, as follows:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1

TABLE A-4. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation, required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Indeno[1,2,3-c,d]pyrene	0.1
-------------------------	-----

- d. The B[a]P Total Potency Equivalents (TPEs) calculated for specific soil samples using Potency Equivalency Factors (PEFs) should be multiplied by an Uncertainty Factor of 3 when evaluating PAH mixtures associated with creosote or coal tar-type environmental releases, prior to evaluating against the human health direct contact soil remediation guideline.
- e. The Index of Additive Cancer Risk (IACR) is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture, as follows:

Fine Soils:

$$IACR = \frac{[Benz(a)anthracene]}{6,4} + \frac{[Benzo(b + j)fluoranthene]}{3,0} + \frac{[Benzo(k)fluoranthene]}{0,64} + \frac{[Benzo(g, h, i)perylene]}{130} + \frac{[Benzo(a)pyrene]}{7,0} + \frac{[Chrysene]}{40} + \frac{[Dibenz(a, h)anthracene]}{4,4} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{51}$$

Coarse Soils:

$$IACR = \frac{[Benz(a)anthracene]}{12} + \frac{[Benzo(b + j)fluoranthene]}{5,8} + \frac{[Benzo(k)fluoranthene]}{1,2} + \frac{[Benzo(g, h, i)perylene]}{250} + \frac{[Benzo(a)pyrene]}{14} + \frac{[Chrysene]}{78} + \frac{[Dibenz(a, h)anthracene]}{8,5} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{98}$$

- f. Overall guideline value for ecological receptors only.
- g. Guideline for protection of aquatic life is below detection limit, groundwater monitoring is required.
- h. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- i. Guideline values adopted directly from CCME (1999 and updates) without change.
- j. Analytical methodology specified in the Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
- k. Eco-contact guidelines from Stantec (2012).
- l. Eco-contact guidelines from Stantec (2008.)
- m. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008).
- n. Guideline for protection of aquatic life (fine soil) is based on a groundwater guideline of 0.10 g/L. See Appendix B for more information

BDL - Below detection limit

NGR - no guideline required, calculated value >1,000,000 mg/kg; or for PAH groundwater protection, calculated value results in groundwater concentration greater than solubility

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
General and Inorganic Parameters																		
pH (in 0.01M CaCl <sub>2</sub> )			6.0-8.5	6.0-8.5	-	-	-	-	-	-	6.0-8.5	6.0-8.5	-	-	-	-	-	-
Cyanide (free)			8.0	8.0	420	-	-	-	-	-	8.0	8.0	-	-	-	-	-	-
Fluoride			2,000	2,000	-	-	-	-	-	-	2,000	2,000	-	-	-	-	-	-
Sulphur (elemental) <sup>a</sup>			500	500	-	-	-	-	-	-	500	500	-	-	-	-	-	-
Metals																		
Antimony			40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-
Arsenic (inorganic)			26	26	35	-	-	-	-	-	26	26	-	-	-	-	-	-
Barium (non-barite)			2,000	2,000	130,000	-	-	-	-	96,000	2,000	2,000	-	-	-	-	-	-
Barite-barium <sup>b</sup>			140,000	140,000	140,000	-	-	-	-	140,000	200,000	200,000	-	-	-	140,000	-	-
Beryllium			8.0	8.0	-	-	-	-	-	-	8.0	8.0	-	-	-	-	-	-
Boron (mg/L in saturated paste extract) <sup>m</sup>			5.0	5.0	230,000	-	-	65	120	110,000	7.9	7.9	-	5.0	5.0	46	-	-
Cadmium			22	22	2,100	-	-	-	-	-	22	22	200	-	-	-	-	-
Chromium (hexavalent)			1.4	1.4	-	-	-	-	-	-	1.4	1.4	-	-	-	-	-	-
Chromium (total)			87	87	2,300	-	-	-	-	-	87	87	-	-	-	-	-	-
Cobalt			300	300	-	-	-	-	-	-	300	300	-	-	-	-	-	-
Copper			91	91	16,000	-	-	-	-	-	91	91	350	-	-	-	-	-
Lead			600	600	8,200	-	-	-	-	-	600	600	830	-	-	-	-	-
Mercury (inorganic)			50	50	99	-	-	-	-	-	50	50	52	-	-	-	-	-

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Molybdenum			40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-
Nickel			89	89	5,100	-	-	-	-	2,500	89	89	240	-	-	290	-	-
Selenium			2.9	2.9	4,050	-	-	-	-	1,100	2.9	2.9	-	-	-	5.0	-	-
Silver			40	40	-	-	-	-	-	-	40	40	-	-	-	-	-	-
Thallium			1.0	1.0	1.0	-	-	-	-	-	3.6	3.6	-	-	-	-	-	-
Tin			300	300	-	-	-	-	-	-	300	300	-	-	-	-	-	-
Uranium			300	300	510	-	-	-	-	300	2,000	2,000	-	-	-	7,100	-	-
Vanadium			130	130	-	-	-	-	-	-	130	130	260	-	-	-	-	-
Zinc			410	410	270,000	-	-	-	-	140,000	450	450	410	-	-	2,900	-	-
Hydrocarbons																		
Benzene			0.046	0.078	550	2.2	0.19	0.046	0.078	5,100	310	180	-	7.9	0.17	440	-	-
Toluene			0.52	0.12	16,000	7,900	700	0.52	0.95	9,900	330	250	-	63,000	0.12	1,100	-	-
Ethylbenzene			0.073	0.14	44,000	13,000	1,100	0.073	0.14	27,000	430	300	-	NGR	540	790	-	-
Xylenes			0.99	1.9	12,000	950	80	0.99	1.9	7,300	230	350	-	NGR	41	930	-	-
Styrene			0.68	0.80	270,000	14,000	1,100	110	210	170,000	-	-	-	0.68	0.80	-	-	-
F1			320	270	30,000	4,500	270	1,100	2,200	30,000	320	320	-	30,000	1,300	3,000	800	700
F2			260	260	30,000	23,000	1,500	1,500	2,900	30,000	260	260	-	30,000	520	2,100	1,000	1,000
F3			2,500	1,700	30,000	-	-	-	-	30,000	2,500	1,700	-	-	-	4,300	5,000	3,500
F4			6,600	3,300	30,000	-	-	-	-	30,000	6,600	3,300	-	-	-	30,000	10,000	10,000

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor	Overall Guideline		Human						Ecological						Other	
Pathway			Direct Soil Contact	Vapour Inhalation	Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact	Nutrient/ Energy Cycling Check		Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Acenaphthene	0.33	0.38	87,000	770,000	43,000	NGR	NGR	79,000	-	-	-	0.33	0.38	-	-	-
Anthracene	1.3	0.0056	430,000	NGR	NGR	NGR	NGR	390,000	32	32	-	1.3	0.0056	36	-	-
Fluoranthene	180	0.055	58,000	NGR	NGR	NGR	NGR	53,000	180	180	-	NGR	0.055	720	-	-
Fluorene	0.40	0.34	53,000	NGR	91,000	NGR	NGR	40,000	-	-	-	0.40	0.34	-	-	-
Naphthalene	0.014	0.017	45,000	1,600	110	28	53	29,000	-	-	-	0.014	0.017	-	-	-
Phenanthrene	0.11	0.061	-	-	-	-	-	-	-	-	-	0.11	0.061	-	-	-
Pyrene	30,000	0.15	40,000	NGR	NGR	NGR	NGR	31,000	-	-	-	NGR	0.15	-	-	-
Carcinogenic PAHs (as B(a)P TPE) °	IACR<1.0 ° and TPE ≤ 23	IACR<1.0 ° and TPE ≤ 23	23 d	NGR	NGR	IACR<1.0 °	IACR<1.0 °	80	-	-	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	6.4	12	-	-	-	-	NGR	NGR	-	-	-
Benzo[b+j]fluoranthene	-	-	-	-	-	3.0	5.8	-	-	-	-	-	-	-	-	-
Benzo[k]fluoranthene	-	-	-	-	-	0.64	1.2	-	-	-	-	-	-	-	-	-
Benzo[g,h,i]perylene	-	-	-	-	-	130	250	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene f	72	72	-	-	-	7.0	14	-	72	72	-	NGR	NGR	290	-	-
Chrysene	-	-	-	-	-	40	78	-	-	-	-	-	-	-	-	-
Dibenz[a,h]anthracene	-	-	-	-	-	4.4	8.5	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	51	98	-	-	-	-	-	-	-	-	-
Chlorinated Aliphatics																
Vinyl chloride	0.014	0.0043	95	0.055	0.0043	0.014	0.020	900	-	-	-	-	-	-	-	-
1,1-Dichloroethene	0.15	0.24	2,700	3.1	0.27	0.15	0.24	1,700	-	-	-	-	-	-	-	-



TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Trichloroethene (Trichloroethylene, TCE)			0.030	0.0023	960	0.030	0.0023	0.054	0.093	530	50	50	-	0.72	0.081	43	-	-
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)			0.26	0.22	4,300	2.9	0.22	0.26	0.46	2,700	-	-	-	0.69	0.77	-	-	-
1,2-Dichloroethane			0.025	0.033	14,000	0.37	0.033	0.025	0.041	130,000	-	-	-	0.12	0.12	-	-	-
Dichloromethane (Methylene chloride)			0.10	0.095	8,500	30	2.4	0.21	0.32	4,900	-	-	-	0.10	0.095	-	-	-
Trichloromethane (Chloroform) <sup>n</sup>			0.16	0.030	2,400	20	1.8	0.53	0.88	1,100	-	-	-	0.16	0.030	-	-	-
Tetrachloromethane (Carbon tetrachloride)			0.037	0.018	650	0.24	0.018	0.037	0.062	400	-	-	-	0.059	0.062	-	-	-
Dibromochloromethane			0.91	1.5	18,000	76	2.5	0.91	1.5	11,000	-	-	-	-	-	-	-	-
Chlorinated Aromatics																		
Chlorobenzene <sup>9</sup>			0.61	0.22	390,000	2.7	0.22	0.61	1.1	240,000	-	-	-	BDL	BDL	-	-	-
1,2-Dichlorobenzene <sup>9</sup>			0.097	0.18	390,000	1700	130	0.097	0.18	240,000	-	-	-	BDL	BDL	-	-	-
1,4-Dichlorobenzene			0.051	0.098	100,000	65	5.0	0.051	0.098	63,000	-	-	-	0.32	0.38	-	-	-
1,2,3-Trichlorobenzene			0.26	0.31	1,200	58	2.7	1.9	3.6	730	-	-	-	0.26	0.31	-	-	-
1,2,4-Trichlorobenzene			0.78	0.93	1,100	51	2.4	2.0	3.9	570	-	-	-	0.78	0.93	-	-	-
1,3,5-Trichlorobenzene			1.9	1.3	1,100	27	1.3	1.9	3.6	670	-	-	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene			0.042	0.05	540	190	7.9	3.1	5.9	1,100	-	-	-	0.042	0.05	-	-	-
1,2,3,5-Tetrachlorobenzene			0.37	0.70	380	23	0.96	0.37	0.70	230	-	-	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene			0.19	0.37	34	12	0.49	0.19	0.37	66	-	-	-	-	-	-	-	-
Pentachlorobenzene			24	5.2	160	1000	70	24	47	320	-	-	-	NGR	5.2	-	-	-

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Hexachlorobenzene			1.8	2.3	5.0	34	2.3	1.8	3.6	47	-	-	-	-	-	-	-	-
2,4-Dichlorophenol			0.0029	0.0034	92,000	NGR	57,000	0.018	0.034	57,000	-	-	-	0.0029	0.0034	-	-	-
2,4,6-Trichlorophenol			0.19	0.37	400	71,000	3,300	0.19	0.37	3,700	-	-	-	0.42	0.50	-	-	-
2,3,4,6-Tetrachlorophenol			0.039	0.047	9,200	110,000	4,400	0.16	0.31	5,700	-	-	-	0.039	0.047	-	-	-
Pentachlorophenol			0.025	0.029	960	NGR	950,000	6.0	12	1,900	28	28	-	0.025	0.029	160	-	-
Dioxins & Furans <sup>h,i</sup>			0.0000040	0.0000040	-	-	-	-	-	0.0000040	-	-	-	-	-	-	-	-
PCBs			1.2	1.2	4.4	-	-	-	-	1.2	33	33	-	-	-	470	-	-
Pesticides																		
Aldicarb <sup>g</sup>			0.041	0.065	160	-	-	0.041	0.065	320	-	-	-	BDL	BDL	-	-	-
Aldrin			0.24	0.46	0.47	330	13	0.24	0.46	4.4	-	-	-	-	-	-	-	-
Atrazine and metabolites			0.0088	0.010	80	-	-	0.10	0.19	160	-	-	-	0.0088	0.010	-	-	-
Azniphos-methyl (Guthion)			0.41	0.75	400	-	-	0.41	0.75	790	-	-	-	-	-	-	-	-
Bromacil <sup>k</sup>			0.009	0.009	15,000	-	-	7.0	10	30,000	0.49	0.20	-	0.009	0.009	1.7	-	-
Bromoxynil			0.044	0.052	80	-	-	0.18	0.35	160	-	-	-	0.044	0.052	-	-	-
Carbaryl <sup>g</sup>			1.9	3.6	1,600	-	-	1.9	3.6	3,200	-	-	-	BDL	BDL	-	-	-
Carbofuran <sup>g</sup>			0.68	1.2	1,600	-	-	0.68	1.2	3,200	-	-	-	BDL	BDL	-	-	-
Chlorothalonil			0.0085	0.010	2,400	-	-	27	53	4,800	-	-	-	0.0085	0.010	-	-	-
Chlorpyrifos <sup>g</sup>			49	95	1,600	-	-	49	95	3,200	-	-	-	BDL	BDL	-	-	-
2,4-D <sup>g</sup>			0.43	0.67	9,200	-	-	0.43	0.67	5,700	-	-	-	BDL	BDL	-	-	-

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway					Direct Soil Contact	Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
DDT			12	12	24	660,000	25,000	89	170	160	12	12	550	-	-	170	-	-
Diazinon <sup>9</sup>			2.2	4.2	320	-	-	2.2	4.2	630	-	-	-	BDL	BDL	-	-	-
Dicamba <sup>9</sup>			0.50	0.79	2,000	-	-	0.50	0.79	4,000	-	-	-	BDL	BDL	-	-	-
Diclofop-methyl			570	2.4	920	-	-	NGR	NGR	570	-	-	-	NGR	2.4	-	-	-
Dieldrin			0.025	0.048	0.50	160	6.1	0.025	0.048	4.7	-	-	-	-	-	-	-	-
Dimethoate			0.0058	0.0055	1,800	-	-	0.077	0.12	1,100	-	-	-	0.0058	0.0055	-	-	-
Dinoseb <sup>9</sup>			2.8	5.5	160	-	-	2.8	5.5	320	-	-	-	BDL	BDL	-	-	-
Diquat			11	21	1,300	-	-	11	21	2,500	-	-	-	-	-	-	-	-
Diuron			1.9	3.5	2,500	-	-	1.9	3.5	4,900	-	-	-	-	-	-	-	-
Endosulfan			0.80	0.0016	5,500	-	-	99	190	3,400	-	-	-	0.80	0.0016	-	-	-
Endrin			2.4	4.7	48	-	-	2.4	4.7	96	-	-	-	-	-	-	-	-
Glyphosate			0.054	0.049	4,800	-	-	0.95	1.4	9,500	-	-	-	0.054	0.049	-	-	-
Heptachlor epoxide			0.039	0.076	0.88	2.4	0.094	0.039	0.076	4.1	-	-	-	-	-	-	-	-
Lindane <sup>9</sup>			0.31	0.60	48	-	-	0.31	0.60	95	-	-	-	BDL	BDL	-	-	-
Linuron			0.051	0.059	320	-	-	0.56	1.1	630	-	-	-	0.051	0.059	-	-	-
Malathion <sup>9</sup>			0.82	1.3	3,200	-	-	0.82	1.3	6,300	-	-	-	BDL	BDL	-	-	-
MCPA <sup>9</sup>			0.42	0.66	8,200	-	-	0.42	0.66	160	-	-	-	BDL	BDL	-	-	-
Methoxychlor			800	0.32	800	-	-	NGR	NGR	1,600	-	-	-	NGR	0.32	-	-	-
Metolachlor			0.048	0.055	800	-	-	1.3	2.4	1,600	-	-	-	0.048	0.055	-	-	-

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit		
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Metribuzin			0.024	0.028	1,300	-	-	7.8	15	2,600	-	-	-	0.024	0.028	-	-	-
Paraquat (as dichloride)			1.1	2.2	160	-	-	1.1	2.2	320	-	-	-	-	-	-	-	-
Phorate			0.075	0.14	32	-	-	0.075	0.14	63	-	-	-	-	-	-	-	-
Picloram			0.024	0.022	3,200	-	-	0.64	0.94	6,300	-	-	-	0.024	0.022	-	-	-
Simazine			0.033	0.038	210	-	-	0.14	0.25	410	-	-	-	0.033	0.038	-	-	-
Tebuthiuron <sup>9,i</sup>			0.60	0.60	11,000	-	-	2.5	3.7	22,000	0.60	0.60	-	BDL	BDL	0.66	-	-
Terbufos			0.080	0.15	8.0	-	-	0.080	0.15	16	-	-	-	-	-	-	-	-
Toxaphene			3.3	6.3	7.3	36,000	1,400	3.3	6.3	69	-	-	-	-	-	-	-	-
Triallate			0.0077	0.0092	12,000	-	-	16	31	7,400	-	-	-	0.0077	0.0092	-	-	-
Trifluralin			0.22	0.045	4,400	-	-	NGR	NGR	2,700	-	-	-	0.22	0.045	-	-	-
Other Organics																		
Aniline <sup>9</sup>			0.36	0.60	6,400	140	9.8	0.36	0.60	4,000	-	-	-	BDL	BDL	-	-	-
Di- <i>n</i> -butyl phthalate			0.54	0.65	9,600	NGR	82,000	70	130	19,000	-	-	-	0.54	0.65	-	-	-
Dichlorobenzidine			0.60	1.2	18	NGR	NGR	0.60	1.2	170	-	-	-	-	-	-	-	-
Diethanolamine <sup>j</sup>			2.0	3.5	1,000	-	-	2.0	3.5	2,000	2,000	2,000	-	500,000	45	15,000	-	-
Diethylene glycol			10	15	100,000	-	-	10	15	200,000	1,500	1,500	-	2,000	65	15,000	-	-
Diisopropanolamine			14	17	590,000	-	-	130	250	370,000	750	750	-	14	17	5,100	-	-
Ethylene glycol			60	62	NGR	NGR	NGR	60	68	NGR	1,800	1,800	2,000	89	62	16,000	-	-
Hexachlorobutadiene			0.026	0.031	590	1.2	0.078	0.50	0.95	5,600	-	-	-	0.026	0.031	-	-	-

**TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS**  
**This table must not be used for Tier 1 assessment and remediation.** Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline			Human					Ecological					Other		
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration		Direct Soil Contact		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type			Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse		Fine	Coarse	-	Fine	Coarse
Building Type					-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-
Unit (unless otherwise indicated)			(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Methanol			37	11	370,000	210,000	18,000	37	42	-	1,600	1,600	-	300	11	-	750	750
Methylmethacrylate			36	17	NGR	280	17	36	49	800,000	-	-	-	-	-	-	-	-
Monoethanolamine <sup>j</sup>			20	10	10,000	-	-	20	40	20,000	1,500	1,500	-	300,000	10	20,000	-	-
MTBE			0.044	0.062	9,200	7.4	0.57	0.044	0.062	5,700	-	-	-	7.1	6.1	-	-	-
Nonylphenol + ethoxylates			14	14	-	-	-	-	-	-	14	14	-	NGR	2,000	82	-	-
Perfluorooctane sulfonate (PFOS)			0.070	0.14	39	-	-	0.070	0.14	30	60	60	-	0.19	0.23	140	-	-
Phenol			0.0028	0.0024	55,000	90,000	5,800	1.6	2.3	34,000	130	130	-	0.0028	0.0024	290	-	-
Sulfolane			0.18	0.21	2,600	-	-	0.18	0.21	5,000	430	430	-	24	18	3,000	-	-
Triethylene glycol			100	150	NGR	-	-	100	150	NGR	7,000	7,000	-	10,000	200	70,000	-	-

Notes:

a. For more information see Guidelines for Landfill Disposal of Sulphur Waste and Remediation of Sulphur Containing Soils (AENV, 2011)

b. True total barium as measured by fusion-XRF or fusion-ICP. For more information see Soil Remediation Guidelines for Barite: Environmental Health and Human Health (AENV, 2009)

c. Human health direct soil contact guidelines for carcinogenic PAHs are based on B[a]P Total Potency Equivalents (TPE). TPEs are calculated by multiplying the soil concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (WHO/IPCS, 1998) scheme, as follows:

TABLE A-5. SURFACE SOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS

This table must not be used for Tier 1 assessment and remediation. Tier 1 soil guidelines are found in Tables 1, 3 and 4.

This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

- d. The B[a]P Total Potency Equivalents (TPEs) calculated for specific soil samples using Potency Equivalency Factors (PEFs) should be multiplied by an Uncertainty Factor of 3 when evaluating PAH mixtures associated with creosote or coal tar-type environmental releases, prior to evaluating against the human health direct contact soil remediation guideline.
- e. The Index of Additive Cancer Risk (IACR) is calculated by dividing the soil concentration of each carcinogenic PAH by its Protection of Domestic Use Aquifer guideline value to calculate a hazard index for each PAH and subsequently summing the hazard indexes for the entire PAH mixture. For example, the IACR for coarse textured soil would be calculated as follows:

Fine Soil:

$$IACR = \frac{[Benz(a)anthracene]}{6.4} + \frac{[Benzo(b + j)fluoranthene]}{3.0} + \frac{[Benzo(k)fluoranthene]}{0.64} + \frac{[Benzo(g, h, i)perylene]}{130} + \frac{[Benzo(a)pyrene]}{7.0} + \frac{[Chrysene]}{40} + \frac{[Dibenz(a, h)anthracene]}{4.4} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{51}$$

Coarse Soil:

$$IACR = \frac{[Benz(a)anthracene]}{12} + \frac{[Benzo(b + j)fluoranthene]}{5.8} + \frac{[Benzo(k)fluoranthene]}{1.2} + \frac{[Benzo(g, h, i)perylene]}{250} + \frac{[Benzo(a)pyrene]}{14} + \frac{[Chrysene]}{78} + \frac{[Dibenz(a, h)anthracene]}{8.5} + \frac{[Indeno(1,2,3 - c, d)pyrene]}{98}$$

- f. Overall guideline value for ecological receptors only.
- g. Guideline for protection of aquatic life is below detection limit, groundwater monitoring is required.
- h. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- i. Guideline values adopted directly from CCME (1999 and updates) without change.
- j. Analytical methodology specified in the Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine (AENV, 2010), or equivalent, must be used. See AENV (2010) for further details.
- k. Eco-contact guidelines from Stantec (2012).
- l. Eco-contact guidelines from Stantec (2008).
- m. Boron must be measured in a saturated paste extract prepared in accordance with Method 15.2.1 (Carter and Gregorich, 2008).
- n. Guideline for protection of aquatic life (fine soil) is based on a groundwater guideline of 0.10 g/L. See Appendix B for more information

BDL - Below detection limit

NGR - no guideline required, calculated value >1,000,000 mg/kg; or for PAH groundwater protection, calculated value results in groundwater concentration greater than solubility.

**TABLE A-6. SUBSOIL REMEDIATION GUIDELINE VALUES FOR NATURAL AREA LAND USE - ALL EXPOSURE PATHWAYS (BTEX AND PHC ONLY)**  
**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human		Ecological								Other	
Pathway			Protection of Domestic Use Aquifer		Direct Soil Contact <sup>a</sup>		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Wildlife Water		Management Limit		
Soil Type	Fine	Coarse	Fine	Coarse	Fine	Coarse		-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	
Building Type			-	-	-	-	-	-	-	-	-	-	-	-	-	
Unit	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	
Benzene	0.046	0.078	0.046	0.078	120	62	-	na	na	7.9	0.17	15	0.33	-	-	
Toluene	0.52	0.12	0.52	0.95	220	150	-	na	na	63,000	0.12	NGR	1,000	-	-	
Ethylbenzene	0.073	0.14	0.073	0.14	240	110	-	na	na	NGR	540	NGR	17,000	-	-	
Xylenes	0.99	1.9	0.99	1.9	130	190	-	na	na	NGR	41	NGR	16,000	-	-	
F1	420	420	1,100	2,200	420	420	-	na	na	30,000	1,300	NGR	30,000	800	700	
F2	300	300	1,500	2,900	300	300	-	na	na	30,000	520	NGR	30,000	1,000	1,000	
F3	2,600	600	-	-	2,600	600	-	na	na	-	-	-	-	3,500	2,500	
F4	10,000	5,600	-	-	11,200	5,600	-	na	na	-	-	-	-	10,000	10,000	

Notes:  
 a. Exclusion of the ecological direct soil contact pathway for F1, F2, F3, and F4 is permitted below 3 metres  
 na = exposure pathway not applicable to subsoil  
 NGR - no guideline required, calculated value >1,000,000 mg/kg

**TABLE A-7. SUBSOIL REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND USE - ALL EXPOSURE PATHWAYS (BTEX AND PHC ONLY)**  
**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological												Other		
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contact <sup>a</sup>		Nutrient/ Energy Cycling Check	Livestock Soil and Food Ingestion	Wildlife Soil and Food Ingestion	Protection of Freshwater Aquatic Life		Protection of Livestock Water		Protection of Wildlife Water		Protection of Irrigation Water		Management Limit		
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse			-	-	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Unit	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Benzene	0.046	0.021	360	0.36	0.34	0.021	0.028	0.046	0.078	120	62	-	na	na	7.9	0.17	0.20	0.21	15	0.33	-	-	-	-	
Toluene	0.52	0.12	690	1,300	1,200	80	110	0.52	0.95	220	150	-	na	na	63,000	0.12	26	29	NGR	1,000	-	-	-	-	
Ethylbenzene	0.073	0.14	1,900	2,100	1,900	120	170	0.073	0.14	240	110	-	na	na	NGR	540	36	42	NGR	17,000	-	-	-	-	
Xylenes	0.99	1.9	510	150	140	9.0	13	0.99	1.9	130	190	-	na	na	NGR	41	160	180	NGR	16,000	-	-	-	-	
F1	420	30	14,000	710	630	30	55	1,100	2,200	420	420	-	na	na	30,000	1,300	6,600	7,300	NGR	30,000	-	-	800	700	
F2	300	160	6,800	3,600	3,300	160	290	1,500	2,900	300	300	-	na	na	30,000	520	16,000	19,000	NGR	30,000	-	-	1,000	1,000	
F3	2,600	600	15,000	-	-	-	-	-	-	2,600	600	-	na	na	-	-	-	-	-	-	-	-	3,500	2,500	
F4	10,000	5,600	21,000	-	-	-	-	-	-	11,200	5,600	-	na	na	-	-	-	-	-	-	-	-	10,000	10,000	

**Notes:**  
 a. Exclusion of the ecological direct soil contact pathway for F1, F2, F3, and F4 is permitted below 3 metres  
 na = exposure pathway not applicable to subsoil  
 NGR - no guideline required, calculated value >1,000,000 mg/kg



TABLE A-8. SUBSOIL REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND LAND USE - ALL EXPOSURE PATHWAYS (BTEX AND PHC ONLY)

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological				Other	
Pathway			Direct Soil Contact		Vapour Inhalation			Protection of Domestic Use Aquifer		Direct Soil Contacta		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Fine	Coarse	Coarse	Fine	Coarse	Fine	Coarse		Fine	Coarse	Fine	Coarse
Building Type			-	Basement	Slab	Basement	Slab	-	-	-	-	-	-	-	-	-
Unit	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Benzene	0.046	0.021	360	0.36	0.34	0.021	0.028	0.046	0.078	120	62	-	7.9	0.17	-	-
Toluene	0.52	0.12	690	1,300	1,200	80	110	0.52	0.95	220	150	-	63,000	0.12	-	-
Ethylbenzene	0.073	0.14	1,900	2,100	1,900	120	170	0.073	0.14	240	110	-	NGR	540	-	-
Xylenes	0.99	1.9	510	150	140	9.0	13	0.99	1.9	130	190	-	NGR	41	-	-
F1	420	30	14,000	710	630	30	55	1,100	2,200	420	420	-	30,000	1300	800	700
F2	300	160	6,800	3,600	3,300	160	290	1,500	2,900	300	300	-	30,000	520	1,000	1,000
F3	2,600	600	15,000	-	-	-	-	-	-	2,600	600	-	-	-	3,500	2,500
F4	10,000	5,600	21,000	-	-	-	-	-	-	11,200	5,600	-	-	-	10,000	10,000

**Notes:**  
a. Exclusion of the ecological direct soil contact pathway for F1, F2, F3, and F4 is permitted below 3 metres  
na = exposure pathway not applicable to subsoil  
NGR - no guideline required, calculated value >1,000,000 mg/kg

**TABLE A-9. SUBSOIL REMEDIATION GUIDELINE VALUES FOR COMMERCIAL LAND USE - ALL EXPOSURE PATHWAYS (BTEx AND PhC ONLY)**  
**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 200 as amended 7).

Receptor			Overall Guideline			Human					Ecological					Other	
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration	Direct Soil Contacta		Nutrient/ Energy Cycling Check	Protection of Freshwater Aquatic Life		Off-Site Migration	Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse			Fine	Coarse	-	Fine	Coarse
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-	-
Unit	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)
Benzene	0.046	0.078	550	2.3	0.26	0.046	0.078	1,100	620	360	-	7.9	0.17	890	-	-	
Toluene	0.52	0.12	1,100	8,100	970	0.52	0.95	9,200	660	500	-	63,000	0.12	2,100	-	-	
Ethylbenzene	0.073	0.14	2,900	13,000	1,500	0.073	0.14	24,000	860	600	-	NGR	540	1,600	-	-	
Xylenes	0.99	1.9	780	970	110	0.99	1.9	6,900	460	700	-	NGR	41	930	-	-	
F1	640	440	22,000	4,700	440	1,100	2,200	30,000	640	640	-	30,000	1300	3,000	800	700	
F2	520	520	10,000	24,000	2,400	1,500	2,900	30,000	520	520	-	30,000	520	2,100	1,000	1,000	
F3	4,300	3,400	23,000	-	-	-	-	30,000	5,000	3,400	-	-	-	4,300	5,000	3,500	
F4	10,000	6,600	30,000	-	-	-	-	30,000	13,200	6,600	-	-	-	30,000	10,000	10,000	

**Notes:**  
 a. Exclusion of the ecological direct soil contact pathway for F1, F2, F3, and F4 is permitted below 3 metres  
 na = exposure pathway not applicable to subsoil  
 NGR - no guideline required, calculated value >1,000,000 mg/kg

**TABLE A-10. SUBSOIL REMEDIATION GUIDELINE VALUES FOR INDUSTRIAL LAND USE - ALL EXPOSURE PATHWAYS (BTX AND PHC ONLY)**  
**This table must not be used for Tier 1 assessment and remediation.** Tier 1 soil guidelines are found in Tables 1, 3 and 4.  
 This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Receptor			Overall Guideline		Human						Ecological						Other			
Pathway			Direct Soil Contact		Vapour Inhalation		Protection of Domestic Use Aquifer		Off-Site Migration		Direct Soil Contacta		Nutrient/ Energy Cycling of Freshwater Aquatic Life		Protection of Freshwater Aquatic Life		Off-Site Migration		Management Limit	
Soil Type	Fine	Coarse	-	Fine	Coarse	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse	-	Fine	Coarse				
Building Type			-	Slab	Slab	-	-	-	-	-	-	-	-	-	-	-				
Unit	(mg/kg)	(mg/kg)	(mg/kg)	mg/kg	mg/kg	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)	(mg/kg)				
Benzene	0.046	0.078	550	2.3	0.26	0.046	0.078	1,100	620	360	-	7.9	0.17	890	-	-				
Toluene	0.52	0.12	16,000	8,100	970	0.52	0.95	9,200	660	500	-	63,000	0.12	2,100	-	-				
Ethylbenzene	0.073	0.14	44,000	13,000	1,500	0.073	0.14	24,000	860	600	-	NGR	540	1,600	-	-				
Xylenes	0.99	1.9	12,000	970	110	0.99	1.9	6,900	460	700	-	NGR	41	930	-	-				
F1	640	440	30,000	4,700	440	1,100	2,200	30,000	640	640	-	30,000	1300	3,000	800	700				
F2	520	520	30,000	24,000	2,400	1,500	2,900	30,000	520	520	-	30,000	520	2,100	1,000	1,000				
F3	4,300	3,400	30,000	-	-	-	-	30,000	5,000	3,400	-	-	-	4,300	5,000	3,500				
F4	10,000	6,600	30,000	-	-	-	-	30,000	13,200	6,600	-	-	-	30,000	10,000	10,000				

**Notes:**  
 a. Exclusion of the ecological direct soil contact pathway for F1, F2, F3, and F4 is permitted below 3 metres  
 na = exposure pathway not applicable to subsoil  
 NGR - no guideline required, calculated value >1,000,000 mg/kg

**Appendix B**  
**GROUNDWATER Remediation Guidelines**  
**All Water Uses**

Appendix B consists of four tables, one for each of the land uses, except that commercial and industrial groundwater guidelines are identical, and separate tables are not required. Each table provides the groundwater remediation guideline for each applicable water use, where available. Tier 1 guidelines must be chosen from Tables 1 to 4, not from Appendix A or B. The only exception occurs when a more sensitive land use borders on, or is less than 30 m from, the site of interest. Under this condition guidelines for specific exposure pathways on the more sensitive land use must be evaluated and if they are lower than the Tier 1 guideline, they must be applied to the site of interest (See Section 3.2 and 5.1.1 and Figure 1).

The four tables are as follows:

- Table B-1. Groundwater Remediation Guideline Values for Natural Area Land Use
- Table B-2. Groundwater Remediation Guideline Values for Agricultural Land Use
- Table B-3. Groundwater Remediation Guideline Values for Residential/Parkland Use
- Table B-4. Groundwater Remediation Guideline Values for Commercial/Industrial Land Use

Note that the groundwater guidelines protective of inhalation under agricultural or residential/parkland use are based on a building with slab-on-grade construction. These values are protective in all cases of both slab-on-grade and basement construction.

All water uses are applicable at Tier 1. However, it may be possible to exclude or modify certain water uses at Tier 2. The companion Tier 2 should be consulted for further information. The information in the tables in this appendix will assist in determining whether a Tier 2 approach for groundwater is likely to be useful at a given site.

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
<b>General and Inorganic Parameters</b>									
pH	6.5-8.5	6.5-8.5	6.5-8.5	-	-	6.5-9.0	6.5-9.0	-	-
Ammonia	see note d	see note d	-	-	-	see note d	see note d	-	-
Bromate	0.010	0.010	0.010	-	-	-	-	-	-
Chloride	120	120	250	-	-	120	120	-	-
Cyanide (free)	0.0050	0.0050	0.20	-	-	0.0050	0.0050	-	-
Fluoride	1.5	1.5	1.5	-	-	-	-	-	-
Nitrate (as nitrogen)	3.0	3.0	10	-	-	3.0	3.0	-	-
Nitrite (as nitrogen)	see note e	see note e	1.0	-	-	see note d	see note d	-	-
Sodium	200	200	200	-	-	-	-	-	-
Sulphate	see note e	see note e	500	-	-	see note d	see note d	-	-
Sulphide – Total (as S) <sup>i</sup>	0.0019	0.0019	0.050	-	-	0.0019	0.0019	-	-
Total Dissolved Solids (TDS)	500	500	500	-	-	-	-	-	-
<b>Metals</b>									
Aluminum	see note d	see note d	2.9	-	-	see note d	see note d	-	-
Antimony	0.0060	0.0060	0.0060	-	-	-	-	-	-
Arsenic	0.0050	0.0050	0.010	-	-	0.005	0.005	-	-
Barium	2.0	2.0	2.0	-	-	-	-	-	-
Boron	1.5	1.5	5.0	-	-	1.5	1.5	-	-
Cadmium	see note e	see note e	0.0070	-	-	see note d	see note d	-	-
Chromium (trivalent)	0.0089	0.0089	-	-	-	0.0089	0.0089	-	-
Chromium (hexavalent)	0.0010	0.0010	-	-	-	0.0010	0.0010	-	-
Chromium (total) <sup>j</sup>	0.050	0.050	0.050	-	-	-	-	-	-

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Copper	0.0070	0.0070	1.0	-	-	0.0070	0.0070	-	-
Iron	0.30	0.30	0.30	-	-	0.30	0.30	-	-
Lead	See note e	See note e	0.0050	-	-	see note d	see note d	-	-
Manganese	0.020	0.020	0.020	-	-	-	-	-	-
Mercury (total) <sup>j</sup>	0.0000050	0.0000050	0.0010	-	-	0.0000050	0.0000050	-	-
Nickel	see note d	see note d	-	-	-	see note d	see note d	-	-
Selenium	0.0020	0.0020	0.050	-	-	0.0020	0.0020	-	-
Silver	0.00010	0.00010	-	-	-	0.00010	0.00010	-	-
Uranium	0.015	0.015	0.020	-	-	0.015	0.015	-	-
Zinc	0.030	0.030	5.0	-	-	0.030	0.030	-	-
<b>Hydrocarbons</b>									
Benzene	0.0050	0.0050	0.0050	100	61	3.6	0.074	6.8	0.14
Toluene	0.024	0.021	0.024	82	59	12,000	0.021	NGR	180
Ethylbenzene	0.0016	0.0016	0.0016	42	20	NGR	41	NGR	NGR
Xylenes	0.020	0.020	0.020	21	31	NGR	2.9	NGR	NGR
Styrene	0.072	0.072	2.8	-	-	0.072	0.072	-	-
F1	2.2	2.2	2.2	6.5	7.1	NGR	9.8	NGR	NGR
F2	1.1	1.1	1.1	1.8	1.8	NGR	1.3	NGR	NGR
Acenaphthene	0.0060	0.0058	1.4	-	-	0.0060	0.0058	NGR	NGR
Anthracene	0.0034	0.000012	NGR	0.025	0.025	0.0034	0.000012	NGR	NGR
Fluoranthene	0.24	0.000057	NGR	0.24	0.24	NGR	0.000057	NGR	NGR
Fluorene	0.0042	0.0030	0.94	-	-	0.0042	0.0030	NGR	NGR
Naphthalene	0.0010	0.0010	0.47	-	-	0.0010	0.0010	NGR	NGR

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Phenanthrene	0.00086	0.00040	-	-	-	0.00086	0.00040	NGR	NGR
Pyrene	0.71	0.000092	0.71	-	-	NGR	0.000092	NGR	NGR
Carcinogenic PAHs (as B(a)P TPE) <sup>a</sup>	0.000040	0.000040	0.000040	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	NGR	NGR	NGR	NGR
Benzo[b+j]fluoranthene	-	-	-	-	-	-	-	NGR	NGR
Benzo[k]fluoranthene	-	-	-	-	-	-	-	NGR	NGR
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>b</sup>	0.0018	0.0018	-	0.0018	0.0018	NGR	NGR	NGR	NGR
Chrysene	-	-	-	-	-	-	-	NGR	NGR
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	NGR	NGR
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	-	-
<b>Halogenated Aliphatics</b>									
Vinyl chloride	0.0020	0.0020	0.0020	-	-	-	-	-	-
1,1-Dichloroethene	0.014	0.014	0.014	-	-	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.0050	0.0050	0.0050	4.4	5.0	0.27	0.029	-	-
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.010	0.010	-	-	0.11	0.11	-	-
1,2-Dichloroethane	0.0050	0.0050	0.0050	-	-	0.10	0.10	-	-
Dichloromethane (Methylene chloride)	0.050	0.050	0.050	-	-	0.098	0.098	-	-
Trichloromethane (Chloroform) <sup>i</sup>	0.080	0.018	0.080	-	-	0.10	0.018	-	-
Tetrachloromethane (Carbon tetrachloride)	0.0020	0.0020	0.0020	-	-	0.013	0.013	-	-
Dibromochloromethane	0.19	0.19	0.19	-	-	-	-	-	-
<b>Chlorinated Aromatics</b>									
Chlorobenzene	0.0013	0.0013	0.030	-	-	0.0013	0.0013	-	-

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
1,2-Dichlorobenzene	0.00070	0.00070	0.0030	-	-	0.00070	0.00070	-	-
1,4-Dichlorobenzene	0.0010	0.0010	0.0010	-	-	0.026	0.026	-	-
1,2,3-Trichlorobenzene	0.0080	0.0080	0.014	-	-	0.0080	0.0080	-	-
1,2,4-Trichlorobenzene	0.015	0.015	0.015	-	-	0.024	0.024	-	-
1,3,5-Trichlorobenzene	0.014	0.014	0.014	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.0018	0.0018	0.032	-	-	0.0018	0.0018	-	-
1,2,3,5-Tetrachlorobenzene	0.0038	0.0038	0.0038	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.0020	0.0020	0.002 0	-	-	-	-	-	-
Pentachlorobenzene	0.0094	0.0069	0.0094	-	-	NGR	0.0069	-	-
Hexachlorobenzene	0.00029	0.00029	0.00029	-	-	-	-	-	-
2,4-Dichlorophenol	0.00020	0.00020	0.00030	-	-	0.00020	0.00020	-	-
2,4,6-Trichlorophenol	0.0020	0.0020	0.0020	-	-	0.018	0.018	-	-
2,3,4,6-Tetrachlorophenol	0.0010	0.0010	0.0010	-	-	0.0010	0.0010	-	-
Pentachlorophenol	0.00051	0.00050	0.03	0.87	0.88	0.00051	0.00050	-	-
Dioxins & Furans <sup>c</sup>	0.00000012	0.00000012	0.00000012	-	-	-	-	-	-
PCBs	0.000094	0.000094	0.000094	-	-	-	-	-	-
<b>Pesticides</b>									
Aldicarb	0.0010	0.0010	0.0090	-	-	0.0010	0.0010	-	-
Aldrin	0.000028	0.000028	0.000028	-	-	-	-	-	-
Atrazine and metabolites	0.0018	0.0018	0.0050	-	-	0.0018	0.0018	-	-
Azniphos-methyl (Guthion)	0.000010	0.000010	0.020	-	-	0.000010	0.000010	-	-
Bromacil <sup>9</sup>	0.0050	0.0050	0.95	0.44	0.30	0.0050	0.0050	-	-
Bromoxynil	0.0050	0.0050	0.0050	-	-	0.0050	0.0050	-	-



**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Carbaryl	0.00020	0.00020	0.090	-	-	0.00020	0.00020	-	-
Carbofuran	0.0018	0.0018	0.090	-	-	0.0018	0.0018	-	-
Chlorothalonil	0.00018	0.00018	0.14	-	-	0.00018	0.00018	-	-
Chlorpyrifos	0.0000046	0.0000020	0.09	-	-	0.0000046	0.0000020	-	-
2,4-D	0.0040	0.0040	0.10	-	-	0.0040	0.0040	-	-
DDT	0.0014	0.0014	0.0014	-	-	-	-	-	-
Diazinon	0.00017	0.00017	0.02	-	-	0.00017	0.00017	-	-
Dicamba	0.010	0.010	0.12	-	-	0.010	0.010	-	-
Diclofop-methyl	0.0090	0.0061	0.0090	-	-	0.56	0.0061	-	-
Dieldrin	0.000029	0.000029	0.000029	-	-	-	-	-	-
Dimethoate	0.0062	0.0062	0.020	-	-	0.0062	0.0062	-	-
Dinoseb	0.000055	0.000050	0.010	-	-	0.000055	0.000050	-	-
Diquat	0.070	0.070	0.070	-	-	-	-	-	-
Diuron	0.15	0.15	0.15	-	-	-	-	-	-
Endosulfan	0.0019	0.0000031	0.057	-	-	0.0019	0.0000031	-	-
Endrin	0.0028	0.0028	0.0028	-	-	-	-	-	-
Glyphosate	0.065	0.065	0.28	-	-	0.065	0.065	-	-
Heptachlor epoxide	0.000052	0.000052	0.000052	-	-	-	-	-	-
Lindane	0.000010	0.000010	0.0028	-	-	0.000010	0.000010	-	-
Linuron	0.0070	0.0070	0.019	-	-	0.0070	0.0070	-	-
Malathion	0.00010	0.00010	0.19	-	-	0.00010	0.00010	-	-
MCPA	0.0026	0.0026	0.10	-	-	0.0026	0.0026	-	-
Methoxychlor	0.90	0.00017	0.90	-	-	NGR	0.00017	-	-
Metolachlor	0.0078	0.0078	0.050	-	-	0.0078	0.0078	-	-

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Metribuzin	0.0010	0.0010	0.080	-	-	0.0010	0.0010	-	-
Paraquat (as dichloride)	0.010	0.010	0.010	-	-	-	-	-	-
Phorate	0.0020	0.0020	0.0020	-	-	-	-	-	-
Picloram	0.029	0.029	0.19	-	-	0.029	0.029	-	-
Simazine	0.010	0.010	0.010	-	-	0.010	0.010	-	-
Tebuthiuron <sup>h</sup>	0.0016	0.0016	0.66	0.20	0.25	0.0016	0.0016	-	-
Terbufos	0.0010	0.0010	0.0010	-	-	-	-	-	-
Toxaphene	0.00043	0.00043	0.00043	-	-	-	-	-	-
Triallate	0.00024	0.00024	0.12	-	-	0.00024	0.00024	-	-
Trifluralin	0.0012	0.00020	0.045	-	-	0.0012	0.00020	-	-
<b>Other Organics</b>									
Aniline	0.0022	0.0022	0.066	-	-	0.0022	0.0022	-	-
Di- <i>n</i> -butyl phthalate	0.019	0.019	0.59	-	-	0.019	0.019	-	-
Dichlorobenzidine	0.0010	0.0010	0.0010	-	-	-	-	-	-
Diethanolamine	0.060	0.060	0.060	-	-	65,000	5.0	-	-
Diethylene glycol	6.0	6.0	6.0	-	-	4,000	200	-	-
Diisopropanolamine	1.6	1.6	3.6	160	160	1.6	1.6	-	-
Ethylene glycol	31	31	31	9,200	16,000	190	190	-	-
Hexachlorobutadiene	0.0013	0.0013	0.0060	-	-	0.0013	0.0013	-	-
Methanol	19	19	19	-	-	630	32	-	-
Methylmethacrylate	13	13	13	-	-	-	-	-	-
Monoethanolamine	0.60	0.60	0.60	-	-	30,000	1.0	-	-
MTBE	0.015	0.015	0.015	-	-	10	10	-	-

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable GW	Eco Soil Contact		Aquatic Life		Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Nitritotriacetic acid	0.40	0.40	0.40	-	-	-	-	-	-
Nonylphenol + ethoxylates	0.0081	0.0081	-	0.0081	0.0081	NGR	0.61	-	-
Perfluorooctane sulfonate (PFOS) <sup>k</sup>	0.00060	0.00060	0.00060	1.4	1.4	0.0068	0.0068	0.052	0.052
Perfluorooctanoic acid (PFOA) <sup>k</sup>	0.00020	0.00020	0.00020	-	-	-	-	-	-
Phenol	0.0040	0.0040	0.57	110	150	0.0040	0.0040	-	-
Sulfolane	0.090	0.090	0.090	1,700	2,800	50	50	-	-
Triethylene glycol	60	60	60	-	-	25,000	550	-	-
Trihalomethanes - total (THMs)	0.10	0.10	0.10	-	-	-	-	-	-

Notes:

- a. B[a]P TPE (Total Potency Equivalents) are calculated by multiplying the groundwater concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (1999) scheme, as follows:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

---

**TABLE B-1. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR NATURAL AREAS - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

- b. For ecological receptors only.
- c. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- d. See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014 and updates) for further guidance on aquatic life pathway.
- e. Tier 1 guideline = lowest of aquatic life guideline and potable GW guideline.
- f. As S, but can be applied to undissociated H<sub>2</sub>S if concerns arise.
- g. Eco-contact guidelines from Stantec (2012)
- h. Eco-contact guidelines from Stantec (2008)
- i. Guideline for protection of aquatic life (fine soil) is set at the maximum concentration of trichloromethane that will support biological degradation (MEMS, 2016).
- j. Total means all chemical species.
- k. As the toxicological effects of PFOA and perfluorooctanoyl sulfonate (PFOS) are considered to be additive, the sum of the ratios of the detected concentrations to the corresponding MACs for PFOS and PFOA should not exceed 1.

NGR - no guideline required, calculated value > solubility or >1,000,000 mg/L

Potable GW = protection of groundwater for potable drinking water

Eco Soil Contact = protection of terrestrial plants and soil invertebrates in areas with shallow groundwater

Aquatic Life = protection of groundwater discharging to a surface water body hosting aquatic life

Wildlife Watering = protection of groundwater discharging to a surface water body from which wildlife may drink

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
<b>General and Inorganic Parameters</b>													
pH	6.5-8.5	6.5-8.5	6.5-8.5	-	-	-	-	6.5-9.0	6.5-9.0	-	-	-	-
Ammonia	see note d	see note d	-	-	-	-	-	see note d	see note d	-	-	-	-
Bromate	0.010	0.010	0.010	-	-	-	-	-	-	-	-	-	-
Chloride	100	100	250	-	-	-	-	120	120	100	-	-	-
Cyanide (free)	0.0050	0.0050	0.20	-	-	-	-	0.0050	0.0050	-	-	-	-
Electrical Conductivity (dS/m)	1.0	1.0								1.0			
Fluoride	1.0	1.0	1.5	-	-	-	-	-	-	1.0	1.0	-	-
Nitrate (as nitrogen)	3.0	3.0	10	-	-	-	-	3.0	3.0	-	-	-	-
Nitrate + Nitrite (as nitrogen)	100	100	-	-	-	-	-	-	-	-	100	-	-
Nitrite (as nitrogen)	see note e	see note e	1.0	-	-	-	-	see note d	see note d	-	10	-	-
Sodium	200	200	200	-	-	-	-	-	-	-	-	-	-
Sodium Adsorption Ratio	5.0	5.0								5.0			
Sulphate	see note e	see note e	500	-	-	-	-	see note d	see note d	-	1000	-	-
Sulphide – Total (as S) <sup>f</sup>	0.0019	0.0019	0.050	-	-	-	-	0.0019	0.0019	-	-	-	-
Total Dissolved Solids (TDS)	500	500	500	-	-	-	-	-	-	-	3000	-	-
<b>Metals</b>													
Aluminum	see note d	see note d	2.9	-	-	-	-	see note d	see note d	5.0	5.0	-	-
Antimony	0.0060	0.0060	0.0060	-	-	-	-	-	-	-	-	-	-
Arsenic	0.0050	0.0050	0.010	-	-	-	-	0.0050	0.0050	0.16	0.025	-	-
Barium	2.0	2.0	2.0	-	-	-	-	-	-	-	-	-	-
Boron	1.0	1.0	5.0	-	-	-	-	1.5	1.5	1.0	5.0	-	-
Cadmium	see note e	see note e	0.0070	-	-	-	-	see note d	see note d	0.0082	0.080	-	-

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	Irrigation		Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Chromium (trivalent)	0.0049	0.0049	-	-	-	-	-	0.0089	0.0089	0.0049	0.050	-	-
Chromium (hexavalent)	0.0010	0.0010	-	-	-	-	-	0.0010	0.0010	0.0080	0.050	-	-
Chromium (total) <sup>j</sup>	0.050	0.050	0.050	-	-	-	-	-	-	-	-	-	-
Copper	0.0070	0.0070	1.0	-	-	-	-	0.007	0.007	0.20	0.50	-	-
Iron	0.30	0.30	0.30	-	-	-	-	0.30	0.30	5.0	-	-	-
Lead	see note e	see note e	0.0050	-	-	-	-	see note d	see note d	0.20	0.10	-	-
Manganese	0.020	0.020	0.020	-	-	-	-	-	-	0.20	-	-	-
Mercury (total) <sup>j</sup>	0.0000050	0.0000050	0.001	-	-	-	-	0.0000050	0.0000050	-	0.003	-	-
Nickel	see note e	see note e	-	-	-	-	-	see note d	see note d	0.20	1.0	-	-
Selenium	0.0020	0.0020	0.050	-	-	-	-	0.0020	0.0020	0.020	0.050	-	-
Silver	0.00010	0.00010	-	-	-	-	-	0.00010	0.00010	0.020	0.050	-	-
Uranium	0.010	0.010	0.020	-	-	-	-	0.015	0.015	0.010	0.20	-	-
Zinc	0.030	0.030	5.0	-	-	-	-	0.030	0.030	1.0	50	-	-
<b>Hydrocarbons</b>													
Benzene	0.0050	0.0050	0.0050	0.57	0.03	100	61	3.6	0.074	-	0.088	6.8	0.14
Toluene	0.024	0.021	0.024	NGR	45	82	59	12,000	0.021	-	4.9	NGR	180
Ethylbenzene	0.0016	0.0016	0.0016	NGR	31	42	20	NGR	41	-	3.2	NGR	NGR
Xylenes	0.020	0.020	0.020	44	2.2	21	31	NGR	2.9	-	13	NGR	NGR
Styrene	0.072	0.072	2.8	NGR	39	-	-	0.072	0.072	-	-	-	-
F1	2.2	0.81	2.2	19	0.81	6.5	7.1	NGR	9.8	-	53	NGR	NGR
F2	1.1	1.1	1.1	NGR	1.5	1.8	1.8	NGR	1.3	-	NGR	NGR	NGR
Acenaphthene	0.0060	0.0058	1.4	NGR	NGR	-	-	0.0060	0.0058	-	NGR	NGR	NGR
Anthracene	0.0034	0.000012	NGR	NGR	NGR	0.025	0.025	0.0034	0.000012	-	NGR	NGR	NGR
Fluoranthene	0.24	0.000057	NGR	NGR	NGR	0.24	0.24	NGR	0.000057	-	NGR	NGR	NGR
Fluorene	0.0042	0.0030	0.94	NGR	NGR	-	-	0.0042	0.0030	-	NGR	NGR	NGR

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Naphthalene	0.0010	0.0010	0.47	NGR	2.7	-	-	0.0010	0.0010	-	NGR	NGR	NGR
Phenanthrene	0.00086	0.00040	-	-	-	-	-	0.00086	0.00040	-	NGR	NGR	NGR
Pyrene	0.71	0.000092	0.71	NGR	NGR	-	-	NGR	0.000092	-	NGR	NGR	NGR
Carcinogenic PAHs (as B(a)P TPE) <sup>a</sup>	0.000040	0.000040	0.000040	-	-	-	-	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	-	-	NGR	NGR	-	NGR	NGR	NGR
Benzo[b+j]fluoranthene	-	-	-	-	-	-	-	-	-	-	NGR	NGR	NGR
Benzo[k]fluoranthene	-	-	-	-	-	-	-	-	-	-	NGR	NGR	NGR
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>b</sup>	0.0018	0.0018	-	-	-	0.0018	0.0018	NGR	NGR	-	NGR	NGR	NGR
Chrysene	-	-	-	-	-	-	-	-	-	-	NGR	NGR	NGR
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	-	-	-	NGR	NGR	NGR
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Halogenated Aliphatics</b>													
Vinyl chloride	0.0020	0.0011	0.0020	0.018	0.0011	-	-	-	-	-	-	-	-
1,1-Dichloroethene	0.014	0.014	0.014	0.68	0.039	-	-	-	-	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.0050	0.00032	0.0050	0.0064	0.00032	4.4	5.0	0.27	0.029	-	0.05	-	-
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.010	0.010	0.25	0.012	-	-	0.11	0.11	-	-	-	-
1,2-Dichloroethane	0.0050	0.0050	0.0050	0.17	0.010	-	-	0.10	0.10	-	0.0050	-	-
Dichloromethane (Methylene chloride)	0.050	0.050	0.050	17	0.94	-	-	0.098	0.098	-	0.050	-	-
Trichloromethane (Chloroform) <sup>i</sup>	0.080	0.018	0.080	7.1	0.41	-	-	0.10	0.018	-	0.10	-	-
Tetrachloromethane (Carbon tetrachloride)	0.0020	0.0015	0.0020	0.030	0.0015	-	-	0.013	0.013	-	0.0050	-	-

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Dibromochloromethane	0.10	0.10	0.19	26	1.1	-	-	-	-	-	0.10	-	-
<b>Chlorinated Aromatics</b>													
Chlorobenzene	0.0013	0.0013	0.030	0.30	0.014	-	-	0.0013	0.0013	-	-	-	-
1,2-Dichlorobenzene	0.00070	0.00070	0.0030	120	5.4	-	-	0.00070	0.00070	-	-	-	-
1,4-Dichlorobenzene	0.0010	0.0010	0.0010	2.8	0.13	-	-	0.026	0.026	-	-	-	-
1,2,3-Trichlorobenzene	0.0080	0.0080	0.014	0.80	0.032	-	-	0.0080	0.0080	-	-	-	-
1,2,4-Trichlorobenzene	0.015	0.015	0.015	0.71	0.028	-	-	0.024	0.024	-	-	-	-
1,3,5-Trichlorobenzene	0.014	0.014	0.014	0.38	0.015	-	-	-	-	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.0018	0.0018	0.032	NGR	0.14	-	-	0.0018	0.0018	-	-	-	-
1,2,3,5-Tetrachlorobenzene	0.0038	0.0038	0.0038	0.41	0.017	-	-	-	-	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.0020	0.0020	0.002 0	0.21	0.0088	-	-	-	-	-	-	-	-
Pentachlorobenzene	0.0094	0.0069	0.0094	NGR	0.038	-	-	NGR	0.0069	-	-	-	-
Hexachlorobenzene	0.00029	0.00029	0.00029	0.012	0.00050	-	-	-	-	-	0.00052	-	-
2,4-Dichlorophenol	0.00020	0.00020	0.00030	NGR	1500	-	-	0.00020	0.00020	-	-	-	-
2,4,6-Trichlorophenol	0.0020	0.0020	0.0020	NGR	54	-	-	0.018	0.018	-	-	-	-
2,3,4,6-Tetrachlorophenol	0.0010	0.0010	0.0010	NGR	NGR	-	-	0.0010	0.0010	-	-	-	-
Pentachlorophenol	0.00051	0.00050	0.030	NGR	NGR	0.87	0.88	0.00051	0.00050	-	-	-	-
Dioxins & Furans <sup>c</sup>	0.00000012	0.00000012	0.00000012	-	-	-	-	-	-	-	-	-	-
PCBs	0.000094	0.000094	0.000094	-	-	-	-	-	-	-	-	-	-
<b>Pesticides</b>													
Aldicarb	0.0010	0.0010	0.0090	-	-	-	-	0.0010	0.0010	0.073	0.011	-	-
Aldrin	0.000028	0.000028	0.000028	0.056	0.0026	-	-	-	-	-	-	-	-



**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Atrazine and metabolites	0.0018	0.0018	0.0050	-	-	-	-	0.0018	0.0018	0.010	0.005	-	-
Azniphos-methyl (Guthion)	0.000010	0.000010	0.020	-	-	-	-	0.000010	0.000010	-	-	-	-
Bromacil <sup>9</sup>	0.00020	0.00020	0.95	-	-	0.44	0.30	0.0050	0.0050	0.00020	1.1	-	-
Bromoxynil	0.00044	0.00044	0.0050	-	-	-	-	0.0050	0.0050	0.00044	0.011	-	-
Carbaryl	0.00020	0.00020	0.090	-	-	-	-	0.00020	0.00020	-	1.1	-	-
Carbofuran	0.0018	0.0018	0.090	-	-	-	-	0.0018	0.0018	-	0.045	-	-
Chlorothalonil	0.00018	0.00018	0.14	-	-	-	-	0.00018	0.00018	0.0093	0.17	-	-
Chlorpyrifos	0.0000046	0.0000020	0.090	-	-	-	-	0.0000046	0.0000020	-	0.024	-	-
2,4-D	0.0040	0.0040	0.10	-	-	-	-	0.0040	0.0040	-	0.10	-	-
DDT	0.0014	0.0014	0.0014	15	0.69	-	-	-	-	-	0.10	-	-
Diazinon	0.00017	0.00017	0.020	-	-	-	-	0.00017	0.00017	-	-	-	-
Dicamba	0.0000080	0.0000080	0.12	-	-	-	-	0.010	0.010	0.0000080	0.12	-	-
Diclofop-methyl	0.00024	0.00024	0.0090	-	-	-	-	0.56	0.0061	0.00024	0.0090	-	-
Dieldrin	0.000029	0.000029	0.000029	0.27	0.013	-	-	-	-	-	-	-	-
Dimethoate	0.0030	0.0030	0.020	-	-	-	-	0.0062	0.0062	-	0.0030	-	-
Dinoseb	0.000055	0.000050	0.010	-	-	-	-	0.000055	0.000050	0.021	0.15	-	-
Diquat	0.070	0.070	0.070	-	-	-	-	-	-	-	-	-	-
Diuron	0.15	0.15	0.15	-	-	-	-	-	-	-	-	-	-
Endosulfan	0.0019	0.0000031	0.057	-	-	-	-	0.0019	0.0000031	-	-	-	-
Endrin	0.0028	0.0028	0.0028	-	-	-	-	-	-	-	-	-	-
Glyphosate	0.065	0.065	0.28	-	-	-	-	0.065	0.065	-	0.28	-	-
Heptachlor epoxide	0.000052	0.000052	0.000052	0.0043	0.00021	-	-	-	-	-	-	-	-
Lindane	0.000010	0.000010	0.0028	-	-	-	-	0.000010	0.000010	-	0.0040	-	-
Linuron	0.00011	0.00011	0.019	-	-	-	-	0.0070	0.0070	0.00011	-	-	-
Malathion	0.00010	0.00010	0.19	-	-	-	-	0.00010	0.00010	-	-	-	-

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
MCPA	0.000040	0.000040	0.10	-	-	-	-	0.0026	0.0026	0.00004	0.025	-	-
Methoxychlor	0.90	0.00017	0.90	-	-	-	-	NGR	0.00017	-	-	-	-
Metolachlor	0.0078	0.0078	0.050	-	-	-	-	0.0078	0.0078	0.028	0.050	-	-
Metribuzin	0.00050	0.00050	0.080	-	-	-	-	0.0010	0.0010	0.00050	0.080	-	-
Paraquat (as dichloride)	0.010	0.010	0.010	-	-	-	-	-	-	-	-	-	-
Parathion	-	-	-	-	-	-	-	0.000013	0.000013	-	-	-	-
Phorate	0.0020	0.0020	0.0020	-	-	-	-	-	-	-	-	-	-
Picloram	0.029	0.029	0.19	-	-	-	-	0.029	0.029	-	0.19	-	-
Simazine	0.00050	0.00050	0.010	-	-	-	-	0.010	0.010	0.00050	0.010	-	-
Tebuthiuron <sup>h</sup>	0.00043	0.00043	0.66	-	-	0.20	0.25	0.0016	0.0016	0.00043	0.13	-	-
Terbufos	0.0010	0.0010	0.0010	-	-	-	-	-	-	-	-	-	-
Toxaphene	0.00043	0.00043	0.00043	6.4	0.31	-	-	-	-	-	-	-	-
Triallate	0.00024	0.00024	0.12	-	-	-	-	0.00024	0.00024	-	0.23	-	-
Trifluralin	0.0012	0.0002	0.045	-	-	-	-	0.0012	0.0002	-	0.045	-	-
<b>Other Organics</b>													
Aniline	0.0022	0.0022	0.066	59	2.8	-	-	0.0022	0.0022	-	-	-	-
Di- <i>n</i> -butyl phthalate	0.019	0.019	0.59	NGR	NGR	-	-	0.019	0.019	-	-	-	-
Dichlorobenzidine	0.0010	0.0010	0.0010	NGR	NGR	-	-	-	-	-	-	-	-
Diethanolamine	0.060	0.060	0.060	-	-	-	-	65,000	5.0	-	-	-	-
Diethylene glycol	6.0	6.0	6.0	-	-	-	-	4,000	200	-	-	-	-
Diisopropanolamine	1.6	1.6	3.6	-	-	160	160	1.6	1.6	3.2	-	-	-
Ethylene glycol	31	31	31	NGR	NGR	9,200	16,000	190	190	-	-	-	-
Hexachlorobutadiene	0.0013	0.0013	0.006	0.031	0.0013	-	-	0.0013	0.0013	-	-	-	-
Methanol	19	19	19	270,000	19,000	-	-	630	32	-	-	-	-
Methylmethacrylate	13	11	13	230	11	-	-	-	-	-	-	-	-

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life		Irrigation	Livestock	Wildlife Watering	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse	All	All	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Monoethanolamine	0.60	0.60	0.60	-	-	-	-	30,000	1.0	-	-	-	-
MTBE	0.015	0.015	0.015	6.1	0.34	-	-	10	10	-	-	-	-
Nitrilotriacetic acid	0.40	0.40	0.40	-	-	-	-	-	-	-	-	-	-
Nonylphenol + ethoxylates	0.0081	0.0081	-	-	-	0.0081	0.0081	NGR	0.61	-	-	-	-
Perfluorooctane sulfonate (PFOS)*	0.00060	0.00060	0.00060	-	-	1.4	1.4	0.0068	0.0068	-	0.060	0.052	0.052
Perfluorooctanoic acid (PFOA)*	0.00020	0.00020	0.00020	-	-	-	-	-	-	-	-	-	-
Phenol	0.0020	0.0020	0.57	73,000	3,700	110	150	0.0040	0.0040	-	0.0020	-	-
Sulfolane	0.090	0.090	0.090	-	-	1,700	2,800	50	50	0.80	-	-	-
Triethylene glycol	60	60	60	-	-	-	-	25,000	550	-	-	-	-
Trihalomethanes - total (THMs)	0.10	0.10	0.10	-	-	-	-	-	-	-	-	-	-

Notes:

- a. B[a]P TPE (Total Potency Equivalents) are calculated by multiplying the groundwater concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence Factor (PEF) to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (1999) scheme, as follow:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

---

**TABLE B-2. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR AGRICULTURAL LAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

- b. For ecological receptors only.
- c. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- d. See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014 and updates) for further guidance on aquatic life pathway.
- e. Tier 1 guideline = lowest of aquatic life guideline and all other guidelines.
- f. As S, but can be applied to undissociated H<sub>2</sub>S if concerns arise.
- g. Eco-contact guidelines from Stantec (2012)
- h. Eco-contact guidelines from Stantec (2008)
- i. Guideline for protection of aquatic life (fine soil) is set at the maximum concentration of trichloromethane that will support biological degradation (MEMS, 2016).
- j. Total means all chemical species.
- k. As the toxicological effects of PFOA and perfluorooctanoyl sulfonate (PFOS) are considered to be additive, the sum of the ratios of the detected concentrations to the corresponding MACs for PFOS and PFOA should not exceed 1.

NGR - no guideline required, calculated value > solubility or >1,000,000 mg/L

Potable GW = protection of groundwater for potable drinking water

Inhalation = protection of volatilization from groundwater and migration into indoor air

Eco Soil Contact = protection of terrestrial plants and soil invertebrates in areas with shallow groundwater

Aquatic Life = protection of groundwater discharging to a surface water body hosting aquatic life

Irrigation = protection of a potential irrigation groundwater source

Livestock Watering = protection of a potential livestock watering groundwater resource

Wildlife Watering = protection of groundwater discharging to a surface water body from which wildlife may drink

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
<b>General and Inorganic Parameters</b>									
pH	6.5-8.5	6.5-8.5	6.5-8.5	-	-	-	-	6.5-9.0	6.5-9.0
Ammonia	see note d	see note d	-	-	-	-	-	see note d	see note d
Bromate	0.010	0.010	0.010	-	-	-	-	-	-
Chloride	120	120	250	-	-	-	-	120	120
Cyanide (free)	0.0050	0.0050	0.20	-	-	-	-	0.0050	0.0050
Fluoride	1.5	1.5	1.5	-	-	-	-	-	-
Nitrate (as nitrogen)	3.0	3.0	10	-	-	-	-	3.0	3.0
Nitrite (as nitrogen)	see note e	see note e	1.0	-	-	-	-	see note d	see note d
Sodium	200	200	200	-	-	-	-	-	-
Sulphate	see note e	see note e	500	-	-	-	-	see note d	see note d
Sulphide - Total (as S) <sup>f</sup>	0.0019	0.0019	0.050	-	-	-	-	0.0019	0.0019
Total Dissolved Solids (TDS)	500	500	500	-	-	-	-	-	-
<b>Metals</b>									
Aluminum	see note d	see note d	2.9	-	-	-	-	see note d	see note d
Antimony	0.0060	0.0060	0.0060	-	-	-	-	-	-
Arsenic	0.0050	0.0050	0.010	-	-	-	-	0.0050	0.0050
Barium	2.0	2.0	2.0	-	-	-	-	-	-
Boron	1.5	1.5	5	-	-	-	-	1.5	1.5
Cadmium	see note e	see note e	0.0070	-	-	-	-	see note d	see note d
Chromium (trivalent)	0.0089	0.0089	-	-	-	-	-	0.0089	0.0089
Chromium (hexavalent)	0.0010	0.0010	-	-	-	-	-	0.0010	0.0010
Chromium (total) <sup>j</sup>	0.050	0.050	0.050	-	-	-	-	-	-

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

This table must not be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Copper	0.0070	0.0070	1.0	-	-	-	-	0.0070	0.0070
Iron	0.30	0.30	0.30	-	-	-	-	0.30	0.30
Lead	see note e	see note e	0.0050	-	-	-	-	see note d	see note d
Manganese	0.020	0.020	0.020	-	-	-	-	-	-
Mercury (total) <sup>i</sup>	0.000005	0.000005	0.001	-	-	-	-	0.000005	0.000005
Nickel	see note d	see note d	-	-	-	-	-	see note d	see note d
Selenium	0.0020	0.0020	0.050	-	-	-	-	0.0020	0.0020
Silver	0.00010	0.00010	-	-	-	-	-	0.00010	0.00010
Uranium	0.015	0.015	0.020	-	-	-	-	0.015	0.015
Zinc	0.030	0.030	5.0	-	-	-	-	0.030	0.030
<b>Hydrocarbons</b>									
Benzene	0.0050	0.0050	0.0050	0.57	0.030	100	61	3.6	0.074
Toluene	0.024	0.021	0.024	NGR	45	82	59	12,000	0.021
Ethylbenzene	0.0016	0.0016	0.0016	NGR	31	42	20	NGR	41
Xylenes	0.020	0.020	0.020	44	2.2	21	31	NGR	2.9
Styrene	0.072	0.072	2.8	NGR	39	-	-	0.072	0.072
F1	2.2	0.81	2.2	19	0.81	6.5	7.1	NGR	9.8
F2	1.1	1.1	1.1	NGR	1.5	1.8	1.8	NGR	1.3
Acenaphthene	0.0060	0.0058	1.4	NGR	NGR	-	-	0.0060	0.0058
Anthracene	0.0034	0.000012	NGR	NGR	NGR	0.025	0.025	0.0034	0.000012
Fluoranthene	0.24	0.000057	NGR	NGR	NGR	0.24	0.24	NGR	0.000057
Fluorene	0.0042	0.0030	0.94	NGR	NGR	-	-	0.0042	0.0030
Naphthalene	0.0010	0.0010	0.47	NGR	2.7	-	-	0.0010	0.0010
Phenanthrene	0.00086	0.00040	-	-	-	-	-	0.00086	0.00040

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Pyrene	0.71	0.000092	0.71	NGR	NGR	-	-	NGR	0.000092
Carcinogenic PAHs (as B(a)P TPE) <sup>a</sup>	0.000040	0.000040	0.000040	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	-	-	NGR	NGR
Benzo[b+j]fluoranthene	-	-	-	-	-	-	-	-	-
Benzo[k]fluoranthene	-	-	-	-	-	-	-	-	-
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>b</sup>	0.0018	0.0018	-	-	-	0.0018	0.0018	NGR	NGR
Chrysene	-	-	-	-	-	-	-	-	-
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	-	-
<b>Halogenated Aliphatics</b>									
Vinyl chloride	0.0020	0.0011	0.0020	0.018	0.0011	-	-	-	-
1,1-Dichloroethene	0.014	0.014	0.014	0.68	0.039	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.0050	0.00032	0.0050	0.0064	0.00032	4.4	5.0	0.27	0.029
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.010	0.010	0.25	0.012	-	-	0.11	0.11
1,2-Dichloroethane	0.0050	0.0050	0.0050	0.17	0.010	-	-	0.10	0.10
Dichloromethane (Methylene chloride)	0.050	0.050	0.050	17	0.94	-	-	0.098	0.098
Trichloromethane (Chloroform) <sup>i</sup>	0.080	0.018	0.080	7.1	0.41	-	-	0.10	0.018
Tetrachloromethane (Carbon tetrachloride)	0.0020	0.0015	0.0020	0.030	0.0015	-	-	0.013	0.013
Dibromochloromethane	0.19	0.19	0.19	26	1.1	-	-	-	-
<b>Chlorinated Aromatics</b>									
Chlorobenzene	0.0013	0.0013	0.03	0.30	0.014	-	-	0.0013	0.0013
1,2-Dichlorobenzene	0.00070	0.00070	0.0030	116	5.4	-	-	0.00070	0.00070

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
1,4-Dichlorobenzene	0.0010	0.0010	0.0010	2.8	0.13	-	-	0.026	0.026
1,2,3-Trichlorobenzene	0.0080	0.0080	0.014	0.80	0.032	-	-	0.0080	0.0080
1,2,4-Trichlorobenzene	0.015	0.015	0.015	0.71	0.028	-	-	0.024	0.024
1,3,5-Trichlorobenzene	0.014	0.014	0.014	0.38	0.015	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.0018	0.0018	0.032	NGR	0.14	-	-	0.0018	0.0018
1,2,3,5-Tetrachlorobenzene	0.0038	0.0038	0.0038	0.41	0.017	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.0020	0.0020	0.0020	0.21	0.0088	-	-	-	-
Pentachlorobenzene	0.0094	0.0069	0.0094	NGR	0.038	-	-	NGR	0.0069
Hexachlorobenzene	0.00029	0.00029	0.00029	0.012	0.00050	-	-	-	-
2,4-Dichlorophenol	0.00020	0.00020	0.00030	NGR	1500	-	-	0.00020	0.00020
2,4,6-Trichlorophenol	0.0020	0.0020	0.0020	NGR	54	-	-	0.018	0.018
2,3,4,6-Tetrachlorophenol	0.0010	0.0010	0.0010	NGR	NGR	-	-	0.0010	0.0010
Pentachlorophenol	0.00051	0.00050	0.030	NGR	NGR	0.87	0.88	0.00051	0.00050
Dioxins & Furans <sup>c</sup>	0.00000012	0.00000012	0.00000012	-	-	-	-	-	-
PCBs	0.000094	0.000094	0.000094	-	-	-	-	-	-
<b>Pesticides</b>									
Aldicarb	0.001	0.001	0.0090	-	-	-	-	0.0010	0.0010
Aldrin	0.000028	0.000028	0.000028	0.056	0.0026	-	-	-	-
Atrazine and metabolites	0.0018	0.0018	0.0050	-	-	-	-	0.0018	0.0018
Azinphos-methyl (Guthion)	0.000010	0.000010	0.020	-	-	-	-	0.000010	0.000010
Bromacil <sup>d</sup>	0.0050	0.0050	0.95	-	-	0.44	0.30	0.0050	0.0050
Bromoxynil	0.0050	0.0050	0.0050	-	-	-	-	0.0050	0.0050
Carbaryl	0.00020	0.00020	0.090	-	-	-	-	0.00020	0.00020
Carbofuran	0.0018	0.0018	0.090	-	-	-	-	0.0018	0.0018



**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Chlorothalonil	0.00018	0.00018	0.14	-	-	-	-	0.00018	0.00018
Chlorpyrifos	0.0000046	0.0000020	0.090	-	-	-	-	0.0000046	0.0000020
2,4-D	0.0040	0.0040	0.10	-	-	-	-	0.0040	0.0040
DDT	0.0014	0.0014	0.0014	15	0.69	-	-	-	-
Diazinon	0.00017	0.00017	0.020	-	-	-	-	0.00017	0.00017
Dicamba	0.010	0.010	0.12	-	-	-	-	0.010	0.010
Diclofop-methyl	0.0090	0.0061	0.0090	-	-	-	-	0.56	0.0061
Dieldrin	0.000029	0.000029	0.000029	0.27	0.013	-	-	-	-
Dimethoate	0.0062	0.0062	0.02	-	-	-	-	0.0062	0.0062
Dinoseb	0.000055	0.000050	0.010	-	-	-	-	0.000055	0.000050
Diquat	0.070	0.070	0.070	-	-	-	-	-	-
Diuron	0.15	0.15	0.15	-	-	-	-	-	-
Endosulfan	0.0019	0.0000031	0.057	-	-	-	-	0.0019	0.0000031
Endrin	0.0028	0.0028	0.0028	-	-	-	-	-	-
Glyphosate	0.065	0.065	0.28	-	-	-	-	0.065	0.065
Heptachlor epoxide	0.000052	0.000052	0.000052	0.0043	0.00021	-	-	-	-
Lindane	0.000010	0.000010	0.0028	-	-	-	-	0.000010	0.000010
Linuron	0.0070	0.0070	0.019	-	-	-	-	0.0070	0.0070
Malathion	0.00010	0.00010	0.19	-	-	-	-	0.00010	0.00010
MCPA	0.0026	0.0026	0.10	-	-	-	-	0.0026	0.0026
Methoxychlor	0.90	0.00017	0.90	-	-	-	-	NGR	0.00017
Metolachlor	0.0078	0.0078	0.050	-	-	-	-	0.0078	0.0078
Metribuzin	0.0010	0.0010	0.080	-	-	-	-	0.0010	0.0010
Paraquat (as dichloride)	0.010	0.010	0.010	-	-	-	-	-	-

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Phorate	0.0020	0.0020	0.0020	-	-	-	-	-	-
Picloram	0.029	0.029	0.19	-	-	-	-	0.029	0.029
Simazine	0.010	0.010	0.010	-	-	-	-	0.010	0.010
Tebuthiuron <sup>h</sup>	0.0016	0.0016	0.66	-	-	0.20	0.25	0.0016	0.0016
Terbufos	0.0010	0.0010	0.0010	-	-	-	-	-	-
Toxaphene	0.00043	0.00043	0.00043	6.4	0.31	-	-	-	-
Triallate	0.00024	0.00024	0.12	-	-	-	-	0.00024	0.00024
Trifluralin	0.0012	0.00020	0.045	-	-	-	-	0.0012	0.00020
<b>Other Organics</b>									
Aniline	0.0022	0.0022	0.066	59	2.8	-	-	0.0022	0.0022
Di- <i>n</i> -butyl phthalate	0.019	0.019	0.59	NGR	NGR	-	-	0.019	0.019
Dichlorobenzidine	0.0010	0.0010	0.0010	NGR	NGR	-	-	-	-
Diethanolamine	0.060	0.060	0.060	-	-	-	-	65,000	5.0
Diethylene glycol	6.0	6.0	6.0	-	-	-	-	4,000	200
Diisopropanolamine	1.6	1.6	3.6	-	-	160	160	1.6	1.6
Ethylene glycol	31	31	31	NGR	NGR	9,200	16,000	190	190
Hexachlorobutadiene	0.0013	0.0013	0.006	0.031	0.0013	-	-	0.0013	0.0013
Methanol	19	19	19	270,000	19,000	-	-	630	32
Methylmethacrylate	13	11	13	230	11	-	-	-	-
Monoethanolamine	0.60	0.60	0.60	-	-	-	-	30,000	1.0
MTBE	0.015	0.015	0.015	6.1	0.34	-	-	10	10
Nitrilotriacetic acid	0.40	0.40	0.40	-	-	-	-	-	-
Nonylphenol + ethoxylates	0.0081	0.0081	-	-	-	0.0081	0.0081	NGR	0.61
Perfluorooctane sulfonate (PFOS)	0.00060	0.00060	0.00060	-	-	1.4	1.4	0.0068	0.0068

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Water Use	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Perfluorooctanoic acid (PFOA)	0.00020	0.00020	0.00020	-	-	-	-	-	-
Phenol	0.0040	0.0040	0.57	73,000	3,700	110	150	0.0040	0.0040
Sulfolane	0.090	0.090	0.090	-	-	1,700	2,800	50	50
Triethylene glycol	60	60	60	-	-	-	-	25,000	550
Trihalomethanes - total (THMs)	0.10	0.10	0.10	-	-	-	-	-	-

Notes:

- a. B[a]P TPE (Total Potency Equivalents) are calculated by multiplying the groundwater concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (1999) scheme, as follows:

Carcinogenic PAH Compound	PEF
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

- b. For ecological receptors only.
- c. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- d. See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014 and updates) for further guidance on aquatic life pathway.
- e. Tier 1 guideline = lowest of aquatic life guideline and potable GW guideline.
- f. As S, but can be applied to undissociated H<sub>2</sub>S if concerns arise.
- g. Eco-contact guidelines from Stantec (2012)

---

**TABLE B-3. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR RESIDENTIAL/PARKLAND - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation**, unless required due to land use considerations (see Sections 3.2 and 5.1.1). Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

- h. Eco-contact guidelines from Stantec (2008)
- i. Guideline for protection of aquatic life (fine soil) is set at the maximum concentration of trichloromethane that will support biological degradation (MEMS, 2016).
- j. Total means all chemical species.
- k. As the toxicological effects of PFOA and perfluorooctanoyl sulfonate (PFOS) are considered to be additive, the sum of the ratios of the detected concentrations to the corresponding MACs for PFOS and PFOA should not exceed 1.

NGR - no guideline required, calculated value > solubility or >1,000,000 mg/L

Potable GW = protection of groundwater for potable drinking water

Inhalation = protection of volatilization from groundwater and migration into indoor air

Eco Soil Contact = protection of terrestrial plants and soil invertebrates in areas with shallow groundwater

Aquatic Life = protection of groundwater discharging to a surface water body hosting aquatic life

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
<b>General and Inorganic Parameters</b>									
pH	6.5-8.5	6.5-8.5	6.5-8.5	-	-	-	-	6.5-9.0	6.5-9.0
Ammonia	see note d	see note d	-	-	-	-	-	see note d	see note d
Bromate	0.010	0.010	0.010	-	-	-	-	-	-
Chloride	120	120	250	-	-	-	-	120	120
Cyanide (free)	0.0050	0.0050	0.20	-	-	-	-	0.0050	0.0050
Fluoride	1.5	1.5	1.5	-	-	-	-	-	-
Nitrate (as nitrogen)	3.0	3.0	10	-	-	-	-	3.0	3.0
Nitrite (as nitrogen)	see note e	see note e	1.0	-	-	-	-	see note d	see note d
Sodium	200	200	200	-	-	-	-	-	-
Sulphate	see note e	see note e	500	-	-	-	-	see note d	see note d
Sulphide - Total (as S) <sup>f</sup>	0.0019	0.0019	0.050	-	-	-	-	0.0019	0.0019
Total Dissolved Solids (TDS)	500	500	500	-	-	-	-	-	-
<b>Metals</b>									
Aluminum	see note d	see note d	2.9	-	-	-	-	see note d	see note d
Antimony	0.0060	0.0060	0.0060	-	-	-	-	-	-
Arsenic	0.0050	0.0050	0.010	-	-	-	-	0.0050	0.0050
Barium	2.0	2.0	2.0	-	-	-	-	-	-
Boron	1.5	1.5	5	-	-	-	-	1.5	1.5
Cadmium	see note e	see note e	0.0070	-	-	-	-	see note d	see note d
Chromium (trivalent)	0.0089	0.0089	-	-	-	-	-	0.0089	0.0089
Chromium (hexavalent)	0.0010	0.0010	-	-	-	-	-	0.0010	0.0010
Chromium (total) <sup>j</sup>	0.050	0.050	0.050	-	-	-	-	-	-
Copper	0.0070	0.0070	1.0	-	-	-	-	0.0070	0.0070
Iron	0.30	0.30	0.30	-	-	-	-	0.30	0.30

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Lead	see note e	see note e	0.0050	-	-	-	-	see note d	see note d
Manganese	0.020	0.020	0.020	-	-	-	-	-	-
Mercury (total)	0.0000050	0.0000050	0.0010	-	-	-	-	0.0000050	0.0000050
Nickel	see note d	see note d	-	-	-	-	-	see note d	see note d
Selenium	0.0020	0.00200	0.05	-	-	-	-	0.0020	0.0020
Silver	0.00010	0.00010	-	-	-	-	-	0.00010	0.00010
Uranium	0.015	0.015	0.020	-	-	-	-	0.015	0.015
Zinc	0.030	0.030	5.0	-	-	-	-	0.030	0.030
<b>Hydrocarbons</b>									
Benzene	0.0050	0.0050	0.0050	3.9	0.37	540	350	3.6	0.074
Toluene	0.024	0.021	0.024	NGR	NGR	240	200	12,000	0.021
Ethylbenzene	0.0016	0.0016	0.0016	NGR	NGR	150	110	NGR	41
Xylenes	0.020	0.020	0.020	NGR	26	74	120	NGR	2.9
Styrene	0.072	0.072	2.8	NGR	NGR	-	-	0.072	0.072
F1	2.2	2.2	2.2	NGR	9.1	9.9	11	NGR	9.8
F2	1.1	1.1	1.1	NGR	17	3.1	3.1	NGR	1.3
Acenaphthene	0.0060	0.0058	1.4	NGR	NGR	-	-	0.0060	0.0058
Anthracene	0.0034	0.000012	NGR	NGR	NGR	0.32	0.32	0.0034	0.000012
Fluoranthene	0.86	0.000057	NGR	NGR	NGR	0.86	0.86	NGR	0.000057
Fluorene	0.0042	0.0030	0.94	NGR	NGR	-	-	0.0042	0.0030
Naphthalene	0.0010	0.0010	0.47	NGR	31	-	-	0.0010	0.0010
Phenanthrene	0.00086	0.00040	-	-	-	-	-	0.00086	0.00040
Pyrene	0.71	0.000092	0.71	NGR	NGR	-	-	NGR	0.000092

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Carcinogenic PAHs (as B(a)P TPE) <sup>a</sup>	0.000040	0.000040	0.000040	-	-	-	-	-	-
Benz[a]anthracene	-	-	-	-	-	-	-	NGR	NGR
Benzo[b+j]fluoranthene	-	-	-	-	-	-	-	-	-
Benzo[k]fluoranthene	-	-	-	-	-	-	-	-	-
Benzo[g,h,i]perylene	-	-	-	-	-	-	-	-	-
Benzo[a]pyrene <sup>b</sup>	0.0066	0.0066	-	-	-	0.0066	0.0066	NGR	NGR
Chrysene	-	-	-	-	-	-	-	-	-
Dibenz[a,h]anthracene	-	-	-	-	-	-	-	-	-
Indeno[1,2,3-c,d]pyrene	-	-	-	-	-	-	-	-	-
<b>Halogenated Aliphatics</b>									
Vinyl chloride	0.0020	0.0020	0.0020	0.12	0.013	-	-	-	-
1,1-Dichloroethene	0.014	0.014	0.014	4.5	0.49	-	-	-	-
Trichloroethene (Trichloroethylene, TCE)	0.0050	0.0038	0.0050	0.044	0.0038	73	83	0.27	0.029
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.010	0.010	1.8	0.14	-	-	0.11	0.11
1,2-Dichloroethane	0.0050	0.0050	0.0050	1.2	0.13	-	-	0.10	0.10
Dichloromethane (Methylene chloride)	0.050	0.050	0.050	110	12	-	-	0.098	0.098
Trichloromethane (Chloroform) <sup>i</sup>	0.080	0.018	0.080	47	5.1	-	-	0.10	0.018
Tetrachloromethane (Carbon tetrachloride)	0.0020	0.0020	0.0020	0.21	0.018	-	-	0.013	0.013
Dibromochloromethane	0.19	0.19	0.19	250	10	-	-	-	-
<b>Chlorinated Aromatics</b>									
Chlorobenzene	0.0013	0.0013	0.030	2.2	0.18	-	-	0.0013	0.0013
1,2-Dichlorobenzene	0.00070	0.00070	0.0030	NGR	64	-	-	0.00070	0.00070
1,4-Dichlorobenzene	0.0010	0.0010	0.0010	20	1.6	-	-	0.026	0.026
1,2,3-Trichlorobenzene	0.0080	0.0080	0.014	6.9	0.33	-	-	0.0080	0.0080
1,2,4-Trichlorobenzene	0.015	0.015	0.015	6.1	0.29	-	-	0.024	0.024

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
1,3,5-Trichlorobenzene	0.014	0.014	0.014	3.3	0.15	-	-	-	-
1,2,3,4-Tetrachlorobenzene	0.0018	0.0018	0.032	NGR	NGR	-	-	0.0018	0.0018
1,2,3,5-Tetrachlorobenzene	0.0038	0.0038	0.0038	NGR	0.16	-	-	-	-
1,2,4,5-Tetrachlorobenzene	0.0020	0.0020	0.0020	NGR	0.080	-	-	-	-
Pentachlorobenzene	0.0094	0.0069	0.0094	NGR	0.44	-	-	NGR	0.0069
Hexachlorobenzene	0.00029	0.00029	0.00029	0.086	0.0057	-	-	-	-
2,4-Dichlorophenol	0.00020	0.00020	0.00030	NGR	NGR	-	-	0.00020	0.00020
2,4,6-Trichlorophenol	0.0020	0.0020	0.0020	NGR	540	-	-	0.018	0.018
2,3,4,6-Tetrachlorophenol	0.0010	0.0010	0.0010	NGR	NGR	-	-	0.0010	0.0010
Pentachlorophenol	0.00051	0.00050	0.030	NGR	NGR	2.2	2.2	0.00051	0.00050
Dioxins & Furans °	0.00000012	0.00000012	0.00000012	-	-	-	-	-	-
PCBs	0.000094	0.000094	0.000094	-	-	-	-	-	-
<b>Pesticides</b>									
Aldicarb	0.0010	0.0010	0.0090	-	-	-	-	0.0010	0.0010
Aldrin	0.000028	0.000028	0.000028	0.63	0.024	-	-	-	-
Atrazine and metabolites	0.0018	0.0018	0.0050	-	-	-	-	0.0018	0.0018
Azniphos-methyl (Guthion)	0.000010	0.000010	0.020	-	-	-	-	0.000010	0.000010
Bromacil °	0.0050	0.0050	0.95	-	-	1.1	0.50	0.0050	0.0050
Bromoxynil	0.0050	0.0050	0.0050	-	-	-	-	0.0050	0.0050
Carbaryl	0.00020	0.00020	0.090	-	-	-	-	0.00020	0.00020
Carbofuran	0.0018	0.0018	0.090	-	-	-	-	0.0018	0.0018
Chlorothalonil	0.00018	0.00018	0.14	-	-	-	-	0.00018	0.00018
Chlorpyrifos	0.0000046	0.0000020	0.090	-	-	-	-	0.0000046	0.0000020
2,4-D	0.0040	0.0040	0.10	-	-	-	-	0.0040	0.0040
DDT	0.0014	0.0014	0.0014	170	6.3	-	-	-	-



**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Diazinon	0.00017	0.00017	0.020	-	-	-	-	0.00017	0.00017
Dicamba	0.010	0.010	0.12	-	-	-	-	0.010	0.010
Diclofop-methyl	0.0090	0.0061	0.0090	-	-	-	-	0.56	0.0061
Dieldrin	0.000029	0.000029	0.000029	3.0	1.2	-	-	-	-
Dimethoate	0.0062	0.0062	0.020	-	-	-	-	0.0062	0.0062
Dinoseb	0.000055	0.000050	0.010	-	-	-	-	0.000055	0.000050
Diquat	0.070	0.070	0.070	-	-	-	-	-	-
Diuron	0.15	0.15	0.15	-	-	-	-	-	-
Endosulfan	0.0019	0.0000031	0.057	-	-	-	-	0.0019	0.0000031
Endrin	0.0028	0.0028	0.0028	-	-	-	-	-	-
Glyphosate	0.065	0.065	0.28	-	-	-	-	0.065	0.065
Heptachlor epoxide	0.000052	0.000052	0.000052	0.051	0.0020	-	-	-	-
Lindane	0.000010	0.000010	0.0028	-	-	-	-	0.000010	0.000010
Linuron	0.0070	0.0070	0.019	-	-	-	-	0.0070	0.0070
Malathion	0.00010	0.00010	0.19	-	-	-	-	0.00010	0.00010
MCPA	0.0026	0.0026	0.10	-	-	-	-	0.0026	0.0026
Methoxychlor	0.90	0.00017	0.90	-	-	-	-	NGR	0.00017
Metolachlor	0.0078	0.0078	0.050	-	-	-	-	0.0078	0.0078
Metribuzin	0.0010	0.0010	0.080	-	-	-	-	0.0010	0.0010
Paraquat (as dichloride)	0.010	0.010	0.010	-	-	-	-	-	-
Phorate	0.0020	0.0020	0.0020	-	-	-	-	-	-
Picloram	0.029	0.029	0.19	-	-	-	-	0.029	0.029
Simazine	0.010	0.010	0.010	-	-	-	-	0.010	0.010
Tebuthiuron <sup>h</sup>	0.0016	0.0016	0.66	-	-	2.6	3.2	0.0016	0.0016
Terbufos	0.0010	0.0010	0.0010	-	-	-	-	-	-
Toxaphene	0.00043	0.00043	0.00043	75	2.9	-	-	-	-

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

This table must **not** be used for Tier 1 assessment and remediation. Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Pathway	Lowest Guideline		Potable	Inhalation		Eco Soil Contact		Aquatic Life	
Soil Type	Fine	Coarse	All	Fine	Coarse	Fine	Coarse	Fine	Coarse
Unit	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)
Triallate	0.00024	0.00024	0.12	-	-	-	-	0.00024	0.00024
Trifluralin	0.0012	0.00020	0.045	-	-	-	-	0.0012	0.00020
<b>Other Organics</b>									
Aniline	0.0022	0.0022	0.066	420	33	-	-	0.0022	0.0022
Di- <i>n</i> -butyl phthalate	0.019	0.019	0.59	NGR	NGR	-	-	0.019	0.019
Dichlorobenzidine	0.0010	0.0010	0.0010	NGR	NGR	-	-	-	-
Diethanolamine	0.060	0.060	0.060	-	-	-	-	65,000	5.0
Diethylene glycol	6.0	6.0	6.0	-	-	-	-	4,000	200
Diisopropanolamine	1.6	1.6	3.6	-	-	320	320	1.6	1.6
Ethylene glycol	31	31	31	NGR	NGR	15,000	26,000	190	190
Hexachlorobutadiene	0.0013	0.0013	0.0060	0.22	0.015	-	-	0.0013	0.0013
Methylmethacrylate	13	13	13	1,600	140	-	-	-	-
Methanol	19	19	19	NGR	250,000	-	-	630	32
Monoethanolamine	0.60	0.60	0.60	-	-	-	-	30,000	1.0
MTBE	0.015	0.015	0.015	40	4.3	-	-	10	10
Nitritotriacetic acid	0.40	0.40	0.40	-	-	-	-	-	-
Nonylphenol + ethoxylates	0.020	0.020	-	-	-	0.020	0.020	NGR	0.61
Perfluorooctane sulfonate (PFOS) <sup>k</sup>	0.00060	0.00060	0.00060	-	-	8.2	8.2	0.0068	0.0068
Perfluorooctanoic acid (PFOA) <sup>k</sup>	0.00020	0.00020	0.00020	-	-	-	-	-	-
Phenol	0.0040	0.0040	0.57	NGR	45,000	730	1000	0.0040	0.0040
Sulfolane	0.090	0.090	0.090	-	-	3,400	5,700	50	50
Triethylene glycol	60	60	60	-	-	-	-	25,000	550
Trihalomethanes - total (THMs)	0.10	0.10	0.10	-	-	-	-	-	-

---

**TABLE B-4. GROUNDWATER REMEDIATION GUIDELINE VALUES FOR COMMERCIAL/INDUSTRIAL - ALL WATER USES**

**This table must not be used for Tier 1 assessment and remediation.** Tier 1 groundwater guidelines are found in Table 2. This table is provided to assist Tier 2 guideline development, using the procedures outlined in the companion Tier 2 document (ESRD 2007 as amended).

Notes:

- a. B[a]P TPE (Total Potency Equivalents) are calculated by multiplying the groundwater concentration of individual carcinogenic PAHs by a standardized Benzo[a]pyrene Potency Equivalence to produce a Benzo[a]pyrene relative potency concentration, and by subsequently summing the relative potency concentrations for the entire PAH mixture. B[a]P PEFs are order of magnitude estimates of carcinogenic potential and are based on the World Health Organization (1999) scheme, as follows:

<b>Carcinogenic PAH Compound</b>	<b>PEF</b>
Benz[a]anthracene	0.1
Benzo(b+j)fluoranthene	0.1
Benzo[k]fluoranthene	0.1
Benzo[ghi]perylene	0.01
Benzo[a]pyrene	1
Chrysene	0.01
Dibenz[a,h]anthracene	1
Indeno[1,2,3-c,d]pyrene	0.1

- b. For ecological receptors only.
- c. Expressed as toxic equivalents (TEQs) based on 2,3,7,8-PCDD (See CCME, 1999 and updates)
- d. See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014 and updates) for further guidance on aquatic life pathway.
- e. Tier 1 guideline = lowest of aquatic life guideline and potable GW guideline.
- f. As S, but can be applied to undissociated H<sub>2</sub>S if concerns arise.
- g. Eco-contact guidelines from Stantec (2012)
- h. Eco-contact guidelines from Stantec (2008)
- i. Guideline for protection of aquatic life (fine soil) is set at the maximum concentration of trichloromethane that will support biological degradation (MEMS, 2016).
- j. Total means all chemical species.
- k. As the toxicological effects of PFOA and perfluorooctanoyl sulfonate (PFOS) are considered to be additive, the sum of the ratios of the detected concentrations to the corresponding MACs for PFOS and PFOA should not exceed 1.

NGR - no guideline required, calculated value > solubility or >1,000,000 mg/L

Potable GW = protection of groundwater for potable drinking water

Inhalation = protection of volatilization from groundwater and migration into indoor air

Eco Soil Contact = protection of terrestrial plants and soil invertebrates in areas with shallow groundwater

Aquatic Life = protection of groundwater discharging to a surface water body hosting aquatic life

# Appendix C

## Protocols for Calculating Alberta Tier 1 Soil and Groundwater Remediation Guidelines

### Contents

1. Introduction .....	134
1.1 Approach .....	134
2. Parameter Values .....	135
2.1 Non-Chemical-Specific Parameters.....	135
2.2 Chemical-Specific Parameters.....	136
3. Human Exposure Pathways.....	139
3.1 Direct Contact.....	139
3.2 Inhalation .....	140
3.3 Offsite Migration.....	143
4. Ecological Exposure Pathways .....	144
4.1 Direct Contact.....	144
4.2 Nutrient and Energy Cycling .....	144
4.3 Soil and Food Ingestion .....	144
4.4 Offsite Migration.....	145
5. Groundwater Pathways.....	146
5.1 Soil Remediation Guidelines.....	146
5.2 Groundwater Remediation Guidelines .....	150
6. Other Considerations .....	151
7. References.....	152

# LIST OF TABLES

Table C-1	Human Receptor Characteristics
Table C-2	Soil and Hydrogeological Parameters
Table C-3	Site Characteristics
Table C-4	Building Parameters
Table C-5	Livestock and Wildlife Receptor Characteristics
Table C-6	Chemical Parameters
Table C-7	Human Toxicity Reference Values
Table C-8	Human Absorption Factors
Table C-9	Human Background Exposure Parameters
Table C-10	Petroleum Hydrocarbon Subfraction Distribution
Table C-11	Surface Water Quality Guidelines

# 1. Introduction

This Appendix provides the protocols, parameter values, and equations used to determine the numerical values for Alberta Tier 1 Soil and Groundwater Remediation Guidelines for each of the exposure pathways described in the main text.

## 1.1 Approach

CCME is a primary source of guidance for use in Alberta. For development of Tier 1 guidelines, Alberta uses *A Protocol for the Derivation of Environmental and Human Health Soil Quality Guidelines* (CCME, 2006a) and *A Protocol for the Derivation of Groundwater Quality Guidelines for Use at Contaminated Sites* (CCME 2015) and *A Protocol for the Derivation of Soil Vapour Quality Guidelines for Human Exposures via Inhalation of Vapours* (CCME 2014) can be seen as a starting point for developing the soil remediation guidelines. The protocols were adapted for Alberta as appropriate.

Where calculated groundwater values are greater than the solubility of the chemical in question, the guideline value is replaced in the appropriate table with “NGR” for “no guideline required”

Parameters used for modelling in Alberta are listed in the tables below. Parameters have been largely adapted from the CCME protocols with some exceptions. See the *Alberta Contaminated Sites Policy Framework* (AEP, 2014) for specific details.

## 2. Parameter Values

Parameter values used in the models that calculate the Alberta Tier 1 Soil and Groundwater Remediation Guidelines fall into two main groups: i) parameters relating to receptor exposure and properties of the site, referred to as “non-chemical-specific parameters”; and, ii) parameters that relate to the chemical properties, toxicity, or background exposure to chemicals, referred to as “chemical-specific parameters”. These two groups of parameters are discussed below.

### 2.1 Non-Chemical-Specific Parameters

Parameter values used in the models in this section relating to human receptor characteristics, soil and hydrogeological parameters, site characteristics, building parameters, and livestock and wildlife receptor characteristics are summarized in Tables C-1 to C-5, respectively. All parameters are adopted directly from CCME with a few exceptions. See the *Contaminated Sites Policy Framework* (AEP, 2014). for more information. The main differences that effect Tier 1 tables are listed here.

In Table C-2, the saturated hydraulic conductivity for fine soils is 32 m/year for all exposure pathways, the lateral hydraulic gradient is 0.028, and the soil permeability to vapour flow is  $10^{-9}$  cm<sup>2</sup> for fine soil and  $6 \times 10^{-8}$  cm<sup>2</sup> for coarse soil. These changes from parameter values in CCME (2006a) are based on decisions made by the Contaminated Sites Working Group of the CCME, and reflect the deliberations that took place during the process of revising the PHC CWS (CCME, 2008a). The groundwater recharge rate is 0.012 m/year in fine soil and 0.06 m/year in coarse soil. Groundwater recharge rates are based on extensive research by Atomic Energy of Canada Limited (AECL, 1990, 1991, 1992, and 1995) and hydrogeological literature for the prairies (e.g., Trudell, 1994; Keller *et al.*, 1986; Woo and Rowsell, 1993). This work is consistent with groundwater recharge being less than 10% of precipitation in Coarse soils and less than 2% of precipitation in Fine soils. Taking these values together with precipitation data for the wettest of the Alberta locations for which climate normals are available (Edson: precipitation = 568 mm/year, based on 1961 to 1990 Canadian Climate Normals, Environment Canada, 2001) and rounding up, gives recharge values of 60 mm/year, and 12 mm/year for Coarse and Fine soils, respectively.

In Table C-4, the height for a residential building is 360 cm, and the air exchange rates for residential and commercial buildings are 0.5 and 0.9 exchanges/hour, respectively. These changes from parameter values in CCME (2006a) are consistent with CCME (2008a) and CCME (2014). In addition, for petroleum hydrocarbons only, there is an adjustment factor of 10 applied in the calculation of the indoor infiltration and inhalation guideline, reflecting empirical evidence that concentrations of petroleum hydrocarbon vapours in buildings are typically less than predicted values by this factor or more due to biodegradation in the subsurface and conservatism in the partitioning relationship as it applies to these substances. This is consistent with CCME (2014). No adjustment factor is used for non-petroleum hydrocarbons, since no empirical evidence is available to support such a factor.

In Table C-5, the cattle body weight and water ingestion rate are the same as were used in CCME (2006a), and the soil ingestion rate is taken from NRC (1996). The meadow vole was selected as the surrogate ecological receptor that would be protective of the majority of wildlife species, based on its small home range and high soil and water ingestion rates relative to its bodyweight. Meadow vole body weight and water ingestion rate were taken from USEPA (1993). Meadow vole soil ingestion rate was calculated from data and equations provided in USEPA (1993) as follows:

$$SIR = FIR \times PSID$$

and

$$FIR = 0.0687 \times BW^{0.822} \times 1000$$

Where: SIR = soil ingestion rate for meadow voles (g/day);  
PSID = proportion of soil in diet for meadow voles (0.024 – USEPA, 1993);  
FIR = food ingestion rate for meadow voles (2.41 g dry weight/day; calculated below);  
BW = meadow vole body weight (0.017 kg – minimum value from the range given in USEPA, 1993); and,  
1000 = conversion factor from kg to g.

Substituting these values in the above equation yields value for FIR of 2.41 g dry weight/day and for SIR of 0.058 g day (Table C-5).

## 2.2 Chemical-Specific Parameters

### 2.2.1 Physical Parameters

Physical parameters for the Alberta Tier 1 substances are summarized in Table C-6, together with an indication of the source for each. The parameters  $K_{oc}$  (organic carbon partition coefficient),  $H'$  (dimensionless Henry's Law constant), and  $D_{air}$  (diffusion coefficient in air) are used in the soil and groundwater guideline models. Solubility was also included, to enable groundwater guideline values greater than solubility to be identified. References for each source are provided at the end of this appendix.

### 2.2.2 Toxicological Parameters

Human toxicity reference values (TRVs) for the Alberta Tier 1 substances are summarized in Table C-7, together with an indication of the source for each. For threshold substances, the applicable TRVs are the tolerable daily intake (TDI) for oral exposure and the tolerable concentration (TC) for inhalation exposure. For non-threshold substances, the oral slope factor and the inhalation unit risk are provided in Table C-7. Where needed in the guideline calculations, the risk-specific dose and the risk-specific concentration were calculated as follows:

$$RsD = \frac{ARL}{SF}$$

$$RsC = \frac{ARL}{UR}$$

Where: RsD = risk-specific dose (mg/kg bw per day);  
ARL = acceptable risk level ( $10^{-5}$ , see below); and,  
SF = slope factor (Table C-7); (mg/kg bw per day) $^{-1}$ ;  
RsC = risk-specific concentration (mg/m<sup>3</sup>);  
UR = unit risk (Table C-7); (mg/m<sup>3</sup>) $^{-1}$ .

The acceptable risk level was set at  $10^{-5}$  based on guidance in Health Canada (Health Canada, 2004) and precedent from Alberta Environment and Parks.

TRVs are selected in accordance with *Guidance for Selecting Toxicity Reference Values for Alberta Tier 1 and Tier 2 Soil and Groundwater Remediation Guidelines* (AEP, 2017).

For volatile chemicals, if an inhalation tolerable concentration or unit risk was not available, one was estimated from the oral tolerable daily intake or the oral slope factor, using the adult body weight and inhalation rate provided in Table C-1. References for each source are provided at the end of this appendix.

### 2.2.3 Human Exposure Parameters

Human absorption factors are summarized in Table C-8.

Human background exposure parameters for the Alberta Tier 1 substances are summarized in Table C-9, together with an indication of the source for each. Note that background exposure rates are not required in the case of non-threshold chemicals. Parameter values include toddler and adult estimated daily intake (EDI), which is the daily dose of chemical that an individual might receive from all non-site-related sources (assumed to be soil, water, air, food, and consumer products for Tier 1 guideline derivation), background air concentration ( $C_a$ ), and soil allocation factor (SAF). The following sources of background exposure information were used, in order of preference:

- CCME guideline documents;
- Assessments carried out as part of the priority substance list (PSL) program under the Canadian Environmental Protection Act (CEPA); and,
- Information in the United States Agency for Toxic Substances and Disease Registry toxicological profiles.

Recognizing that a receptor might be exposed to contaminants from other sources than site-related soil, a soil allocation factor is used to ensure that exposure to contaminated soil represents only a portion of the overall allowable exposure. For the purposes of guideline calculation, exposure is assumed to be possible from five environmental compartments: soil, water, air, food, and consumer products (CCME, 2006a). Values for the soil allocation factor (SAF) were selected based on the following logic:



If a value of SAF other than the default value of 0.2 had previously been used in a guideline document, then that value was retained (e.g., hydrocarbon compounds, sulfolane, diisopropanolamine).

If the substance could exist in all five of the environmental compartments, then the default SAF of 0.2 was used.

If background exposure information was available and it was unlikely that the substance would appear in one or more of the five environmental compartments because of its chemical or physical properties, then an SAF was derived by dividing 1 by the remaining number of environmental compartments. For instance, if one compartment was removed, then an SAF of 0.25 was used to reflect exposure divided between the remaining four compartments.

References for each source are provided at the end of this appendix.

## 2.2.4 Petroleum Hydrocarbon Fractions

Petroleum hydrocarbons are a complex mixture of substances. To facilitate the calculation of risk-based soil and groundwater remediation guidelines, each fraction has been divided into several sub-fractions on the basis of chemical structure (aliphatic vs. aromatic) and carbon chain length.

Soil or groundwater remediation guidelines for each PHC fraction were developed by combining guidelines for each individual sub-fraction according to the proportion (by mass) of each sub-fraction within the fraction, according to the equation below:

$$SGRG \text{ (mg / kg – bw / day)} = \frac{1}{\sum_i \frac{F_i}{SGRG_i}}$$

Where: SGRG = soil (mg/kg) or groundwater (mg/L) remediation guideline;

$F_i$  = the assumed proportion of the fraction in soil or groundwater made up of sub-fraction  $i$  (dimensionless, see below; values in Table C-10); and,

$SGRG_i$  = soil (mg/kg) or groundwater (mg/L) remediation guideline for sub-fraction “ $i$ ” (mg/kg). For soil, maximum concentration set at  $1 \times 10^6$  mg/kg.

The assumed proportion of each sub-fraction differs for soil and groundwater, since each sub-fraction partitions differently between soil and groundwater. Assumed sub-fraction distributions for soil and groundwater are provided in Table C-10. The assumed sub-fraction distributions in soil are adopted directly from CCME (2008a). The assumed sub-fraction distributions in groundwater are calculated from the soil values by making standard equilibrium partitioning assumptions and using the following equations:

$$G_i = F_{i(\text{soil})} \frac{\rho_b}{\theta_w + (K_{oc} \times f_{oc} \times \rho_b) + (H' \times \theta_a)}$$

and

$$F_{i(\text{groundwater})} = \frac{G_i}{\sum_i G_i}$$

Where:  $G_i$  = proportion of sub-fraction  $i$  in groundwater (before normalization; dimensionless);  
 $F_{i(\text{soil})}$  = proportion of sub-fraction  $i$  in soil (Table C-10; dimensionless);  
 $F_{i(\text{groundwater})}$  = proportion of sub-fraction  $i$  in groundwater (normalized; Table C-10; dimensionless);  
 $\rho_b$  = dry soil bulk density (g/cm<sup>3</sup>);  
 $\theta_w$  = moisture-filled porosity (dimensionless);  
 $K_{oc}$  = organic carbon partition coefficient (L/kg);  
 $f_{oc}$  = fraction of organic carbon (g/g);  
 $H'$  = dimensionless Henry’s Law Constant (dimensionless); and,  
 $\theta_a$  = vapour-filled porosity (dimensionless).

Guideline values for individual exposure pathways have been capped at a maximum value of 30,000 mg/kg to maintain consistency with the *Canada-Wide Standard for Petroleum Hydrocarbons in Soil* (CCME, 2008a).

### 2.2.5 Carcinogenic Polycyclic Aromatic Hydrocarbons

Polycyclic aromatic hydrocarbons (PAHs) commonly occur as complex mixtures that may include compounds known or suspected to be carcinogenic. Soil and groundwater remediation guidelines have been calculated for benzo[a]pyrene. The benzo[a]pyrene guideline is applied to other carcinogenic PAHs through the use of Potency Equivalence Factors (PEFs). PEFs establish the toxicity of individual carcinogenic PAHs relative to benzo[a]pyrene. The notes in Appendix A and B explain the use of PEFs in more detail.

PEFs are also used in the calculation of soil guidelines for the protection of drinking water. The Risk Specific Dose (RSD) for benzo[a]pyrene is multiplied by the appropriate PEF to derive a RSD for each carcinogenic PAH. This value is used in the calculation of individual soil guidelines for the protection of drinking water as explained above in Section 5.2.4. These guidelines are then used with site-specific soil concentrations to calculate the Index of Additive Cancer Risk (IACR) as explained in the notes in Appendix A.

### 2.2.6 Water Quality Guidelines

Water quality guidelines for the Alberta Tier 1 chemicals are required as a starting point for calculating soil or groundwater remediation guidelines that are protective of groundwater uses. Values are summarized in Table C-11.

TRVs are selected in accordance with *Guidance for Selecting Toxicity Reference Values for Alberta Tier 1 and Tier 2 Soil and Groundwater Remediation Guidelines* (AEP, 2017). The following rules were applied when choosing primary reference sources for use in Alberta:

The values for human health were taken from the following sources, in order of preference:

- *Canadian Drinking Water Quality Guidelines* (HC, 2021c);
- If Canadian drinking water quality guidelines were not available, drinking water guidelines were calculated from the oral TRVs summarized in Table C-7 using the Health Canada approach (Health Canada, 1995a). The adult exposure parameters used in the calculation are summarized in Table C-1. The proportion of total intake normally ingested in water was assumed to be equivalent to the soil allocation factor (Table C-9) because both are derived in the same manner and are typically equivalent.

The values for ecological receptors were taken from the following sources, in order of preference:

- *Environmental Quality Guidelines for Alberta Surface Waters* (ESRD, 2014); and
- Livestock or wildlife watering guidelines calculated from published daily threshold exposure doses (DTEDs) and the ecological exposure parameters presented in Table C-5. DTEDs are selected in accordance with *Guidance for Selecting Toxicity Reference Values for Alberta Tier 1 and Tier 2 Soil and Groundwater Remediation Guidelines* (AEP, 2017).

### 3. Human Exposure Pathways

#### 3.1 Direct Contact

The model used to calculate the soil remediation guidelines protective of the human direct soil contact (soil ingestion, dermal contact, and particulate inhalation) is taken without change from CCME (2006a). Based on guidance in CCME (2006a), exposure via particulate inhalation was not considered for volatile compounds ( $IR_s$  was set to 0 kg/day for volatile chemicals in the equations below). Parameter values were discussed Section 2, and parameter values are summarized in Tables C-1 to C-9. Separate calculations are made for non-threshold and threshold chemicals.

##### Threshold Substances

$$SRG_{HH} = \frac{(TDI - EDI) \times SAF \times BW}{[(AF_G \times SIR) + (AF_L \times IR_s \times ET_2) + (AF_S \times SR)] \times ET_1} + [BSC]$$

Where:  $SRG_{HH}$  = human health-based soil remediation guideline (mg/kg);  
 $TDI$  = tolerable daily intake (mg/kg bw per day);  
 $EDI$  = estimated daily intake (mg/kg bw per day);  
 $SAF$  = soil allocation factor (dimensionless);  
 $BW$  = adult or toddler body weight (kg);  
 $AF_G$  = absorption factor for gut (dimensionless);  
 $AF_L$  = absorption factor for lung (dimensionless);  
 $AF_S$  = absorption factor for skin (dimensionless);  
 $SIR$  = adult or toddler soil ingestion rate (kg/day);  
 $IR_s$  = inhalation of particulate matter re-suspended from soil (kg/day);  
 $SR$  = adult or toddler soil dermal contact rate (kg/day; calculated below);  
 $ET_1$  = exposure term 1 (dimensionless) (days/week  $\div$  7  $\times$  weeks/year  $\div$  52);  
 $ET_2$  = exposure term 2 (dimensionless) (hours/day  $\div$  24); and,  
 $BSC$  = background soil concentration (mg/kg).

##### Non-Threshold Substances

$$SRG_{HH} = \frac{(RsD) \times BW}{[(AF_G \times SIR) + (AF_L \times IR_s \times ET_2) + (AF_S \times SR)] \times ET_1} + [BSC]$$

Where:  $SRG_{HH}$  = preliminary human health-based soil remediation guideline (mg/kg);  
 $R_sD$  = risk-specific dose (mg/kg bw per day);  
 $BW$  = adult body weight (kg);  
 $AF_G$  = absorption factor for gut (dimensionless);  
 $AF_L$  = absorption factor for lung (dimensionless);  
 $AF_S$  = absorption factor for skin (dimensionless);  
 $SIR$  = adult soil ingestion rate (kg/day);  
 $IR_s$  = inhalation of particulate matter re-suspended from soil (kg/day);  
 $SR$  = adult soil dermal contact rate (kg/day);  
 $ET_1$  = exposure term 1 (dimensionless) (days/week  $\div$  7  $\times$  weeks/year  $\div$  52);  
 $ET_2$  = exposure term 2 (dimensionless) (hours/day  $\div$  24); and,  
 $BSC$  = background soil concentration (mg/kg).

Note that in contrast to the CCME (2006a) protocol, an exposure term based on the exposure scenario is used for commercial and industrial land use for non-threshold substances.

The soil dermal contact rate ( $SR$ ) is the mass of contaminated soil which is assumed to contact the skin each day. This parameter is calculated as follows (CCME, 2006a):

$$SR = \{(SA_H \times DL_H) + (SA_O \times DL_O)\} \times EF$$

Where: SR = soil dermal contact rate (kg/day);  
 SA<sub>H</sub> = exposed surface area of hands (m<sup>2</sup>);  
 DL<sub>H</sub> = dermal loading of soil to hands (kg/m<sup>2</sup> per event);  
 SA<sub>O</sub> = area of exposed body surfaces other than hands (m<sup>2</sup>);  
 DL<sub>O</sub> = dermal loading of soil to other surfaces (kg/m<sup>2</sup> per event); and,  
 EF = exposure frequency (events/day).

The soil dermal contact rate is calculated separately for toddlers and adults using the parameters in Table C-1.

## 3.2 Inhalation

Soil and groundwater guidelines protective of the indoor infiltration and inhalation pathway were calculated for volatile organic compounds using the equations from the CCME (2006a) protocol without change for soil and adapted as appropriate for groundwater.

### 3.2.1 Adjustment Factor

Consistent with the approach taken in CCME (2008a, 2014), an adjustment factor of 10 is applied in the equations below for petroleum hydrocarbons (see Section 2), to account for empirical evidence that measured indoor air concentrations are typically lower by at least this factor than concentrations predicted from the models below. The adjustment factor takes the value of 1 for all non-petroleum hydrocarbon chemicals, reflecting the lack of any empirical data to support such a factor for these chemicals.

### 3.2.2 Assumptions

Assumptions implicit in the model include the following:

- contaminant vapour immediately above the groundwater table is assumed to be in equilibrium with contaminant concentrations in the groundwater based on Henry's Law;
- the soil is physically and chemically homogeneous;
- cracks in the building floor slab are filled with dry material of the underlying soil type;
- the moisture content is uniform throughout the unsaturated zone;
- decay of the contaminant source is not considered (*i.e.*, infinite source mass);
- attenuation of the contaminant in the unsaturated zone is not considered, except in the case of hydrocarbons, where the adjustment factor accounts empirically for this and other processes;
- interactions of the contaminant with other chemicals or soil minerals are not considered.

### 3.2.3 Soil

#### Threshold Substances

$$SRG_I = \frac{(TC - C_a) \times [\theta_w + (K_{oc} \times f_{oc} \times \rho_b) + (H' \times \theta_a)] \times SAF \times DF_i \times 10^3 \times AF}{H' \times \rho_b \times ET \times 10^6} + BSC$$

Where: SRG<sub>I</sub> = soil remediation guideline for indoor infiltration (mg/kg);  
 TC = tolerable concentration (mg/m<sup>3</sup>);  
 C<sub>a</sub> = background air concentration (mg/m<sup>3</sup>);  
 θ<sub>w</sub> = moisture-filled porosity (dimensionless);  
 K<sub>oc</sub> = organic carbon partition coefficient (L/kg);  
 f<sub>oc</sub> = fraction of organic carbon (g/g);  
 ρ<sub>b</sub> = dry soil bulk density (g/cm<sup>3</sup>);  
 H' = Henry's Law Constant (dimensionless);  
 θ<sub>a</sub> = vapour-filled porosity (dimensionless);  
 SAF = soil allocation factor (dimensionless);  
 DF<sub>i</sub> = dilution factor from soil gas to indoor air (calculated below);  
 10<sup>3</sup> = conversion factor from kg to g;  
 AF = adjustment factor (10, hydrocarbons; 1, non-petroleum hydrocarbons);  
 ET = exposure term (dimensionless);  
 10<sup>6</sup> = conversion factor from m<sup>3</sup> to cm<sup>3</sup>; and,

BSC = background soil concentration (mg/kg).

### Non-Threshold Substances

$$SRG_i = \frac{RsC \times [\theta_w + (K_{oc} \times f_{oc} \times \rho_b) + (H' \times \theta_a)] \times DF_i \times 10^3 \times AF}{H' \times \rho_b \times ET \times 10^6} + BSC$$

Where: SRG<sub>i</sub> = soil remediation guideline for indoor infiltration (mg/kg);  
 RsC = risk-specific concentration (mg/m<sup>3</sup>);  
 θ<sub>w</sub> = moisture-filled porosity (dimensionless);  
 K<sub>oc</sub> = organic carbon partition coefficient (L/kg);  
 f<sub>oc</sub> = fraction of organic carbon (g/g);  
 ρ<sub>b</sub> = dry soil bulk density (g/cm<sup>3</sup>);  
 H' = Henry's Law Constant (dimensionless);  
 θ<sub>a</sub> = vapour-filled porosity (dimensionless);  
 DF<sub>i</sub> = dilution factor from soil gas to indoor air (calculated below);  
 10<sup>3</sup> = conversion factor from kg to g;  
 AF = adjustment factor (10, hydrocarbons; 1, non-petroleum hydrocarbons);  
 ET = exposure term (dimensionless);  
 10<sup>6</sup> = conversion factor from m<sup>3</sup> to cm<sup>3</sup>; and,  
 BSC = background soil concentration (mg/kg).

Note that in contrast to the CCME (2006a) protocol, an exposure term of 0.2747 (corresponding to exposure to site-related contaminants for 10 hours/day, 5 days/week, and 48 weeks/year) is used for commercial and industrial land use for non-threshold substances.

### 3.2.4 Groundwater

Groundwater remediation guidelines were calculated using equations analogous to those above, but with adaptations to account for a groundwater, rather than soil, source.

### Threshold Substances

$$GWRG_i = \frac{(TC - C_a) \times SAF \times DF_i \times AF}{H' \times ET \times 10^3}$$

Where: GWRG<sub>i</sub> = groundwater remediation guideline for indoor infiltration (mg/L);  
 TC = tolerable concentration (mg/m<sup>3</sup>);  
 C<sub>a</sub> = background air concentration (mg/m<sup>3</sup>);  
 SAF = soil allocation factor (dimensionless);  
 DF<sub>i</sub> = dilution factor from soil gas to indoor air (calculated below);  
 AF = adjustment factor (10, hydrocarbons; 1, non-petroleum hydrocarbons);  
 H' = Henry's Law Constant (dimensionless);  
 ET = exposure term (dimensionless); and,  
 10<sup>3</sup> = conversion factor from m<sup>3</sup> to L.

### Non-Threshold Substances

$$GWRG_i = \frac{RsC \times DF_i \times AF}{H' \times ET \times 10^3}$$

Where: GWRG<sub>i</sub> = groundwater remediation guideline for indoor infiltration (mg/L);  
 RsC = risk-specific concentration (mg/m<sup>3</sup>);  
 DF<sub>i</sub> = dilution factor from soil gas to indoor air (calculated below);  
 AF = adjustment factor (10, hydrocarbons; 1, non-petroleum hydrocarbons);  
 H' = Henry's Law Constant (dimensionless);

ET = exposure term (dimensionless); and,  
 $10^3$  = conversion factor from  $m^3$  to L;

Note that in contrast to the CCME (2006a) protocol, an exposure term of 0.2747 (corresponding to exposure to site-related contaminants for 10 hours/day, 5 days/week, and 48 weeks/year) is used for commercial and industrial land use for non-threshold substances.

### 3.2.5 Dilution Factor Calculation

This section presents the equations (CCME, 2006a) that were used to calculate the dilution factor in the above equations. For the calculation of Tier 1 guidelines for soil and groundwater, calculated values of  $Q_{soil}$  were used rather than defaults from CCME (2014) Where direct soil vapour measurements are taken, see CCME (2014) for appropriate inputs.

The dilution factor ( $DF_i$ ) was calculated as follows:

$$DF_i = \frac{1}{\alpha}$$

Where:  $DF_i$  = dilution factor from soil gas concentration to indoor air concentration (unitless); and,  
 $\alpha$  = attenuation coefficient (unitless; see derivation below).

The attenuation coefficient,  $\alpha$ , was calculated using the following equation:

$$\alpha = \frac{\left( \frac{D_T^{eff} A_B}{Q_B L_T} \right) \exp\left( \frac{Q_{soil} L_{crack}}{D_{crack} A_{crack}} \right)}{\exp\left( \frac{Q_{soil} L_{crack}}{D_{crack} A_{crack}} \right) + \left( \frac{D_T^{eff} A_B}{Q_B L_T} \right) + \left( \frac{D_T^{eff} A_B}{Q_{soil} L_T} \right) \left[ \exp\left( \frac{Q_{soil} L_{crack}}{D_{crack} A_{crack}} \right) - 1 \right]}$$

Where:

$\alpha$  = attenuation coefficient (dimensionless);  
 $D_T^{eff}$  = effective porous media diffusion coefficient ( $cm^2/s$ ; calculated below);  
 $A_B$  = building area ( $cm^2$ );  
 $Q_B$  = building ventilation rate ( $cm^3/s$ ; calculated below);  
 $L_T$  = distance from contaminant source to foundation (cm);  
 $Q_{soil}$  = volumetric flow rate of soil gas into the building ( $cm^3/s$ ; calculated below);  
 $L_{crack}$  = thickness of the foundation (cm);  
 $D_{crack}$  = effective vapour diffusion coefficient through the crack ( $cm^2/s$ ; calculated below); and,  
 $A_{crack}$  = area of cracks through which contaminant vapours enter the building ( $cm^2$ ).

$$D_T^{eff} \approx D_a \times \left( \frac{\theta_a^{10/3}}{\theta_t^2} \right)$$

Where:  $D_T^{eff}$  = overall effective porous media diffusion coefficient based on vapour-phase concentrations for the region between the source and foundation ( $cm^2/s$ );  
 $D_a$  = diffusion coefficient in air ( $cm^2/s$ );  
 $\theta_a$  = soil vapour-filled porosity (dimensionless); and,  
 $\theta_t$  = soil total porosity (dimensionless).

$D_{crack}$  is calculated in exactly the same way as  $D_T^{eff}$ , with the exception that the assumption is made that the soil material in the cracks is dry (CCME, 2006a), and accordingly, the soil air filled porosity is the same as the soil total porosity, and the equation becomes:

$$D_{crack} \approx D_a \times \left( \frac{\theta_t^{10/3}}{\theta_t^2} \right)$$

Where:  $D_{crack}$  = effective porous media diffusion coefficient in floor cracks ( $\text{cm}^2/\text{s}$ );  
 $D_a$  = diffusion coefficient in air ( $\text{cm}^2/\text{s}$ );  
 $\theta_t$  = total porosity (dimensionless).

$$Q_B = \frac{L_B W_B H_B ACH}{3,600}$$

Where:  $Q_B$  = building ventilation rate ( $\text{cm}^3/\text{s}$ );  
 $L_B$  = building length (cm);  
 $W_B$  = building width (cm);  
 $H_B$  = building height (cm);  
ACH = air exchanges per hour ( $\text{h}^{-1}$ ); and,  
3,600 = conversion factor from hours to seconds.

$$Q_{soil} = \frac{2\pi\Delta P k_v X_{crack}}{\mu \ln \left[ \frac{2Z_{crack}}{r_{crack}} \right]}$$

Where  $Q_{soil}$  = volumetric flow rate of soil gas into the building ( $\text{cm}^3/\text{s}$ );  
 $\Delta P$  = pressure differential ( $\text{g}/\text{cm}\cdot\text{s}^2$ );  
 $k_v$  = soil vapour permeability to vapour flow ( $\text{cm}^2$ );  
 $X_{crack}$  = length of idealized cylinder (cm);  
 $\mu$  = vapour viscosity ( $0.000173 \text{ g}/\text{cm}\cdot\text{s}$ ; CCME, 2008a);  
 $Z_{crack}$  = distance below grade to idealized cylinder (cm); and,  
 $r_{crack}$  = radius of idealized cylinder (cm; calculated as  $A_{crack}/X_{crack}$ ).

### 3.3 Offsite Migration

“Offsite Migration” guidelines are calculated to check that the guideline set for commercial and industrial land use will not result in adjacent more sensitive land being contaminated at levels above the applicable guideline for the sensitive land due to wind and/or water transport of contaminated soil from the commercial or industrial site. The guideline is calculated using the equation provided in the CCME (2006a) protocol:

$$SRG_{OM} = (14.3 \times SRG_A) - (13.3 \times BSC)$$

Where  $SRG_{OM}$  = soil remediation guideline protective of offsite migration ( $\text{mg}/\text{kg}$ );  
 $SRG_A$  = soil remediation guideline for human direct soil contact for agricultural land use ( $\text{mg}/\text{kg}$ ); and,  
BSC = background soil concentration ( $\text{mg}/\text{kg}$ ).

A similar off-site migration check is calculated for ecological health using the ecological direct contact soil remediation guideline for agricultural land use.

## 4. Ecological Exposure Pathways

### 4.1 Direct Contact

#### 4.1.1 Soil

The exposure pathway considering direct contact of plants, soil invertebrates and microbes with contaminated soil is an important primary pathway in CCME (2006a), and is also important in the framework under which the Alberta Tier 1 guidelines were developed. Existing CCME soil quality guideline values were adopted for this exposure pathway. The reader is referred to CCME (2006a) for information on the protocol used to develop the existing guidelines for this exposure pathway.

For petroleum hydrocarbons, details of the guideline derivation processes are documented in CCME (2008a). Additional data allowed for the calculation of subsoil ecological direct contact guidelines for petroleum hydrocarbons. To maintain consistency with the CCME approach for benzene, toluene, ethylbenzene, and xylenes, subsoil ecocontact guidelines for petroleum hydrocarbons were set at twice the surface soil guidelines.

#### 4.1.2 Groundwater

The direct contact of shallow groundwater with plants and soil invertebrates pathway is applicable whenever groundwater is present within 3 m of the ground surface. It applies to all land uses, and is based on the corresponding soil guideline. The approach for this exposure pathway for three different chemical classes is provided below.

Non-polar organic compounds can partition between soil organic carbon, pore water, and pore vapour, based on well-established partitioning equations. The guideline for this exposure pathway for these chemicals is calculated from existing eco soil contact soil remediation guidelines using standard assumptions for the partitioning of the contaminant between soil and pore water. Separate guidelines are calculated for coarse and fine soils, using the following equation:

$$GWRG_{DC} = SRG_{DC} \frac{\rho_b}{\theta_w + (K_{oc} \times f_{oc} \times \rho_b) + (H' \times \theta_a)}$$

Where:  $GWRG_{DC}$  = groundwater remediation guideline protective of direct contact with plants and soil invertebrates in areas of shallow groundwater (mg/L);  
 $SRG_{DC}$  = soil remediation guideline protective of direct contact with plants and soil invertebrates (mg/kg);  
 $\rho_b$  = dry soil bulk density (g/cm<sup>3</sup>);  
 $\theta_w$  = moisture-filled porosity (dimensionless);  
 $K_{oc}$  = organic carbon partition coefficient (L/kg);  
 $f_{oc}$  = fraction of organic carbon (g/g);  
 $H'$  = dimensionless Henry's Law Constant (dimensionless); and,  
 $\theta_a$  = vapour-filled porosity (dimensionless).

Soil remediation guidelines protective of ecological soil contact are available for selected chemicals in Appendix A.

Groundwater guidelines for salts are not calculated in this document.

The potential interactions between metals or polar organic compounds and soils are complex in that they can be highly dependant on various environmental conditions including pH, clay mineralogy, redox conditions, and metal species present. Attempting to set groundwater guidelines for these chemicals for this pathway would involve significant uncertainty, and accordingly, it is recommended that concerns with potential adverse effects on surface soil biota from metals and polar organic compounds in shallow groundwater be addressed on a site-specific basis by analyzing soil samples.

### 4.2 Nutrient and Energy Cycling

Existing CCME soil quality guideline values were adopted for this exposure pathway. The reader is referred to CCME (2006a) for information on the protocol used to develop the existing guidelines for this exposure pathway.

### 4.3 Soil and Food Ingestion

The ingestion of contaminants in soil and contaminants bioaccumulated from soil into fodder by livestock or wildlife (soil and food ingestion pathway) forms part of the exposure scenario for natural areas or agricultural land use. Where available, existing CCME guideline values for metals for livestock soil and food ingestion were adopted without change. In general, these



guidelines specifically considered livestock species, and so were not extrapolated to wildlife. Existing CCME guidelines for polycyclic aromatic hydrocarbons, polychlorinated biphenyls (PCBs), dioxins and furans, DDT, and perfluorooctane sulfonate (PFOS) were also adopted without change for this exposure pathway. However, soil and food ingestion guidelines for these four compounds/groups considered primary, secondary and tertiary consumers, and accordingly the guidelines were considered protective of both livestock and wildlife.

Soil ingestion guidelines were calculated for benzene, toluene, ethylbenzene, and xylenes (BTEX) and petroleum hydrocarbon fractions F1 to F4. Petroleum hydrocarbons and BTEX are not considered to bioaccumulate into potential food species, and accordingly, this guideline considered only soil ingestion for petroleum hydrocarbons. The following equation was used to calculate the soil ingestion guideline for PHCs. Livestock guidelines were calculated using parameters for a cow, as an economically important livestock species. Wildlife guidelines were calculated using parameters for a meadow vole. The meadow vole was selected since small animals are typically maximally exposed to contaminants. Wildlife soil ingestion guidelines calculated using the meadow vole are expected to be protective for the majority of wildlife species.

$$SRG_{SI-L/W} = \frac{0.75 \times DTED \times BW_{L/W}}{SIR_{L/W} \times BF}$$

Where:

$SRG_{SI-L/W}$	=	soil remediation guideline for soil ingestion - livestock or wildlife (mg/kg);
0.75	=	allocation factor (dimensionless);
DTED	=	daily threshold effect dose (mg/kg-bw/day);
$BW_{L/W}$	=	body weight - livestock or wildlife (kg);
$SIR_{L/W}$	=	soil ingestion rate - livestock or wildlife (kg/day); and,
BF	=	bioavailability factor (1.0; assumed).

## 4.4 Offsite Migration

“Offsite Migration” guidelines are calculated to check that the guideline set for commercial and industrial land use will not result in adjacent more sensitive land being contaminated at levels above the applicable guideline for the sensitive land due to wind and/or water transport of contaminated soil from the commercial or industrial site. The guideline is calculated using the equation provided in the CCME (2006a) protocol:

$$SRG_{OM} = (14.3 \times SRG_A) - (13.3 \times BSC)$$

Where:

$SRG_{OM}$	=	soil remediation guideline protective of offsite migration (mg/kg);
$SRG_A$	=	soil remediation guideline for ecological direct contact for agricultural land use (mg/kg); and,
BSC	=	background soil concentration (mg/kg).

A similar off-site migration check is calculated for human health using the human direct contact soil remediation guideline for agricultural land use.

## 5. Groundwater Pathways

This section provides the protocols used to calculate soil and groundwater remediation objectives protective of exposure pathways involving groundwater. The following receptors are considered:

- humans (potable drinking water sourced from groundwater);
- livestock (drinking water from a watering well, dugout, or surface water body potentially connected to contaminated groundwater);
- agricultural crops (irrigated from potentially contaminated groundwater);
- aquatic life (via lateral groundwater transport and discharge into a surface water body); and,
- wildlife (drinking water from a surface water body potentially connected to contaminated groundwater).

In the first three cases, it is assumed that the water well or dugout could potentially be installed at any location, and hence it is assumed that there is no lateral offset between the location where the contaminated soil or groundwater is measured and the receptor.

In the last two cases, a minimum lateral separation of 10 m is assumed between the location where the contaminated soil or groundwater is measured and the receptor (location of surface water body). In cases where contamination is present within 10 m of a surface water body, a Tier 2 or Exposure Control approach is required.

Surface water quality guidelines protective of the above water uses are provided in Table C-11, where available.

### 5.1 Soil Remediation Guidelines

Soil remediation guidelines for groundwater pathways were calculated using the model and equations from the CCME (2006a) protocol. Soil remediation guidelines for the protection of groundwater are not calculated for inorganic substances due to the uncertainties associated with the partitioning of metals between the adsorbed and dissolved phase as noted above; these substances should be assessed through site-specific groundwater sampling where these pathways are applicable.

#### 5.1.1 Model Assumptions

Assumptions implicit in the model include the following:

- the soil is physically and chemically homogeneous;
- moisture content is uniform throughout the unsaturated zone;
- infiltration rate is uniform throughout the unsaturated zone;
- depletion of the contaminant source is not considered (*i.e.*, infinite source mass);
- flow in the unsaturated zone is assumed to be one dimensional and downward only (vertical recharge) with dispersion, sorption-desorption, and biological degradation;
- contaminant is not present as an immiscible phase product;
- maximum possible concentration in the leachate is equivalent to the solubility limit of the chemical in water under the defined site conditions;
- groundwater aquifer is unconfined;
- groundwater flow is uniform and steady;
- co-solubility and oxidation/reduction effects are not considered;
- attenuation of the contaminant in the saturated zone is assumed to be one-dimensional with respect to sorption-desorption, dispersion, and biological degradation;
- dispersion in groundwater is assumed to occur in the longitudinal and transverse directions only and diffusion is not considered;
- mixing of the leachate with the groundwater is assumed to occur through mixing of leachate and groundwater mass fluxes; and
- dilution of the plume by groundwater recharge down-gradient of the source is not included.

#### 5.1.2 Guideline Calculation

The soil remediation guideline protective of groundwater uses is calculated in the same way for all five groundwater uses noted at the start of this section, using the corresponding surface water quality guideline as the starting point for each, with 2 exceptions. The first exception is that the lateral offset between the point at which the contaminated soil is measured and the

surface water body (parameter “x” in the dilution factor 4 equation below) is assumed to be 10 m for aquatic life and wildlife watering, and zero for the other water uses. Therefore, dilution factor 4 is only active for aquatic life and wildlife watering and cannot be applied for other pathways. The second exception is that in the calculation of dilution factor 3 for the potable groundwater pathway only, the average thickness of the mixing zone ( $Z_d$ ) takes the fixed value of 2 m, reflecting the likely minimum screen length for a viable drinking water well. It should be noted that this second point reflects Alberta Environment and Parks policy and is not consistent with CCME (2006a).

The model considers four processes:

- partitioning of the substance from soil to pore water (leachate);
- transport of the leachate from the base of contamination to the groundwater table;
- mixing of the leachate with groundwater; and,
- transport of the substance in groundwater down-gradient to a discharge point.

For each of these four processes, a dilution factor was calculated (DF1 through DF4, respectively). DF1 has units of (mg/kg)/(mg/L) or L/kg. The other three dilution factors are dimensionless [units of (mg/L)/(mg/L)]. The overall dilution factor is used to calculate the soil concentration that is protective of groundwater using the following equations:

$$SRG_{GR} = SWQG_{FL} \times DF$$

$$DF = DF1 \times DF2 \times DF3 \times DF4$$

Where:  $SRG_{GR}$  = soil remediation guideline protective of groundwater pathways (mg/kg);  
 $SWQG_{FL}$  = corresponding surface water quality guideline (drinking water, aquatic life, livestock or wildlife watering, or irrigation) (mg/L);  
 $DF$  = overall dilution factor (L/kg);  
 $DF1$  = dilution factor for process 1 (L/kg);  
 $DF2$  = dilution factor for process 2 (dimensionless);  
 $DF3$  = dilution factor for process 3 (dimensionless); and,  
 $DF4$  = dilution factor for process 4 (dimensionless).

### Dilution Factor 1

Dilution factor 1 (DF1) is the ratio of the concentration of a contaminant in soil to the concentration in leachate that is in contact with the soil. This “dilution factor” represents the three phase partitioning between contaminant sorbed to soil, contaminant dissolved in pore water (*i.e.*, as leachate), and contaminant present as soil vapour. DF1 is calculated using the following equation:

$$DF1 = K_{oc} \times f_{oc} + \frac{(\theta_w + H' \times \theta_a)}{\rho_b}$$

Where:  $DF1$  = dilution factor 1 (L/kg);  
 $K_{oc}$  = organic carbon-water partition coefficient (L/kg);  
 $f_{oc}$  = fraction organic carbon (g/g);  
 $\theta_w$  = water filled porosity (dimensionless);  
 $H'$  = dimensionless Henry’s Law constant (dimensionless);  
 $\theta_a$  = air filled porosity (dimensionless); and,  
 $\rho_b$  = dry soil bulk density (g/cm<sup>3</sup>).

## Dilution Factor 2

Dilution factor 2 (DF2) is the ratio of the concentration of a contaminant in leachate that is in contact with the soil, to the concentration in pore water just above the groundwater table. DF2 takes the value 1.00 (*i.e.*, no dilution) for generic guidelines because it is assumed at Tier 1 that the contaminated soil extends down to the water table.

## Dilution Factor 3

Dilution factor 3 (DF3) is the ratio of the concentration of a chemical in pore water just above the groundwater table, to the concentration in groundwater beneath the source. This dilution factor reflects a decrease in concentration as leachate mixes with uncontaminated groundwater. DF3 is a function of groundwater velocity, infiltration rate, source length, and mixing zone thickness. The mixing zone thickness is calculated as being due to two processes: i) mixing due to dispersion, and ii) mixing due to infiltration rate. The equations used are as follows:

$$DF3 = 1 + \frac{Z_d \times V}{I \times X}$$

$$Z_d = r + s$$

$$r = 0.01 \times X$$

$$s = d_a \left\{ 1 - \exp \left( \frac{-2.178 \times X \times I}{V \times d_a} \right) \right\}$$

$$V = K \times i$$

Where: DF3 = dilution factor 3 (dimensionless);  
Z<sub>d</sub> = average thickness of mixing zone (m);  
V = Darcy velocity in groundwater (m/year);  
I = infiltration rate (m/year);  
X = length of contaminated soil parallel to groundwater flow (m);  
r = mixing depth due to dispersion (m);  
s = mixing depth due to infiltration rate (m);  
d<sub>a</sub> = unconfined aquifer thickness (m);  
K = aquifer hydraulic conductivity (m/year); and,  
i = lateral hydraulic gradient in aquifer (dimensionless).

Note that the parameter Z<sub>d</sub> takes the fixed value of 2 m for the drinking water pathway, but is calculated as above for all other pathways.

## Dilution Factor 4

Dilution factor 4 (DF4) accounts for the processes of dispersion and biodegradation as groundwater travels downgradient from beneath the source of contamination, and is the ratio of the concentration of a chemical in groundwater beneath the source, to the concentration in groundwater at a distance (10 m for Tier 1 for aquatic life and wildlife watering) downgradient of the source. For distances less than 10 m, a value of 1 should be used for DF4.

$$DF4 = \frac{4}{\exp(A) \times \operatorname{erfc}(B) \times [\operatorname{erf}(C) - \operatorname{erf}(D)]}$$

$$A = \frac{x}{2D_x} \left\{ 1 - \left( 1 + \frac{4L_s D_x}{v} \right)^{1/2} \right\}$$

$$B = \frac{x - vt \left(1 + \frac{4L_s D_x}{v}\right)^{1/2}}{2(D_x vt)^{1/2}}$$

$$C = \frac{y + Y/2}{2(D_y x)^{1/2}}$$

$$D = \frac{y - Y/2}{2(D_y x)^{1/2}}$$

$$L_s = \frac{0.6931}{t_{1/2s}} (e^{-0.07d})$$

$$v = \frac{V}{\theta_t R_s}$$

$$R_s = 1 + \frac{\rho_b K_{oc} f_{oc}}{\theta_t}$$

$$D_x = 0.1x$$

$$D_y = 0.01x$$

Where: DF4 = dilution factor 4 (dimensionless);  
erf = the error function;  
A = group A (dimensionless);  
C = group C (dimensionless);  
D = group D (dimensionless);  
x = distance to source (10 m, aquatic life and wildlife watering, 0 m other water uses);  
D<sub>x</sub> = dispersivity in the direction of groundwater flow (m);  
L<sub>s</sub> = decay constant (1/year);  
v = velocity of the contaminant (m/year);  
t = transport time (500 yr);  
y = distance to receptor perpendicular to groundwater flow (0 m);  
Y = source width perpendicular to groundwater flow (m);  
D<sub>y</sub> = dispersivity perpendicular to the direction of groundwater flow (m);  
t<sub>1/2s</sub> = decay half-life of contaminant in saturated zone of aquifer (years);  
d = water table depth (m);  
V = Darcy velocity in groundwater (m/year);  
θ<sub>t</sub> = total soil porosity (dimensionless);  
R<sub>s</sub> = retardation factor in saturated zone (dimensionless);  
ρ<sub>b</sub> = dry soil bulk density (g/cm<sup>3</sup>);  
K<sub>oc</sub> = organic carbon partition coefficient (mL/g); and,  
f<sub>oc</sub> = fraction organic carbon (g/g).

## 5.2 Groundwater Remediation Guidelines

It is assumed that a dugout could potentially be constructed at any location on agricultural land and, accordingly, the livestock watering and irrigation water quality guidelines are applicable as groundwater remediation guidelines across all agricultural land. (see Table C-10). Furthermore, it is assumed that a water well could be constructed anywhere within a Domestic Use Aquifer (DUA). Accordingly, drinking water quality guidelines must be applied as groundwater remediation guidelines within the entire DUA. Therefore any modification of these guidelines is considered Exposure Control.

For aquatic life or wildlife watering, it is assumed that there is a minimum 10 m lateral separation between the point of measurement and the surface water body.

### 5.2.1 Model Assumptions

Assumptions implicit in the model include the following:

- the soil/aquifer material in the saturated zone is physically and chemically homogeneous;
- decay of the contaminant source is not considered (*i.e.*, infinite source mass);
- the contaminant is not present as a free phase product;
- groundwater flow is uniform and steady;
- co-solubility and oxidation/reduction effects are not considered;
- dispersion is assumed to occur in the longitudinal and transverse directions only and diffusion is not considered; and,
- dilution of the plume by groundwater recharge down-gradient of the source is not included.

### 5.2.2 Guideline Calculation – Aquatic Life and Wildlife Watering

The groundwater remediation guideline protective of aquatic life and wildlife watering is calculated using the following equations.

$$\text{GWRG}_{\text{GR}} = \text{SWQG}_{\text{FL}} \times \text{DF}_4$$

Where:  $\text{GWRG}_{\text{GR}}$  = groundwater remediation guideline protective of groundwater pathways (mg/L);  
 $\text{SWQG}_{\text{FL}}$  = corresponding surface water quality guideline (aquatic life, or wildlife watering) (mg/L);  
 $\text{DF}_4$  = dilution factor for process 4 (dimensionless).

Dilution factor 4 is calculated in the same way as described above for soil remediation guidelines

For inorganic substances, the aquatic life surface water quality guidelines have been adopted directly because the processes that affect their transport in groundwater do not meet the model assumptions.

## 6. Other Considerations

In addition to the exposure pathways described above, there are a range of other considerations relating to soil and groundwater contamination management including:

- aesthetics/nuisance issues, including smell, taste, and colour.
- concerns with damage to buried infrastructure (concrete foundations, metal pilings or pipelines, fibre-optic communication cables, power cables, polymer piping and joints, etc.); and,
- safety concerns with the possible build-up of flammable vapours or the exposure of workers to dangerous atmospheres, particularly in confined spaces.

In this document, soil ecocontact guideline values that were derived under the CCME (2006a) protocol and previous versions implicitly incorporated the above soil management considerations in the ecological subsoil guidelines. However, where management limits are available that separate these considerations out, these have been adopted in the Alberta Tier 1 guidelines. Currently, management limits are only available for petroleum hydrocarbons and methanol. Please see CCME (2008a, 2017) for more information.

Regardless of whether specific numerical values have been specified for the above considerations, the proponent is responsible for ensuring that these concerns are addressed by the site remediation and risk management activities.

## 7. References

Alberta Environment and Parks references are available at:

<https://www.alberta.ca/ministry-environment-parks.aspx>.

- AECL (Atomic Energy of Canada Limited), 1990. Hydrogeology of surficial materials of permit areas D and F and the Lee River Study Area in the Whiteshell Research Area. Applied Geoscience Branch, Whiteshell Nuclear Research Establishment, Pinawa, Manitoba. TR-498, January 1990.
- AECL (Atomic Energy of Canada Limited), 1991. Hydrometeorological data for the Dead Creek Watershed, Southeastern Manitoba, 1982-1989. Applied Geoscience Branch, Whiteshell Nuclear Research Establishment, Pinawa, Manitoba. TR-540, April 1991.
- AECL (Atomic Energy of Canada Limited), 1992. Hydrogeology and hydrochemistry for the Rice Creek Watershed of the Whiteshell Research Area 1986-1990. Applied Geoscience Branch, Whiteshell Nuclear Research Establishment, Pinawa, Manitoba. TR-570, December 1992.
- AECL (Atomic Energy of Canada Limited), 1995. Hydrogeology and hydrochemistry of the Atikokan Research Area of Northwestern Ontario. Whiteshell Laboratories, Pinawa, Manitoba. TR-713, December 1992.
- AENV (Alberta Environment), 2009. Soil Remediation Guidelines for Barite: Environmental Health and Human Health. Environmental Assurance Division, Alberta Environment.
- AENV (Alberta Environment), 2010a. Soil and Groundwater Remediation Guidelines for Monoethanolamine and Diethanolamine. Environmental Assurance Division. Alberta Environment.
- AENV (Alberta Environment), 2010b. Soil and Groundwater Remediation Guidelines for Diethylene Glycol and Triethylene Glycol. Environmental Assurance Division. Alberta Environment.
- AEP (Alberta Environment and Parks), 2016. Soil Remediation Guidelines for Boron – Environmental and Human Health. Land Policy Branch. Policy and Planning Division.
- AEP (Alberta Environment and Parks), 2017. Guidance for Selecting Toxicity Reference Values for Alberta Tier 1 and Tier 2 Soil and Groundwater Remediation Guidelines. Land Policy Branch, Policy and Planning Division.
- ATSDR (Agency for Toxic Substances and Disease Registry), 1995. Toxicological Profile for Polycyclic Aromatic Hydrocarbons. U.S. Department of Health and Human Services. Public Health Service. August 1995.
- ATSDR (Agency for Toxic Substances and Disease Registry), 2005. Toxicological Profile for Naphthalene, 1-Methylnaphthalene, and 2-Methylnaphthalene. U.S. Department of Health and Human Services. Public Health Service. August 2005.
- BCMELP (British Columbia Ministry of Environment, Lands, and Parks), 1996. Overview of CSST Procedures for the Derivation of Soil Quality Matrix Standards for Contaminated Sites.
- CCME (Canadian Council of Ministers of the Environment). 1997. Canadian Soil Quality Guidelines for the Protection of Environmental and Human Health: Phenol. Fact sheet. Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 1999 and updates. Canadian Environmental Quality Guidelines. Canadian Council of Ministers of the Environment, Winnipeg. Published 1999, updated through 2006.
- CCME (Canadian Council of Ministers of the Environment), 2002. Canadian Soil Quality Guidelines for the Protection of Environmental and Human Health. Polychlorinated Dibenzo-*p*-dioxins and Polychlorinated Dibenzofurans. Fact Sheet. Canadian Council of Ministers of the Environment.
- CCME (Canadian Council of Ministers of the Environment), 2006a. A Protocol for the Derivation of Environmental and Human Health Soil Quality Guidelines. Canadian Council of Ministers of the Environment.
- CCME (Canadian Council of Ministers of the Environment), 2006b. Canadian Soil Quality Guidelines for Diisopropanolamine (DIPA) (Environmental and Human Health Effects). Scientific Supporting Document.
- CCME (Canadian Council of Ministers of the Environment), 2006c. Canadian Soil Quality Guidelines for Sulfolane (Environmental and Human Health Effects). Scientific Supporting Document.



- CCME (Canadian Council of Ministers of the Environment), 2007. Canadian Soil Quality Guidelines for Trichloroethylene (Environmental and Human Health Effects). Scientific Supporting Document. 85pp.
- CCME (Canadian Council of Ministers of the Environment), 2008a. Canada-Wide Standards for Petroleum Hydrocarbons (PHCs) in Soil: Scientific Rationale.
- CCME (Canadian Council of Ministers of the Environment), 2010. Canadian Soil Quality Guidelines Carcinogenic and Other Polycyclic Aromatic Hydrocarbons (PAHs) (Environmental and Human Health Effects). Scientific Supporting Document. PN 1445. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2015. Scientific Criteria Document for Canadian Soil Quality Guidelines for the Protection of Environmental and Human Health: Nickel. PN 1540. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2017. Scientific Criteria Document for the Development of the Canadian Soil Quality Guidelines for Methanol, Protection of Environmental and Human Health.. PN 1573. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- CCME (Canadian Council of Ministers of the Environment), 2021. Scientific Criteria Document for the Development of the Canadian Soil and Groundwater Quality Guidelines for the Protection of Environmental and Human Health: Perfluorooctane Sulfonate (PFOS). PN 1625. Canadian Council of Ministers of the Environment, Winnipeg, Manitoba.
- EC (Environment Canada), 1999a. Canadian Soil Quality Guidelines for DDT. Environmental Effects. Scientific Supporting Document. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa. December 1999.
- EC (Environment Canada), 1999b. Canadian Soil Quality Guidelines for Ethylene Glycol. Environmental Effects. Scientific Supporting Document. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa. Based on 1997 Assessment. Revised December 1999.
- EC (Environment Canada), 2000. Canadian Sediment Quality Guidelines for the Protection of Aquatic Life and Canadian Tissue Residue Guidelines for the Protection of Wildlife Consumers of Aquatic Biota: Polychlorinated Dibenzo-P-Dioxins and polychlorinated Dibenzofurans (PCDD/Fs). Technical Supporting Document. 2 vols. Report # 1-2. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa.
- EC (Environment Canada), 2001. Canadian Soil Quality Guidelines for Polychlorinated Biphenyls (PCBs). Scientific Supporting Document. Report # 1-2. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa. August, 2001.
- EC (Environment Canada), 2002. Canadian Environmental Quality Guidelines for Nonylphenol and its Ethoxylates. Scientific Supporting Document, Water Sediment and Soil. Report # 1-3. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa. August, 2002.
- EC (Environment Canada), 2004a. Canadian Soil Quality Guidelines for Benzene. Technical Supporting Document. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa.
- EC (Environment Canada), 2004b. Canadian Soil Quality Guidelines for Toluene, Ethylbenzene and Xylenes (TEX). Technical Supporting Document. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada, Ottawa.
- ESRD (Alberta Environment and Sustainable Resource Development), 2014 and updates. Environmental Quality Guidelines for Alberta Surface Waters. Water Policy Branch, Policy Division. Edmonton.
- Gustafson, J.B., Tell, J.G., and Orem, D., 1997. Total Petroleum Hydrocarbon Criteria Working Group Series, Volume 3, Selection of Representative TPH Fractions Based on Fate and Transport Considerations. Amherst Scientific Publishers, Massachusetts.
- HC (Health Canada). 1986a. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Chlorpyrifos. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1986b. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Diazinon. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1986c. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Malathion. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1986d. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Metolachlor. Health Canada. Ottawa, Ontario.

- HC (Health Canada). 1986e. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Paraquat. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1986f. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Phorate. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1986g. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Simazine. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1987a. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Chlorophenols. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1987b. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Diclofop-methyl. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1987c. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Diuron. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1987d. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Glyphosate. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1989a. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Azinophos-Methyl. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1989b. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Bromoxynil. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1989c. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Diquat. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1990. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Picloram. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1991a. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Carbaryl. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1991b. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Carbofuran. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1992a. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Trifluralin. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1993. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Atrazine. Water and Air Quality Bureau, Healthy Environments and Consumer Safety Branch, Health Canada. Ottawa, Ontario H128-1/10-620E.
- HC (Health Canada). 1995a. Guidelines for Canadian Drinking Water Quality. Part I: Approach to the Derivation of Drinking Water Guidelines. Federal–Provincial–Territorial Committee on Drinking Water, February 1995.
- HC (Health Canada). 1995b. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Terbufos. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 1996. Canadian Environmental Protection Act, Priority Substances List, Supporting Documentation: Health-Based Tolerable Daily Intakes/Concentrations and Tumourigenic Doses/Concentrations for Priority Substances (unedited version). Health Canada. Environmental Health Centre. Ottawa, Ontario. August 1996.
- HC (Health Canada). 2004. Federal Contaminated Site Risk Assessment in Canada. Part I: Guidance on Human Health Preliminary Quantitative Risk Assessment (PQRA). Prepared by: Environmental Health Assessment Services Safe Environments Programme, September 2004.
- HC (Health Canada). 2006. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Trihalomethanes. Federal–Provincial–Territorial Committee on Drinking Water.
- HC (Health Canada). 2009a. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – 2,4-dichlorophenoxyacetic Acid. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 2009b. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Dimethoate. Health Canada. Ottawa, Ontario.

- HC (Health Canada). 2010. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – 2-Methyl-4-chlorophenoxyacetic acid (MCPA). Water and Air Quality Bureau, Healthy Environments and Consumer Safety Branch, Health Canada. Ottawa, Ontario H128-1/10-620E.
- HC (Health Canada). 2014. Guidelines for Canadian Drinking Water Quality, Supporting Documents – Dicamba. Health Canada. Ottawa, Ontario.
- HC (Health Canada). 2017. Summary Document Indoor Air Reference Levels for Chronic Exposure to Volatile Organic Compounds. Health Canada, Ottawa, Ontario. October 2017.
- HC (Health Canada). 2021a. Federal Contaminated Site Risk Assessment in Canada, Toxicological Reference Values (TRVs) Version 3.0. March 2021.
- HC (Health Canada). 2021b. Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Metribuzin. Water and Air Quality Bureau, Healthy Environments and Consumer Safety Branch, Health Canada. Ottawa, Ontario. H144-13/19-2021E.
- HC (Health Canada). 2021c. Guidelines for Canadian Drinking Water - Summary Table. Water and Air Quality Bureau, Healthy Environments and Consumer Safety Branch, Health Canada, Ottawa, Ontario.
- Keller, C.K., Van der Kamp, G., and Cherry, J.A., 1986. Fracture permeability and groundwater flow in clayey till near Saskatoon, Saskatchewan. *Canadian Geotechnical Journal*, **23**, 229-240.
- MEMS (Millennium EMS Solutions Ltd.), 2016. Disinfection By-product Chloroform in Shallow Groundwater Within the City of Calgary. Prepared for the City of Calgary.
- NRC (National Research Council), 1996. Nutrient Requirements of Beef Cattle. National Academy Press, Washington, DC.
- ORNL (Oak Ridge National Laboratory), 2006. Risk Assessment Information System. On-Line Database, available at [rais.ornl.gov](https://rais.ornl.gov) (Accessed June 22, 2021)
- SRC (Syracuse Research Corporation), 2006. PhysProp Physical Properties Database, available online at [www.srcinc.com/what-we-do/efdb.aspx](http://www.srcinc.com/what-we-do/efdb.aspx) (Accessed June 22, 2021)
- Trudell, M.R., 1994. Soil salinization associated with shallow groundwater flow systems in thin, fractured till moraine. Ph.D. thesis submitted to the Earth Sciences Department of the University of Waterloo, Ontario, Canada.
- US EPA (United States Environmental Protection Agency), 1993. Wildlife Exposure Factors Handbook. Office of Research and Development. December 1993 EPA/600/R-93/187.
- US EPA (United States Environmental Protection Agency), 1994. Chemical Summary for Methyl-tert-Butyl Ether. August 1994 EPA 749 –F-94-017a.
- US EPA (United States Environmental Protection Agency), 1996a. Soil Screening Guidance: Technical Background.
- US EPA (United States Environmental Protection Agency), 1996b. Reregistration Eligibility Decision (RED) Bromacil. EPA 738-R-96-013.
- US EPA (United States Environmental Protection Agency), 2006. Integrated Risk Information System (IRIS) On-Line Database available at [www.epa.gov/iris](http://www.epa.gov/iris) (Accessed June 22, 2021)
- Woo, M.K., and Rowsell, R.D., 1993. Hydrology of a prairie slough. *Journal of Hydrology*, **146**, 175-207.
- WHO/IPCS, 1998. Environmental Health Criteria 202: Selected Non-Heterocyclic Polycyclic Aromatic Hydrocarbon. International Program on Chemical Safety, United Nations Environmental Program, World Health Organization. Geneva.

**TABLE C-1. HUMAN RECEPTOR CHARACTERISTICS**

Parameter	Symbol	Unit	Toddler	Adult
Body Weight	BW	kg	16.5	70.7
Air Inhalation Rate	IR	m <sup>3</sup> /d	9.3	15.8
Soil Inhalation Rate	IR <sub>s</sub>	kg/d	7.1 x 10 <sup>-9</sup>	1.2 x 10 <sup>-8</sup>
Water Ingestion Rate	WIR	L/d	0.6	1.5
Soil Ingestion Rate	SIR	kg/d	0.00008	0.00002
Skin Surface Area				
- Hands	SA <sub>H</sub>	m <sup>2</sup>	0.043	0.089
- Other	SA <sub>O</sub>	m <sup>2</sup>	0.258	0.25
Dermal Loading to Skin				
- Hands	DL <sub>H</sub>	kg/m <sup>2</sup> -event	0.001	0.001
- Other	DL <sub>O</sub>	kg/m <sup>2</sup> -event	0.0001	0.0001
Dermal Exposure Frequency	EF	events/d	1	1
Exposure Term, agricultural and residential/parkland				
Exposure Term, commercial and industrial	ET	-	1	1
Exposure Term, commercial and industrial	ET	-	0.2747	0.2747
Exposure Term, agricultural and residential/parkland	ET <sub>1</sub>	-	1	1
Exposure Term, commercial and industrial	ET <sub>1</sub>	-	0.6593	0.6593
Exposure Term, agricultural and residential/parkland	ET <sub>2</sub>	-	1	1
Exposure Term, commercial and industrial	ET <sub>2</sub>	-	0.4167	0.4167

Notes:

All values from CCME (2006a)

**TABLE C-2. SOIL AND HYDROGEOLOGICAL PARAMETERS**

Parameter	Symbol	Unit	Fine Soil	Coarse Soil	Notes
Soil Bulk Density	$\rho_b$	$g/cm^3$	1.4	1.7	
Soil Total Porosity	$\theta_t$	$cm^3/cm^3$	0.47	0.36	
Soil Moisture-Filled Porosity	$\theta_w$	$cm^3/cm^3$	0.168	0.119	
Soil Vapour-Filled Porosity	$\theta_a$	$cm^3/cm^3$	0.302	0.241	
Fraction of Organic Carbon	$f_{oc}$	$mass/mass$	0.005	0.005	
Saturated Hydraulic Conductivity	K	$m/y$	32	320	
Hydraulic Gradient	i	$m/m$	0.028	0.028	
Recharge (Infiltration) Rate	I	$m/y$	0.012	0.06	1
Soil Permeability to Vapour Flow	$k_v$	$cm^2$	$10^{-9}$	$6 \times 10^{-8}$	2

Notes:

All parameters values from CCME (2006a) except as noted

1. See Section 2

2. Fine grained value from CCME (2008a)

**TABLE C-3. SITE CHARACTERISTICS**

Parameter	Symbol	Unit	Value
Contaminant Source Width Perpendicular to Groundwater Flow	Y	m	10
Contaminant Source Length Parallel to Groundwater Flow	X	m	10
Contaminant Source Depth	Z	m	3
Distance to Surface Water	x	m	10
Distance to Potable Water User	x	m	0
Distance to Agricultural Water User	x	m	0
Distance from Contamination to Building Slab	$L_T$	cm	30
Depth to Groundwater (water table)	d	m	3
Depth of Unconfined Aquifer	$d_a$	m	5
Transport time	t	yr	500

Notes:

All values from CCME (2006a)

**TABLE C-4. BUILDING PARAMETERS**

Parameter	Symbol	Unit	Residential Basement		Residential Slab-on-Grade		Commercial Slab-on-Grade		Notes
			Other Hydrocarbons	Petroleum Hydrocarbons	Other Hydrocarbons	Petroleum Hydrocarbons	Other Hydrocarbons	Petroleum Hydrocarbons	
Adjustment Factor			1	10	1	10	1	10	1,2
Building Length	$L_B$	cm	1,225	1,225	1,225	1,225	2,000	2,000	
Building Width	$W_B$	cm	1,225	1,225	1,225	1,225	1,500	1,500	
Building Height (including basement)	$H_B$	cm	360	360	360	360	300	300	2
Area of Substructure	$A_B$	cm <sup>2</sup>	2.7x10 <sup>6</sup>	2.7x10 <sup>6</sup>	1.5x10 <sup>6</sup>	1.5x10 <sup>6</sup>	3.0x10 <sup>6</sup>	3.0x10 <sup>6</sup>	
Thickness of Floor Slab	$L_{crack}$	cm	11.25	11.25	11.25	11.25	11.25	11.25	
Depth of Floor Slab Below Ground	$Z_{crack}$	cm	244	244	11.25	11.25	11.25	11.25	
Distance from Source to Slab:	$L_T$	cm							
surface soil			30	30	30	30	30	30	
subsoil			30	30	139	139	139	139	
Crack Area	$A_{crack}$	cm <sup>2</sup>	994.5	994.5	994.5	994.5	1,846	1,846	
Crack Length	$X_{crack}$	cm	4,900	4,900	4,900	4,900	7,000	7,000	
Air Exchange Rate	ACH	exch/hr	0.5	0.5	0.5	0.5	0.9	0.9	2
Pressure Differential	$\Delta P$	g/cm·s <sup>2</sup>	40	40	40	40	20	20	

Notes:

All parameters values from CCME (2006a) except as noted.

1. An application factor of 10 is applied to the calculation for hydrocarbons only to account for empirical evidence of reduction in predicted indoor air concentrations. No data are available to support such a correction for non-hydrocarbons, and accordingly no application factor is used for non-hydrocarbons.
2. From CCME (2008a)

**TABLE C-5. LIVESTOCK AND WILDLIFE RECEPTOR CHARACTERISTICS**

<b>Parameter</b>	<b>Symbol</b>	<b>Unit</b>	<b>Livestock (Cow) <sup>a</sup></b>	<b>Wildlife (Meadow Vole)</b>
Body Weight	BW	kg	550 <sup>a</sup>	0.017 <sup>c</sup>
Soil Ingestion Rate	SIR	kg/d	0.747 <sup>b</sup>	0.000058 <sup>d</sup>
Water Ingestion Rate	WIR	L/d	100 <sup>a</sup>	0.00357 <sup>a</sup>

Notes:

a. CCME (2006a)

b. NRC (1996)

c. US EPA (1993)

d. calculated; see text



TABLE C-6. CHEMICAL PARAMETERS

	Koc	Source	H'	Source	Da	Source	Solubility	Source	Half Life	Source
	ml/g		dimensionless		cm2/s		mg/L		yr	
Metals										
Boron										
See AEP (2016)										
Hydrocarbons										
Benzene	81	EC (2004a)	0.225	EC (2004a)	8.80E-02	EC (2004a)	1780	Gustafson et al (1997)	1	BCMELP (1996)
Toluene	234	EC (2004b)	0.274	EC (2004b)	8.70E-02	EC (2004b)	515	Gustafson et al (1997)	0.288	BCMELP (1996)
Ethylbenzene	537	EC (2004b)	0.358	EC (2004b)	7.50E-02	EC (2004b)	152	Gustafson et al (1997)	0.312	BCMELP (1996)
Xylenes	586	EC (2004b)	0.252	EC (2004b)	7.80E-02	EC (2004b)	198	Gustafson et al (1997)	0.501	BCMELP (1996)
Styrene	461	Gustafson et al (1997)	0.123	Gustafson et al (1997)	7.10E-02	Gustafson et al (1997)	300	Gustafson et al (1997)		
F1									1.95	CCME (2008a)
F2									4.79	CCME (2008a)
Aliphatic C <sub>6</sub> -C <sub>8</sub>	3,981	CCME (2008a)	50	CCME (2008a)	5.00E-02	CCME (2008a)	5.4	CCME (2008a)		
Aliphatic C <sub>&gt;8</sub> -C <sub>10</sub>	31,623	CCME (2008a)	80	CCME (2008a)	5.00E-02	CCME (2008a)	0.43	CCME (2008a)		
Aliphatic C <sub>&gt;10</sub> -C <sub>12</sub>	251,189	CCME (2008a)	120	CCME (2008a)	5.00E-02	CCME (2008a)	0.034	CCME (2008a)		
Aliphatic C <sub>&gt;12</sub> -C <sub>16</sub>	5.01E+06	CCME (2008a)	520	CCME (2008a)	5.00E-02	CCME (2008a)	0.00076	CCME (2008a)		
Aliphatic C <sub>&gt;16</sub> -C <sub>21</sub>	6.31E+08	CCME (2008a)	4900	CCME (2008a)	5.00E-02	CCME (2008a)	0.0000025	CCME (2008a)		
Aliphatic C <sub>&gt;21</sub> -C <sub>34</sub>	1.0E+13	CCME (2008a)	5.6E+05	CCME (2008a)	5.00E-02	CCME (2008a)				
Aliphatic C <sub>&gt;34</sub>	1.6E+08	CCME (2008a)	1.2E+08	CCME (2008a)	5.00E-02	CCME (2008a)				
Aromatic C <sub>&gt;8</sub> -C <sub>10</sub>	1,585	CCME (2008a)	0.48	CCME (2008a)	5.00E-02	CCME (2008a)	65	CCME (2008a)		
Aromatic C <sub>&gt;10</sub> -C <sub>12</sub>	2,512	CCME (2008a)	0.14	CCME (2008a)	5.00E-02	CCME (2008a)	25	CCME (2008a)		
Aromatic C <sub>&gt;12</sub> -C <sub>16</sub>	5,012	CCME (2008a)	0.053	CCME (2008a)	5.00E-02	CCME (2008a)	5.8	CCME (2008a)		
Aromatic C <sub>&gt;16</sub> -C <sub>21</sub>	15,849	CCME (2008a)	0.013	CCME (2008a)	5.00E-02	CCME (2008a)	0.65	CCME (2008a)		
Aromatic C <sub>&gt;21</sub> -C <sub>34</sub>	125,893	CCME (2008a)	0.00067	CCME (2008a)	5.00E-02	CCME (2008a)	0.0066	CCME (2008a)		
Aromatic C <sub>&gt;34</sub>	1.8E+06	CCME (2008a)	0.000018	CCME (2008a)	5.00E-02	CCME (2008a)				
Acenaphthene	2,818	CCME (2010)	6.56E-03	CCME (2010)	4.21E-02	USEPA (1996a)	3.9	CCME (2010)		
Anthracene	19,953	CCME (2010)	0.0015	CCME (2010)	3.24E-02	USEPA (1996a)	0.057	CCME (2010)		
Fluoranthene	41,687	CCME (2010)	0.000609	CCME (2010)	3.03E-02	USEPA (1996a)	0.26	CCME (2010)		
Fluorene	4,898	CCME (2010)	0.00337	CCME (2010)	3.63E-02	USEPA (1996a)	1.9	CCME (2010)		

TABLE C-6. CHEMICAL PARAMETERS

	Koc	Source	H'	Source	Da	Source	Solubility	Source	Half Life	Source
	ml/g		dimensionless		cm2/s		mg/L		yr	
Naphthalene	708	CCME (2010)	0.020441	CCME (2010)	5.90E-02	USEPA (1996a)	31.7	CCME (2010)		
Phenanthrene	6,607	CCME (2010)	0.000986	CCME (2010)	2.72E-02	USEPA (1996a)	1.15	CCME (2010)		
Pyrene	69,183	CCME (2010)	0.000466	CCME (2010)	2.72E-02	USEPA (1996a)	1.35	CCME (2010)		
Benz[a]anthracene	199,526	CCME (2010)	0.000142	CCME (2010)	5.01E-02	USEPA (1996a)	0.0094	CCME (2010)		
Benzo[b+j]fluoranthene	93,325	CCME (2010)	0.000468	CCME (2010)	2.26E-02	USEPA (1996a)	0.00375	CCME (2010)		
Benzo[k]fluoranthene	19,953	CCME (2010)	0.0000351	CCME (2010)	2.26E-02	USEPA (1996a)	0.0008	CCME (2010)		
Benzo[g,h,i]perylene	407,380	CCME (2010)	0.00000597	CCME (2010)	4.48E-02	ORNL (2006)	0.000026	CCME (2010)		
Benzo[a]pyrene	2,187,762	CCME (2010)	0.0000478	CCME (2010)	4.30E-02	USEPA (1996a)	0.0016	CCME (2010)		
Chrysene	125,892	CCME (2010)	0.004	CCME (2010)	2.48E-02	USEPA (1996a)	0.00415	CCME (2010)		
Dibenz[a,h]anthracene	1,380,384	CCME (2010)	0.000000622	CCME (2010)	2.02E-02	USEPA (1996a)	0.00249	CCME (2010)		
Indeno[1,2,3-c,d]pyrene	1,584,893	CCME (2010)	0.0000677	CCME (2010)	1.90E-02	USEPA (1996a)	0.000022	CCME (2010)		
Halogenated Aliphatics										
Vinyl chloride	18.6	USEPA (1996a)	1.11	USEPA (1996a)	1.06E-01	USEPA (1996a)	2760	USEPA (1996a)		
1,1-Dichloroethene	65	USEPA (1996a)	1.07	USEPA (1996a)	1.04E-01	USEPA (1996a)	2250	USEPA (1996a)		
Trichloroethene (Trichloroethylene, TCE)	94	USEPA (1996a)	0.422	USEPA (1996a)	7.90E-02	USEPA (1996a)	1100	USEPA (1996a)	2.19	CCME (2007)
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	265	USEPA (1996a)	0.754	USEPA (1996a)	7.20E-02	USEPA (1996a)	200	USEPA (1996a)		
1,2-Dichloroethane	38	USEPA (1996a)	0.0401	USEPA (1996a)	1.04E-01	USEPA (1996a)	8520	USEPA (1996a)		
Dichloromethane (Methylene chloride)	23.74	ORNL (2006)	0.133	ORNL (2006)	1.01E-01	ORNL (2006)	1300	ORNL (2006)		
Trichloromethane (Chloroform)	53.0	USEPA (1996a)	0.15	USEPA (1996a)	1.04E-01	USEPA (1996a)	7920	USEPA (1996a)	0.178	MEMS (2016)
Tetrachloromethane (Carbon tetrachloride)	152	USEPA (1996a)	1.25	USEPA (1996a)	7.80E-02	USEPA (1996a)	793	USEPA (1996a)		
Dibromochloromethane	35	ORNL (2006)	0.032	ORNL (2006)	1.96E-02	ORNL (2006)	2700	ORNL (2006)		
Chlorinated Aromatics										
Chlorobenzene	224	USEPA (1996a)	0.152	USEPA (1996a)	7.30E-02	USEPA (1996a)	472	USEPA (1996a)		
1,2-Dichlorobenzene	379	USEPA (1996a)	0.0779	USEPA (1996a)	6.90E-02	USEPA (1996a)	156	USEPA (1996a)		
1,3-Dichlorobenzene	434	ORNL (2006)	0.108	ORNL (2006)		USEPA (1996a)	125	ORNL (2006)		

TABLE C-6. CHEMICAL PARAMETERS

	Koc	Source	H'	Source	Da	Source	Solubility	Source	Half Life	Source
	ml/g		dimensionless		cm2/s		mg/L		yr	
1,4-Dichlorobenzene	616	USEPA (1996a)	0.0996	USEPA (1996a)	6.90E-02	USEPA (1996a)	73.8	USEPA (1996a)		
1,2,3-Trichlorobenzene			No data, assumed to be equal to other trichlorobenzene isomers							
1,2,4-Trichlorobenzene	1,659	USEPA (1996a)	0.0582	USEPA (1996a)	3.00E-02	USEPA (1996a)	49	ORNL (2006)		
1,3,5-Trichlorobenzene			No data, assumed to be equal to other trichlorobenzene isomers							
1,2,3,4-Tetrachlorobenzene			No data, assumed to be equal to other tetrachlorobenzene isomers							
1,2,3,5-Tetrachlorobenzene			No data, assumed to be equal to other tetrachlorobenzene isomers							
1,2,4,5-Tetrachlorobenzene	1,186	ORNL (2006)	0.0409	ORNL (2006)	2.11E-02	ORNL (2006)	0.595	ORNL (2006)		
Pentachlorobenzene	32,148	USEPA (1996a)	0.0287	ORNL (2006)	5.70E-02	ORNL (2006)	0.831	ORNL (2006)		
Hexachlorobenzene	80,000	USEPA (1996a)	0.0541	USEPA (1996a)	5.42E-02	USEPA (1996a)	6.2	USEPA (1996a)		
Dichlorophenol	718	ORNL (2006)	0.0000895	ORNL (2006)	3.46E-02	ORNL (2006)	4500	ORNL (2006)		
Trichlorophenol	1,186	ORNL (2006)	0.0000662	ORNL (2006)	2.91E-02	ORNL (2006)	1200	ORNL (2006)		
Tetrachlorophenol	2,002	ORNL (2006)	0.000361	ORNL (2006)	2.17E-02	ORNL (2006)	23	ORNL (2006)		
Pentachlorophenol	2,500	ORNL (2006)	1.00E-06	ORNL (2006)	5.60E-02	ORNL (2006)	14	ORNL (2006)		
Dioxins & Furans <sup>2</sup>										
PCBs <sup>2</sup>										
Pesticides										
Aldicarb	32	ORNL (2006)	5.39E-05	ORNL (2006)	3.05E-02	ORNL (2006)	6030	ORNL (2006)		
Aldrin	106,000	ORNL (2006)	0.0018	ORNL (2006)	1.32E-02	ORNL (2006)	0.017	ORNL (2006)		
Atrazine	230	ORNL (2006)	9.65E-08	ORNL (2006)			35	ORNL (2006)		
Azinphos-methyl (Guthion)	231	SRC (2006) <sup>1</sup>	9.96E-07	SRC (2006)			20.9	SRC (2006)		
Bromacil	66.6	ORNL (2006)	5.27E-09	ORNL (2006)			815	ORNL (2006)		
Bromoxynil	435	ORNL (2006)	5.40E-09	ORNL (2006)	2.01E-02	ORNL (2006)	130	ORNL (2006)		
Carbaryl	242	ORNL (2006)	1.78E-07	ORNL (2006)	2.78E-02	ORNL (2006)	110	ORNL (2006)		
Carbofuran	71	ORNL (2006)	1.26E-07	ORNL (2006)			320	ORNL (2006)		
Chlorothalonil	2,392	ORNL (2006)	8.18E-05	ORNL (2006)			0.6	ORNL (2006)		
Chlorpyrifos	6,829	ORNL (2006)	0.00012	ORNL (2006)			1.12	ORNL (2006)		
2,4-D	29	ORNL (2006)	1.45E-06	ORNL (2006)	2.31E-02	ORNL (2006)	677	ORNL (2006)		

TABLE C-6. CHEMICAL PARAMETERS

	Koc	Source	H'	Source	Da	Source	Solubility	Source	Half Life	Source
	ml/g		dimensionless		cm2/s		mg/L		yr	
DDT	794,328	EC (1999a)	3.40E-04	ORNL (2006)	1.37E-02	ORNL (2006)	0.0055	ORNL (2006)		
Diazinon	1,337	ORNL (2006)	4.62E-06	ORNL (2006)	2.06E-02	ORNL (2006)	40	ORNL (2006)		
Dicamba	29	ORNL (2006)	8.91E-08	ORNL (2006)			8310	ORNL (2006)		
Diclofop-methyl	17,092	SRC (2006) <sup>1</sup>	8.21E-05	SRC (2006)			0.8	SRC (2006)		
Dieldrin	10,600	ORNL (2006)	0.000409	ORNL (2006)	1.25E-02	ORNL (2006)	0.25	ORNL (2006)		
Dimethoate	25	ORNL (2006)	4.29E-09	ORNL (2006)			25000	ORNL (2006)		
Dinoseb	3,544	ORNL (2006)	1.86E-05	ORNL (2006)			52	ORNL (2006)		
Diquat	1,933	ORNL (2006)	5.81E-12	ORNL (2006)			708000	ORNL (2006)		
Diuron	136	ORNL (2006)	2.06E-08	ORNL (2006)			42	ORNL (2006)		
Endosulfan	22,000	ORNL (2006)	0.00266	ORNL (2006)	1.15E-02	ORNL (2006)	0.45	ORNL (2006)		
Endrin	10,811	USEPA (1996a)	0.000308	USEPA (1996a)	1.25E-02	USEPA (1996a)	0.25	USEPA (1996a)		
Glyphosate	19	ORNL (2006)	1.67E-17	ORNL (2006)	4.37E-02	ORNL (2006)	12000	ORNL (2006)		
Heptachlor epoxide	9,528	USEPA (1996a)	0.0447	USEPA (1996a)	1.12E-02	USEPA (1996a)	0.18	USEPA (1996a)		
Lindane	1,352	USEPA (1996a)	0.000574	USEPA (1996a)	1.42E-02	USEPA (1996a)	6.8	USEPA (1996a)		
Linuron	350	ORNL (2006)	2.56E-07	ORNL (2006)			75	ORNL (2006)		
Malathion	31	ORNL (2006)	2.00E-07	ORNL (2006)			143	ORNL (2006)		
MCPA	29	ORNL (2006)	5.44E-08	ORNL (2006)			630	ORNL (2006)		
Methoxychlor	80,000	USEPA (1996a)	0.000648	USEPA (1996a)	1.56E-02	USEPA (1996a)	0.045	USEPA (1996a)		
Metolachlor	292	ORNL (2006)	3.68E-07	ORNL (2006)			530	ORNL (2006)		
Metribuzin	1,196	ORNL (2006)	4.78E-09	ORNL (2006)			1050	ORNL (2006)		
Paraquat (as dichloride)	1,405	ORNL (2006)	1.32E-11	ORNL (2006)			620000	ORNL (2006)		
Phorate	444	ORNL (2006)	0.000179	ORNL (2006)			50	ORNL (2006)		
Picloram	18	ORNL (2006)	2.18E-12	ORNL (2006)			430	ORNL (2006)		
Simazine	149	ORNL (2006)	3.85E-08	ORNL (2006)			6.2	ORNL (2006)		
Tebuthiuron	23	ORNL (2006)	4.91E-09	ORNL (2006)			2500	ORNL (2006)		
Terbufos	979	ORNL (2006)	0.000981	ORNL (2006)			5.07	ORNL (2006)		
Toxaphene	95,816	USEPA (1996a)	0.000246	USEPA (1996a)	1.16E-02	USEPA (1996a)	0.74	USEPA (1996a)		

TABLE C-6. CHEMICAL PARAMETERS

	Koc	Source	H'	Source	Da	Source	Solubility	Source	Half Life	Source
	ml/g		dimensionless		cm2/s		mg/L		yr	
Triallate	1,641	ORNL (2006)	0.000789	ORNL (2006)			4	ORNL (2006)		
Trifluralin	9,682	ORNL (2006)	0.00421	ORNL (2006)	1.49E-02	ORNL (2006)	0.184	ORNL (2006)		
Other Organics										
Aniline	45	ORNL (2006)	8.26E-05	ORNL (2006)	7.00E-02	ORNL (2006)	36000	ORNL (2006)		
Di- <i>n</i> -butyl phthalate	1,460	ORNL (2006)	0.000074	ORNL (2006)	4.38E-02	ORNL (2006)	11.2	ORNL (2006)		
Dichlorobenzidine	7,489	ORNL (2006)	2.09E-09	ORNL (2006)	1.94E-02	ORNL (2006)	3.1	ORNL (2006)		
Diethanolamine	***1.9	AENV (2010a)	2.2E-12	AENV (2010a)			miscible	AENV (2010a)	0.75	AENV (2010a)
Diethylene glycol	0.018	AENV (2010b)	5.3E-9	AENV (2010b)			miscible		0.68	AENV (2010b)
Diisopropanolamine	*** 2.2	CCME (2006b)	7.00E-06	CCME (2006b)			870000	CCME (2006b)		
Ethylene glycol	0.0072	EC (1999b)	2.50E-06	EC (1999b)	0.108	ORNL (2006)	miscible	EC (1999b)		
Hexachlorobutadiene	994	ORNL (2006)	0.421	ORNL (2006)	5.61E-02	ORNL (2006)	3.2	ORNL (2006)		
Methanol	0.27	CCME (2017)	0.0002	CCME (2017)	0.15	CCME (2017)	miscible		0.67	CCME (2017)
Methylmethacrylate	10	ORNL (2006)	0.0138	ORNL (2006)	7.70E-02	ORNL (2006)	15000	ORNL (2006)		
Monoethanolamine	***2.21	AENV (2010a)	1.7E-6	AENV (2010a)			miscible	AENV (2010a)	0.75	AENV (2010a)
MTBE	12	USEPA (1994)	0.023	USEPA (1994)	1.02E-01	ORNL (2006)	51000	USEPA (1994)		
Nonylphenol	141,254	EC (2002)	0.005	EC (2002)			5.43	EC (2002)		
Perfluorooctane sulfonate (PFOS)	1,445	CCME (2021)	1.44E-7	CCME (2021)			550	CCME (2021)		
Phenol	12	CCME (1999)	1.60E-05	CCME (1999)	0.082	ORNL (2006)	87,000	CCME (1999)		
Sulfolane	1.2	CCME (2006c)	3.60E-08	CCME (2006c)			miscible	CCME (2006c)		
Triethylene glycol	0.0051	AENV (2010b)	5.3E-9	AENV (2010b)			miscible		0.48	AENV (2010b)

Notes:  
K<sub>oc</sub> = organic carbon water partition coefficient  
H' = dimensionless Henry's Law Coefficient  
D<sub>air</sub> = diffusion coefficient in air  
na = not applicable or not available

\*\*\*Value presented is a mean K<sub>d</sub>, rather than a K<sub>oc</sub> since compound sorbs to clays in preference to organic carbon.

<sup>1</sup>Calculated using the equation K<sub>oc</sub> = 0.41xK<sub>ow</sub>

**TABLE C-6. CHEMICAL PARAMETERS**

<sup>2</sup>PCBs, dioxins and furans are groups of chemicals with a wide range of chemical properties. Chemical properties are not provided for these groups.

CCME (1999) refers to the Canadian Environmental Quality Guidelines (CEQG) and updates, including the scientific supporting documents that are summarized in the CEQG.

**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

	Threshold TRV				Non-Threshold TRV			
	Oral TDI	Source	Inhalation TC	Source	Oral SF	Source	Inhalation UR	Source
	mg/kg-d		mg/m3		(mg/kg-d)-1		(mg/m3)-1	
<b>Metals</b>								
Arsenic (inorganic)					1.8	HC (2021a)		
Barite-barium	0.2	AENV (2009)						
Boron	0.2	AEP (2016)						
Nickel (see note 2)	0.011	CCME (2015)	0.00002	CCME (2015)			0.0000013	CCME (2015)
<b>Hydrocarbons</b>								
Benzene					0.083	HC (2021a)	0.016	HC (2021a)
Toluene	0.0097	HC (2021a)	2.3	HC (2021a)				
Ethylbenzene	0.022	HC (2021a)	2	HC (2021a)				
Xylenes	0.013	HC (2021a)	0.1	HC (2021a)				
Styrene	0.12	HC (1996)	0.85	HC (2017)				
Aliphatic C <sub>6</sub> -C <sub>8</sub>	5.0	CCME (2008a)	18.4	CCME (2008a)				
Aliphatic C <sub>&gt;8</sub> -C <sub>10</sub>	0.1	CCME (2008a)	1.0	CCME (2008a)				
Aliphatic C <sub>&gt;10</sub> -C <sub>12</sub>	0.1	CCME (2008a)	1.0	CCME (2008a)				
Aliphatic C <sub>&gt;12</sub> -C <sub>16</sub>	0.1	CCME (2008a)	1.0	CCME (2008a)				
Aliphatic C <sub>&gt;16</sub> -C <sub>21</sub>	2.0	CCME (2008a)						
Aliphatic C <sub>&gt;21</sub> -C <sub>34</sub>	2.0	CCME (2008a)						
Aliphatic C <sub>&gt;34</sub>	20	CCME (2008a)						
Aromatic C <sub>&gt;8</sub> -C <sub>10</sub>	0.04	CCME (2008a)	0.2	CCME (2008a)				
Aromatic C <sub>&gt;10</sub> -C <sub>12</sub>	0.04	CCME (2008a)	0.2	CCME (2008a)				
Aromatic C <sub>&gt;12</sub> -C <sub>16</sub>	0.04	CCME (2008a)	0.2	CCME (2008a)				
Aromatic C <sub>&gt;16</sub> -C <sub>21</sub>	0.03	CCME (2008a)						
Aromatic C <sub>&gt;21</sub> -C <sub>34</sub>	0.03	CCME (2008a)						
Aromatic C <sub>&gt;34</sub>	0.03	CCME (2008a)						
Naphthalene	0.02	HC (2021a)	0.01	HC (2021a)				

**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

	Threshold TRV				Non-Threshold TRV			
	Oral TDI	Source	Inhalation TC	Source	Oral SF	Source	Inhalation UR	Source
	mg/kg-d		mg/m3		(mg/kg-d)-1		(mg/m3)-1	
Acenaphthene	0.06	US EPA (2006)	0.27	see note 1				
Fluorene	0.04	US EPA (2006)	0.18	see note 1				
Fluoranthene	0.04	US EPA (2006)	0.18	see note 1				
Anthracene	0.3	US EPA (2006)	1.34	see note 1				
Pyrene	0.03	HC (2021a)	0.13	see note 1				
Benzo(a)pyrene	0.0000667	HC (2021a)			1.289	HC (2021a)		
<b>Halogenated Aliphatics</b>								
Vinyl chloride					0.48	HC (2021a)	0.0088	HC (2021a)
1,1-Dichloroethene	0.003	HC (2021a)	0.2	US EPA (2006)				
Trichloroethene (Trichloroethylene, TCE)	0.00146	HC (2021a)	0.002	HC (2021a)	0.000811	HC (2021a)	0.0041	HC (2021a)
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.0047	HC (2021a)	0.04	HC (2021a)				
1,2-Dichloroethane					0.0033	HC (2021a)	0.026	US EPA (2006)
Dichloromethane (Methylene chloride)	0.014	HC (2021a)	0.6	HC (2021a)	0.002	HC (2021a)	0.00001	HC (2021a)
Trichloromethane (Chloroform)	0.0062	HC (2006)	0.3	HC (2017)				
Tetrachloromethane (Carbon tetrachloride)	0.00071	HC (2021a)	0.00318	see note 1				
Dibromochloromethane	0.02	US EPA (2006)	0.08949	see note 1				
<b>Chlorinated Aromatics</b>								
Chlorobenzene	0.43	HC (2021a)	0.01	HC (2021a)				
1,2-Dichlorobenzene	0.43	HC (2021a)	1.92411	see note 1				
1,4-Dichlorobenzene	0.11	HC (2021a)	0.06	HC (2021a)				
1,2,3-Trichlorobenzene	0.0015	HC (1996)	0.00671	see note 1				
1,2,4-Trichlorobenzene	0.0016	HC (1996)	0.007	HC (1996)				
1,3,5-Trichlorobenzene	0.0015	HC (1996)	0.0036	HC (1996)				
1,2,3,4-Tetrachlorobenzene	0.0034	HC 1996)	0.01521	see note 1				



**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

	Threshold TRV				Non-Threshold TRV			
	Oral TDI	Source	Inhalation TC	Source	Oral SF	Source	Inhalation UR	Source
	mg/kg-d		mg/m3		(mg/kg-d)-1		(mg/m3)-1	
1,2,3,5-Tetrachlorobenzene	0.00041	HC (1996)	0.00183	see note 1				
1,2,4,5-Tetrachlorobenzene	0.00021	HC (1996)	0.00094	see note 1				
Pentachlorobenzene	0.001	HC (1996)	0.00447	see note 1				
Hexachlorobenzene	0.0008	US EPA (2006)	0.00358	see note 1	1.6	US EPA (2006)	0.46	US EPA (2006)
2,4-Dichlorophenol	0.1	HC (1987a)	0.447468	see note 1				
2,4,6-Trichlorophenol					0.02	HC (1987a)	0.0045	see note 1
2,3,4,6-Tetrachlorophenol	0.01	HC (1987a)	0.04475	see note 1				
Pentachlorophenol	0.006	HC (1987a)	0.02685	see note 1				
Dioxins and Furans	2.3E-09	HC (2021a)						
PCBs	0.00001	HC (2021a)						
<b>Pesticides</b>								
Aldicarb	0.001	US EPA (2006)						
Aldrin	0.00003	US EPA (2006)			17	US EPA (2006)	4.9	US EPA (2006)
Atrazine and metabolites	0.0005	HC (1993)						
Azinphos-methyl (Guthion)	0.0025	HC (1989a)						
Bromacil	0.1	US EPA (1996b)						
Bromoxynil	0.0005	HC (1989b)						
Carbaryl	0.01	HC (1991a)						
Carbofuran	0.01	HC (1991b)						
Chlorothalonil	0.015	US EPA (2006)						
Chlorpyrifos	0.01	HC (1986a)						
2,4-D	0.01	HC (2009a)						
DDT	0.0005	US EPA (2006)			0.34	US EPA (2006)	0.097	US EPA (2006)
Diazinon	0.002	HC (1986b)						
Dicamba	0.0125	HC (2014)						
Diclofop-methyl	0.001	HC (1987b)						

**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

	Threshold TRV			Non-Threshold TRV				
	Oral TDI	Source	Inhalation TC	Source	Oral SF	Source	Inhalation UR	Source
	mg/kg-d		mg/m3		(mg/kg-d)-1		(mg/m3)-1	
Dieldrin	0.00005	US EPA (2006)			16	US EPA (2006)	4.6	US EPA (2006)
Dimethoate	0.002	HC (2009b)						
Dinoseb	0.001	US EPA (2006)						
Diquat	0.008	HC (1989c)						
Diuron	0.0156	HC (1987c)						
Endosulfan	6.00E-03	US EPA (2006)						
Endrin	3.00E-04	US EPA (2006)						
Glyphosate	0.03	HC (1987d)						
Heptachlor epoxide	1.30E-05	US EPA (2006)			9.1	US EPA (2006)	2.6	US EPA (2006)
Lindane	3.00E-04	US EPA (2006)						
Linuron	2.00E-03	US EPA (2006)						
Malathion	0.02	HC (1986c)						
MCPA	0.012	HC (2010)						
Methoxychlor	0.005	US EPA (2006)						
Metolachlor	0.005	HC (1986d)						
Metribuzin	0.0083	HC (2021b)						
Paraquat (as dichloride)	0.001	HC (1986e)						
Phorate	0.0002	HC (1986f)						
Picloram	0.02	HC (1990)						
Simazine	0.0013	HC (1986g)						
Tebuthiuron	7.00E-02	US EPA (2006)						
Terbufos	0.00005	HC (1995b)						
Toxaphene					1.1	US EPA (2006)	0.32	US EPA (2006)
Triallate	1.30E-02	US EPA (2006)						
Trifluralin	0.0048	HC (1992a)						

**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

	Threshold TRV				Non-Threshold TRV			
	Oral TDI	Source	Inhalation TC	Source	Oral SF	Source	Inhalation UR	Source
	mg/kg-d		mg/m3		(mg/kg-d)-1		(mg/m3)-1	
<b>Other Organics</b>								
Aniline	0.007	HC (1996)	0.001	HC (2017)				
Di- <i>n</i> -butyl phthalate	0.063	HC (1996)	0.28191	see note 1				
Dichlorobenzidine					0.45	US EPA (2006)	0.1	see note 1
Diethanolamine	0.005	AENV (2010a)						
Diethylene glycol	0.5	AENV (2010b)						
Diisopropanolamine	0.39	CCME (2006b)						
Ethylene glycol	2	US EPA (2006)	8.94937	see note 1				
Hexachlorobutadiene					0.078	US EPA (2006)	0.022	US EPA (2006)
Methanol	2.0	CCME (2017)	20	CCME (2017)				
Methylmethacrylate	1.4	US EPA (2006)	0.7	US EPA (2006)				
Monoethanolamine	0.05	AENV (2010a)						
MTBE	0.01	HC (1996)	0.037	HC (1996)				
<b>Nonylphenol</b>								
Perfluorooctane sulfonate (PFOS)	0.00006	HC (2021a)						
Perfluorooctanoic acid (PFOA)	0.000021	CCME (2021a)						
Phenol	0.06	CCME (1997)	0.26848	see note 1				
Sulfolane	0.0097	CCME (2006c)						
Triethylene glycol	5.0	AENV (2010b)						

**Notes:**

TRV = toxicity reference value

TDI = tolerable daily intake

TC = tolerable concentration

SF = slope factor

UR = unit risk

HC = Health Canada

---

**TABLE C-7. HUMAN TOXICITY REFERENCE VALUES**

<sup>1</sup>Estimated from the oral TDI assuming an adult body weight of 70.7 kg, and an inhalation rate of 15.8 m<sup>3</sup>/d.

<sup>2</sup>Human ingestion and dermal contact guideline for agricultural, residential/parkland and commercial land uses based on 10% of estimated daily intake rather than TDI. Human inhalation guidelines calculated separately. See CCME (2015).

TABLE C-8. HUMAN ABSORPTION FACTORS

Absorption Factors						
	Gut	Source	Skin	Source	Lung	Source
Metals						
Arsenic (inorganic)	1.00		0.03	HC (2021a)	1.00	
Barite-barium	1.00		0	AENV (2009)	1.00	
Boron	1.00		0	AEP (2016)	1.00	
Nickel	1.00		0.09	HC (2021a)	1.00	
Hydrocarbons						
Benzene	1.00		0.03	HC (2021a)	1.00	
Toluene	1.00		0.03	HC (2021a)	1.00	
Ethylbenzene	1.00		0.03	HC (2021a)	1.00	
Xylenes	1.00		0.03	HC (2021a)	1.00	
Styrene	1.00		0.03	HC (2021a)	1.00	
Aliphatic C <sub>6</sub> -C <sub>8</sub>	1.00		0.03	HC (2021a)	1.00	
Aliphatic C <sub>&gt;8</sub> -C <sub>10</sub>	1.00		0.03	HC (2021a))	1.00	
Aliphatic C <sub>&gt;10</sub> -C <sub>12</sub>	1.00		0.20	CCME (2008a)	1.00	
Aliphatic C <sub>&gt;12</sub> -C <sub>16</sub>	1.00		0.20	CCME (2008a)	1.00	
Aliphatic C <sub>&gt;16</sub> -C <sub>21</sub>	1.00		0.20	CCME (2008a)	1.00	
Aliphatic C <sub>&gt;21</sub> -C <sub>34</sub>	1.00		0.20	CCME (2008a)	1.00	
Aliphatic C <sub>&gt;34</sub>	1.00		0.20	CCME (2008a)	1.00	
Aromatic C <sub>&gt;8</sub> -C <sub>10</sub>	1.00		0.03	HC (2021a)	1.00	
Aromatic C <sub>&gt;10</sub> -C <sub>12</sub>	1.00		0.20	CCME (2008a)	1.00	
Aromatic C <sub>&gt;12</sub> -C <sub>16</sub>	1.00		0.20	CCME (2008a)	1.00	
Aromatic C <sub>&gt;16</sub> -C <sub>21</sub>	1.00		0.20	CCME (2008a)	1.00	
Aromatic C <sub>&gt;21</sub> -C <sub>34</sub>	1.00		0.20	CCME (2008a)	1.00	
Aromatic C <sub>&gt;34</sub>	1.00		0.20	CCME (2008a)	1.00	
Naphthalene	1.00		0.03	HC (2021a)	1.00	
Acenaphthene	1.00		0.148	HC (2021a)	1.00	
Fluorene	1.00		0.148	HC (2021a)	1.00	
Fluoranthene	1.00		0.148	HC (2021a))	1.00	
Anthracene	1.00		0.148	HC (2021a)	1.00	
Pyrene	1.00		0.148	HC (2021a)	1.00	
Benzo(a)pyrene	1.00		0.148	HC (2021a)	1.00	
Halogenated Aliphatics						
Vinyl chloride	1.00		0.03	HC (2021a)	1.00	
1,1-Dichloroethene	1.00		0.03	HC (2021a)	1.00	
Trichloroethene (Trichloroethylene, TCE)	1.00		0.03	HC (2021a)	1.00	
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	1.00		0.03	HC (2021a)	1.00	
1,2-Dichloroethane	1.00		0.03	HC (2021a)	1.00	
Dichloromethane (Methylene chloride)	1.00		0.03	HC (2021a)	1.00	

TABLE C-8. HUMAN ABSORPTION FACTORS

	Absorption Factors					
	Gut	Source	Skin	Source	Lung	Source
Trichloromethane (Chloroform)	1.00		0.03	HC (2021a)	1.00	
Tetrachloromethane (Carbon tetrachloride)	1.00		0.03	HC (2021a)	1.00	
Dibromochloromethane	1.00		0.03	HC (2021a)	1.00	
Chlorinated Aromatics						
Chlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2-Dichlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,4-Dichlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2,3-Trichlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2,4-Trichlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,3,5-Trichlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2,3,4-Tetrachlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2,3,5-Tetrachlorobenzene	1.00		0.03	HC (2021a)	1.00	
1,2,4,5-Tetrachlorobenzene	1.00		1.00		1.00	
Pentachlorobenzene	1.00		1.00		1.00	
Hexachlorobenzene	1.00		1.00		1.00	
Dichlorophenol	1.00		0.03	HC (2021a)	1.00	
Trichlorophenol	1.00		0.03	HC (2021a)	1.00	
Tetrachlorophenol	1.00		0.03	HC (2021a)	1.00	
Pentachlorophenol	1.00		1.00		1.00	
Dioxins and Furans	1.00		0.03	HC (2021a)	1.00	
PCBs	1.00		0.14	HC (2021a)	1.00	
Pesticides						
Aldicarb	1.00		1.00		1.00	
Aldrin	1.00		1.00		1.00	
Atrazine and metabolites	1.00		1.00		1.00	
Azniphos-methyl (Guthion)	1.00		1.00		1.00	
Bromacil	1.00		1.00		1.00	
Bromoxynil	1.00		1.00		1.00	
Carbaryl	1.00		1.00		1.00	
Carbofuran	1.00		1.00		1.00	
Chlorothalonil	1.00		1.00		1.00	
Chlorpyrifos	1.00		1.00		1.00	
2,4-D	1.00		0.03	HC (2021a)	1.00	
DDT	1.00		1.00		1.00	
Diazinon	1.00		1.00		1.00	
Dicamba	1.00		1.00		1.00	
Diclofop-methyl	1.00		0.03	HC (2021a)	1.00	
Dieldrin	1.00		1.00		1.00	
Dimethoate	1.00		0.03	HC (2021a)	1.00	
Dinoseb	1.00		1.00		1.00	

TABLE C-8. HUMAN ABSORPTION FACTORS

	Absorption Factors					
	Gut	Source	Skin	Source	Lung	Source
Diquat	1.00		1.00		1.00	
Diuron	1.00		1.00		1.00	
Endosulfan	1.00		0.03	HC (2021a)	1.00	
Endrin	1.00		1.00		1.00	
Glyphosate	1.00		1.00		1.00	
Heptachlor epoxide	1.00		1.00		1.00	
Lindane	1.00		1.00		1.00	
Linuron	1.00		1.00		1.00	
Malathion	1.00		1.00		1.00	
MCPA	1.00		1.00		1.00	
Methoxychlor	1.00		1.00		1.00	
Metolachlor	1.00		1.00		1.00	
Metribuzin	1.00		1.00		1.00	
Paraquat (as dichloride)	1.00		1.00		1.00	
Phorate	1.00		1.00		1.00	
Picloram	1.00		1.00		1.00	
Simazine	1.00		1.00		1.00	
Tebuthiuron	1.00		1.00		1.00	
Terbufos	1.00		1.00		1.00	
Toxaphene	1.00		1.00		1.00	
Triallate	1.00		0.03	HC (2021a)	1.00	
Trifluralin	1.00		0.03	HC (2021a)	1.00	
<b>Other Organics</b>						
Aniline	1.00		0.03	HC (2021a)	1.00	
Di- <i>n</i> -butyl phthalate	1.00		1.00		1.00	
Dichlorobenzidine	1.00		1.00		1.00	
Diethanolamine	1.00		1.00		1.00	
Diethylene glycol	1.00		1.00		1.00	
Diisopropanolamine	1.00		0.03	HC (2021a)	1.00	
Ethylene glycol	1.00		0.03	HC (2021a)	1.00	
Hexachlorobutadiene	1.00		0.03	HC (2021a)	1.00	
Methanol	1.00		0.03	HC (2021a)	1.00	
Methylmethacrylate	1.00		0.03	HC (2021a)	1.00	
Monoethanolamine	1.00		1.00		1.00	
MTBE	1.00		0.03	HC (2021a)	1.00	
Nitriloacetic acid	1.00		1.00		1.00	
Nonylphenol	1.00		1.00		1.00	
Perfluorooctane sulfonate (PFOS)	1.00		0.10	CCME (2021)	1.00	
Phenol	1.00		0.03	HC (2021a)	1.00	
Sulfolane	1.00		1.00		1.00	

**TABLE C-8. HUMAN ABSORPTION FACTORS**

	Absorption Factors					
	Gut	Source	Skin	Source	Lung	Source
Triethylene glycol	1.00		1.00		1.00	

**Notes:**

Sources only provided where an absorption factor other than 1.0 is used.

Dermal absorption factors for compounds with a boiling point >250°C are considered volatile for the purposes of dermal absorption. Dermal absorption factors for these compounds are set to 0.03 (HC, 2021a).

CCME (1999) refers to the Canadian Environmental Quality Guidelines (CEQG) and updates, including the scientific supporting documents that are summarized in the CEQG.

HC = Health Canada



**TABLE C-9. HUMAN BACKGROUND EXPOSURE PARAMETERS**

	Toddler EDI	Adult EDI	Source	Ca	Source	BSC	Source	SAF
	mg/kg-d	mg/kg-d		mg/m3		mg/kg		
<b>Metals</b>								
Arsenic (inorganic)	na	na	CCME (1999)	na	CCME (1999)	10	CCME (1999)	0.2
Barite-barium	0.014	0.014	AENV (2009)	na	AENV (2009)	325	AENV (2009)	0.25
Boron	0.048	0.018	AEP (2016)	na	AEP (2016)	10	AEP (2016)	0.25
Nickel	0.0106	0.0038	CCME (2015)	na	-	26.8	CCME (2015)	0.2
<b>Hydrocarbons</b>								
Benzene	na	na	-	na	-	0	assumed	na
Toluene	0.0028	0.0028	EC (2004b)	0.0442	EC (2004b)	0	assumed	0.5
Ethylbenzene	0.0029	0.0029	EC (2004b)	0.0075	EC (2004b)	0	assumed	0.5
Xylenes	0.0079	0.0079	EC (2004b)	0.00182	EC (2004b)	0	assumed	0.5
Styrene	0.00071	0.00027	PSL	0.00028	PSL	0	assumed	0.5
Aliphatic C <sub>6</sub> -C <sub>8</sub>	0.02334	0.02334	CCME (2008a)	0.09111	CCME (2008a)	0	assumed	0.5
Aliphatic C <sub>&gt;8</sub> -C <sub>10</sub>	0.0103	0.0103	CCME (2008a)	0.03881	CCME (2008a)	0	assumed	0.5
Aliphatic C <sub>&gt;10</sub> -C <sub>12</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.5
Aliphatic C <sub>&gt;12</sub> -C <sub>16</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.5
Aliphatic C <sub>&gt;16</sub> -C <sub>21</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.6
Aliphatic C <sub>&gt;21</sub> -C <sub>34</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.6
Aliphatic C <sub>&gt;34</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.8
Aromatic C <sub>&gt;8</sub> -C <sub>10</sub>	0.00938	0.00938	CCME (2008a)	0.03745	CCME (2008a)	0	assumed	0.5
Aromatic C <sub>&gt;10</sub> -C <sub>12</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.5
Aromatic C <sub>&gt;12</sub> -C <sub>16</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.5
Aromatic C <sub>&gt;16</sub> -C <sub>21</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.6
Aromatic C <sub>&gt;21</sub> -C <sub>34</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.6
Aromatic C <sub>&gt;34</sub>	0	0	CCME (2008a)	0	CCME (2008a)	0	assumed	0.8
Naphthalene	0.000535	0.00021231	ATSDR (2005)	0.00095	ATSDR (2005)	0	assumed	0.5
Acenaphthene	0	0	assumed	0	assumed	0	assumed	0.5
Fluorene	0.00902	0.00358	ATSDR (1995)	0.016	ATSDR (1995)	0	assumed	0.5

**TABLE C-9. HUMAN BACKGROUND EXPOSURE PARAMETERS**

	Toddler EDI	Adult EDI	Source	Ca	Source	BSC	Source	SAF
	mg/kg-d	mg/kg-d		mg/m3		mg/kg		
Fluoranthene	0	0	assumed	0	assumed	0	assumed	0.5
Anthracene	0.00502	0.00199	ATSDR (1995)	0.0089	ATSDR (1995)	0	assumed	0.5
Pyrene	0.00620	0.00246	ATSDR (1995)	0.011	ATSDR (1995)	0	assumed	0.5
Benzo(a)pyrene	0.0000059	0.0000059	CCME (2010)	na	-	0.07	CCME (2010)	0.5
<b>Halogenated Aliphatics</b>								
Vinyl chloride	0	0	assumed	0	assumed	0	assumed	0.2
1,1-Dichloroethene	0	0	assumed	0	assumed	0	assumed	0.2
Trichloroethene (Trichloroethylene)	0.00053	0.00041	CCME (2007)	0.0014	CCME (2007)	0	assumed	0.2
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0	0	assumed	0	assumed	0	assumed	0.2
1,2-Dichloroethane	0.0006	0.0005	PSL	0.0018	PSL	0	assumed	0.2
Dichloromethane (Methylene chloride)	0.00558	0.00471	PSL	0.0063	PSL	0	assumed	0.2
Trichloromethane (Chloroform)	0.004315	0.00361	PSL	0.0063	PSL	0	assumed	0.2
Tetrachloromethane (Carbon tetrachloride)	0	0	assumed	0	assumed	0	assumed	0.2
Dibromochloromethane	0	0	assumed	0	assumed	0	assumed	0.2
<b>Chlorinated Aromatics</b>								
Chlorobenzene	0.000122	0.000066	PSL	0.00016	PSL	0	assumed	0.2
1,2-Dichlorobenzene	0.00004	0.00003	PSL	0.1	PSL	0	assumed	0.2
1,4-Dichlorobenzene	0.0014	0.0009	PSL	0.0028	PSL	0	assumed	0.2
1,2,3-Trichlorobenzene	0.00023	0.00024	PSL	0.0008	PSL	0	assumed	0.2
1,2,4-Trichlorobenzene	0.0006	0.00045	PSL	0.0018	PSL	0	assumed	0.2
1,3,5-Trichlorobenzene	0.00032	0.00025	PSL	0.0008	PSL	0	assumed	0.2
1,2,3,4-Tetrachlorobenzene	0.0000007	0.00000025	PSL	0.00000017	PSL	0	assumed	0.2
1,2,3,5-Tetrachlorobenzene	0.00000045	0.00000015	PSL	0.00000017	PSL	0	assumed	0.2
1,2,4,5-Tetrachlorobenzene	0.0000007	0.0000002	PSL	0.00000017	PSL	0	assumed	0.2
Pentachlorobenzene	0.000002	0.0000005	PSL	0.0000001	PSL	0	assumed	0.2
Hexachlorobenzene	0.0000178	0.0000028	PSL	0.00000015	PSL	0	assumed	0.2

**TABLE C-9. HUMAN BACKGROUND EXPOSURE PARAMETERS**

	Toddler EDI	Adult EDI	Source	Ca	Source	BSC	Source	SAF
	mg/kg-d	mg/kg-d		mg/m3		mg/kg		
2,4-Dichlorophenol	0	0	assumed	0	assumed	0	assumed	0.2
2,4,6-Trichlorophenol	0	0	assumed	0	assumed	0	assumed	0.2
2,3,4,6-Tetrachlorophenol	0	0	assumed	0	assumed	0	assumed	0.2
Pentachlorophenol	0	0	assumed	0	assumed	0	assumed	0.2
Dioxins and Furans	7.1E-09	1.33E-09	CCME (2002)	0	assumed	0	assumed	0.25
PCBs	7.69E-06	0.00000254	EC (2001)	0	assumed	0	assumed	0.2
<b>Pesticides</b>								
Aldicarb	0	0	assumed	0	assumed	0	assumed	0.2
Aldrin	0	0	assumed	0	assumed	0	assumed	0.2
Atrazine and metabolites	0	0	assumed	0	assumed	0	assumed	0.2
Azinphos-methyl (Guthion)	0	0	assumed	0	assumed	0	assumed	0.2
Bromacil	0	0	assumed	0	assumed	0	assumed	0.2
Bromoxynil	0	0	assumed	0	assumed	0	assumed	0.2
Carbaryl	0	0	assumed	0	assumed	0	assumed	0.2
Carbofuran	0	0	assumed	0	assumed	0	assumed	0.2
Chlorothalonil	0	0	assumed	0	assumed	0	assumed	0.2
Chlorpyrifos	0	0	assumed	0	assumed	0	assumed	0.2
2,4-D	0	0	assumed	0	assumed	0	assumed	0.2
DDT	0	0	assumed	0	assumed	0	assumed	0.2
Diazinon	0	0	assumed	0	assumed	0	assumed	0.2
Dicamba	0	0	assumed	0	assumed	0	assumed	0.2
Diclofop-methyl	0	0	assumed	0	assumed	0	assumed	0.2
Dieldrin								
Dimethoate	0	0	assumed	0	assumed	0	assumed	0.2
Dinoseb	0	0	assumed	0	assumed	0	assumed	0.2
Diquat	0	0	assumed	0	assumed	0	assumed	0.2
Diuron	0	0	assumed	0	assumed	0	assumed	0.2

**TABLE C-9. HUMAN BACKGROUND EXPOSURE PARAMETERS**

	Toddler EDI	Adult EDI	Source	Ca	Source	BSC	Source	SAF
	mg/kg-d	mg/kg-d		mg/m3		mg/kg		
Endosulfan	0	0	assumed	0	assumed	0	assumed	0.2
Endrin	0	0	assumed	0	assumed	0	assumed	0.2
Glyphosate	0	0	assumed	0	assumed	0	assumed	0.2
Heptachlor epoxide	0	0	assumed	0	assumed	0	assumed	0.2
Lindane	0	0	assumed	0	assumed	0	assumed	0.2
Linuron	0	0	assumed	0	assumed	0	assumed	0.2
Malathion	0	0	assumed	0	assumed	0	assumed	0.2
MCPA	0	0	assumed	0	assumed	0	assumed	0.2
Methoxychlor	0	0	assumed	0	assumed	0	assumed	0.2
Metolachlor	0	0	assumed	0	assumed	0	assumed	0.2
Metribuzin	0	0	assumed	0	assumed	0	assumed	0.2
Paraquat (as dichloride)	0	0	assumed	0	assumed	0	assumed	0.2
Phorate	0	0	assumed	0	assumed	0	assumed	0.2
Picloram	0	0	assumed	0	assumed	0	assumed	0.2
Simazine	0	0	assumed	0	assumed	0	assumed	0.2
Tebuthiuron	0	0	assumed	0	assumed	0	assumed	0.2
Terbufos	0	0	assumed	0	assumed	0	assumed	0.2
Toxaphene	0	0	assumed	0	assumed	0	assumed	0.2
Triallate	0	0	assumed	0	assumed	0	assumed	0.2
Trifluralin	0	0	assumed	0	assumed	0	assumed	0.2
<b>Other Organics</b>								
Aniline	0	0	assumed	0	assumed	0	assumed	0.2
Di- <i>n</i> -butyl phthalate	0.005	0.0019	PSL	0.00285	PSL	0	assumed	0.2
Dichlorobenzidine	0	0	assumed	0	assumed	0	assumed	0.2
Diethanolamine	0	0	assumed	0	assumed	0	assumed	0.25
Diethylene glycol	0	0	assumed	0	assumed	0	assumed	0.25
Diisopropanolamine	0	0	assumed	0	assumed	0	assumed	0.33

**TABLE C-9. HUMAN BACKGROUND EXPOSURE PARAMETERS**

	Toddler EDI	Adult EDI	Source	Ca	Source	BSC	Source	SAF
	mg/kg-d	mg/kg-d		mg/m3		mg/kg		
Ethylene glycol	0.0344	0.0167	PSL	0	PSL	0	assumed	0.33
Hexachlorobutadiene	0.00012	0.00003	PSL	0.00006	PSL	0	assumed	0.2
Methanol	1.6	1.6	CCME (2017)	0.04	CCME (2017)	0	Assumed	0.2
Methylmethacrylate	1.13E-07	0	PSL	2.44E-07	PSL	0	assumed	0.2
Monoethanolamine	0	0	assumed	0	assumed	0	assumed	0.25
MTBE	0.00000067	0.0000005	PSL	0.0000015	PSL	0	assumed	0.2
Nonylphenol	na	na	-	na	-	0	assumed	na
Perfluorooctane sulfonate (PFOS)	0.0000038	0.0000023	CCME (2021)	na	-	0	assumed	0.2
Phenol	0	0	assumed	0	assumed	0	assumed	0.2
Sulfolane	0	0	assumed	0	assumed	0	assumed	0.33
Triethylene glycol	0	0	assumed	0	assumed	0	assumed	0.25

**Notes:**

na = not available or not applicable

EDI = estimated daily intake

C<sub>a</sub> = background indoor air concentration

SAF = soil allocation factor

PSL = Priority Substance List assessment under Canadian Environmental Protection Act (CEPA) for corresponding substance.

CCME (1999) refers to the Canadian Environmental Quality Guidelines (CEQG) and updates, including the scientific supporting documents that are summarized in the CEQG.

**TABLE C-10. PETROLEUM HYDROCARBON SUBFRACTION DISTRIBUTION**

Soil				
TPH Sub-fraction (Proportion of Total Fraction Mass)				
Fraction	Fraction 1	Fraction 2	Fraction 3	Fraction 4
<b>Aliphatics</b>				
C6-C8	0.55			
C>8-C10	0.36			
C>10-C12		0.36		
C>12-C16		0.44		
C>16-C21			0.56	
C>21-C34			0.24	
C>34				0.8
<b>Aromatics</b>				
C>7-C8				
C>8-C10	0.09			
C>10-C12		0.09		
C>12-C16		0.11		
C>16-C21			0.14	
C>21-C34			0.06	
C>34				0.2
Sum of all subfractions	1	1	1	1
<b>Groundwater</b>				
TPH Sub-fraction (Proportion of Total Fraction Mass)				
Fine Soil		Coarse Soil		
Fraction	Fraction 1	Fraction 2	Fraction 1	Fraction 2
<b>Aliphatics</b>				
C6-C8	0.5768		0.6047	
C>8-C10	0.0663		0.0632	
C>10-C12		0.0239		0.024
C>12-C16		0.0015		0.0015
<b>Aromatics</b>				
C>7-C8				
C>8-C10	0.3569		0.3321	
C>10-C12		0.6029		0.6034
C>12-C16		0.3718		0.3711
Sum of all subfractions	1	1	1	1

Notes:

Source: CCME (2008a)

---

**TABLE C-10. PETROLEUM HYDROCARBON SUBFRACTION DISTRIBUTION**

Subfraction distribution in groundwater not required for F3 and F4 due to low aqueous solubility  
Groundwater values were calculated from the soil subfraction distributions above based on equilibrium partitioning assumptions, see text.

**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
<b>General and Inorganic Parameters</b>						
Aluminum	2.9	see note 2	5	5		
Ammonia		see note 2				
Bromate	0.01					
Chloride	250	120	100			
Cyanide (free)	0.2	0.0052				
Electrical Conductivity (dS/m)			1			
Fluoride	1.5		1	1		
Nitrate (as nitrogen)	10	3				
Nitrate + Nitrite (as nitrogen)				100		
Nitrite (as nitrogen)	1.0	see note 2		10		
Sodium	200					
Sodium Adsorption Ratio (SAR)			5			
Sulphate	500	see note 2		1000		
Sulphide - Total (as S) <sup>3</sup>	0.05	0.002				
Total Dissolved Solids (TDS)	500		500	3000		
<b>Metals</b>						
Antimony	0.006					
Arsenic (inorganic)	0.01	0.005	0.16	0.025		
Barium	2.0					
Boron	5	1.5	1.0	5		
Cadmium	0.007	see note 2	0.0082	0.08		
Chromium (trivalent)		0.0089	0.0049	0.05		
Chromium (hexavalent)		0.001	0.008	0.05		
Chromium (total) <sup>8</sup>	0.05					
Copper	1	0.007	0.2	0.5		
Iron	0.3	0.3	5			
Lead	0.005	see note 2	0.2	0.1		
Manganese	0.02		0.2			
Mercury (total) <sup>8</sup>	0.001	0.000005		0.003		



**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
Nickel		see note 2	0.2	1		
Selenium	0.05	0.002	0.02	0.05		
Silver		0.0001	0.02	0.05		
Uranium	0.02	0.015	0.01	0.2		
Zinc	5	0.03	1	50		
<b>Hydrocarbons</b>						
Benzene	0.005	0.04		0.088	0.076	0.08
Toluene	0.024	0.0005		4.91	4.25	4.46
Ethylbenzene	0.0016	0.09		3.20	2.77	2.91
Xylenes	0.02	0.03		13.1	11.3	11.9
Styrene	2.8	0.072				
Aliphatic C <sub>6</sub> -C <sub>8</sub>	140	0.047		53.6	46.4	48.72
Aliphatic C <sub>&gt;8</sub> -C <sub>10</sub>	2.5	0.0076		53.6	46.4	48.72
Aromatic C <sub>&gt;8</sub> -C <sub>10</sub>	0.84	0.14		53.6	46.4	48.72
F1		see note 4		53.6	46.4	48.72
Aliphatic C <sub>&gt;10</sub> -C <sub>12</sub>	2.75	0.0012		49.2	42.6	44.73
Aliphatic C <sub>&gt;12</sub> -C <sub>16</sub>	2.75	0.000074		49.2	42.6	44.73
Aromatic C <sub>&gt;10</sub> -C <sub>12</sub>	1.1	0.096		49.2	42.6	44.73
Aromatic C <sub>&gt;12</sub> -C <sub>16</sub>	1.1	0.055		49.2	42.6	44.73
F2		see note 4		49.2	42.6	44.73
F3				79.7	69.0	72.45
F4				42.0	36.4	38.22
Acenaphthene	1.414	0.0058		NGR	NGR	70
Anthracene	7.07	0.000012		NGR	NGR	200
Fluoranthene	0.94	0.00004		NGR	NGR	50
Fluorene	0.94	0.003		NGR	NGR	50
Naphthalene	0.47	0.001		NGR	NGR	28.6
Phenanthrene		0.0004		NGR	NGR	140
Pyrene	0.71	0.000025		NGR	NGR	25
Benz[a]anthracene		0.000018		NGR	NGR	20

**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
Benzo[b+j]fluoranthene				NGR	NGR	20
Benzo[k]fluoranthene				NGR	NGR	20
Benzo[g,h,i]perylene						
Benzo[a]pyrene	0.00004	0.000015		NGR	NGR	2
Chrysene				NGR	NGR	20
Dibenz[a,h]anthracene						
Indeno[1,2,3-c,d]pyrene						
<b>Halogenated Aliphatics</b>						
Vinyl chloride	0.002					
1,1-Dichloroethene	0.014					
Trichloroethene (Trichloroethylene, TCE)	0.005	0.021		0.05		
Tetrachloroethene (Tetrachloroethylene, Perchloroethylene, PCE)	0.010	0.111				
1,2-Dichloroethane	0.005	0.1		0.005		
Dichloromethane (Methylene chloride)	0.05	0.098		0.05		
Trichloromethane (Chloroform) <sup>7</sup>	0.08	0.0018		0.1		
Tetrachloromethane (Carbon tetrachloride)	0.002	0.0133		0.005		
Dibromochloromethane	0.189			0.1		
<b>Chlorinated Aromatics</b>						
Chlorobenzene	0.03	0.0013				
1,2-Dichlorobenzene	0.003	0.0007				
1,3-Dichlorobenzene		0.15				
1,4-Dichlorobenzene	0.001	0.026				
1,2,3-Trichlorobenzene	0.014	0.008				
1,2,4-Trichlorobenzene	0.015	0.024				
1,3,5-Trichlorobenzene	0.014					
1,2,3,4-Tetrachlorobenzene	0.032	0.0018				
1,2,3,5-Tetrachlorobenzene	0.0038					
1,2,4,5-Tetrachlorobenzene	0.0020					
Pentachlorobenzene	0.0094	0.006				

**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
Hexachlorobenzene	0.000568			0.00052		
2,4-Dichlorophenol	0.0003	0.0002				
2,4,6-Trichlorophenol	0.002	0.018				
2,3,4,6-Tetrachlorophenol	0.001	0.001				
Pentachlorophenol	0.03	0.0005				
Dioxins and Furans	2.7E-08					
PCBs	0.000094					
<b>Pesticides</b>						
Aldicarb	0.009	0.001	0.073	0.011		
Aldrin	0.000028					
Atrazine and metabolites	0.005	0.0018	0.01	0.005		
Azinphos-methyl (Guthion)	0.02	0.00001				
Bromacil <sup>5</sup>	0.95	0.005	0.0002	1.1		
Bromoxynil	0.005	0.005	0.00044	0.011		
Carbaryl	0.09	0.0002		1.1		
Carbofuran	0.09	0.0018		0.045		
Chlorothalonil	0.14	0.00018	0.0093	0.17		
Chlorpyrifos	0.09	0.000002		0.024		
2,4-D	0.1	0.004		0.1		
DDT	0.0014					
Diazinon	0.02	0.00017				
Dicamba	0.12	0.01	0.000008	0.122		
Diclofop-methyl	0.009	0.0061	0.00024	0.009		
Dieldrin	0.000029					
Dimethoate	0.02	0.0062		0.003		
Dinoseb	0.01	0.00005	0.021	0.15		
Diquat	0.07					
Diuron	0.15					
Endosulfan	0.057	0.000003				
Endrin	0.0028					

**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
Glyphosate	0.28	0.065		0.28		
Heptachlor epoxide	0.000052					
Lindane	0.0028	0.00001		0.004		
Linuron	0.019	0.007	0.00011			
Malathion	0.19	0.0001				
MCPA	0.1	0.0026	0.00004	0.025		
Methoxychlor	0.9	0.00003				
Metolachlor	0.05	0.0078	0.028	0.05		
Metribuzin	0.08	0.001	0.0005	0.08		
Paraquat (as dichloride)	0.01					
Phorate	0.002					
Picloram	0.19	0.029		0.19		
Simazine	0.01	0.01	0.0005	0.01		
Tebuthiuron	0.66	0.0016	0.00043	0.13		
Terbufos	0.001					
Toxaphene	0.00043					
Triallate	0.123	0.00024		0.23		
Trifluralin	0.045	0.0002		0.045		
<b>Other Organics</b>						
Aniline	0.066	0.0022				
Di- <i>n</i> -butyl phthalate	0.59	0.019				
Dichlorobenzidine	0.001					
Diethanolamine	0.06	0.45				
Diethylene glycol	6.0	150				
Diisopropanolamine	3.6	1.6	3.2			
Ethylene glycol	31	190				
Hexachlorobutadiene	0.006	0.0013				
Methanol <sup>6</sup>	19	23				
Methylmethacrylate	13					
Monoethanolamine	0.6	0.075				

**TABLE C-11. SURFACE WATER QUALITY GUIDELINES**

	Drinking Water	Aquatic Life	Irrigation	Livestock Water	Wildlife Water	DTED <sup>1</sup>
	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/L)	(mg/kg-bw/d)
MTBE	0.015	10				
Nitriloacetic acid	0.4					
Nonylphenol + ethoxylates		0.0066				
Perfluorooctane sulfonate (PFOS)	0.0006	0.0068		0.060	0.052	0.0543
Perfluorooctanoic acid (PFOA)	0.0002					
Phenol	0.57	0.004		0.002		
Sulfolane	0.09	50	0.8			
Triethylene glycol	60	350				
Trihalomethanes - total (THMs)	0.1					

Notes:

See text for guideline sources.

<sup>1</sup> DTED = daily threshold effect dose from CCME (1999, 2008a, 2021). Included where used to calculate livestock and wildlife watering guidelines according to:

$$WQG = \frac{DTED \times BW \times AF}{WIR}$$

Where:

WQG = water quality guideline (mg/L)  
 DTED = daily threshold effect dose (mg/kg-bw/d)  
 BW = body weight (kg)  
 AF = allocation factor of 0.2 (unitless)  
 WIR = water ingestion rate (L/d)

<sup>2</sup> See Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014).

<sup>3</sup> Surface water guidelines based on H<sub>2</sub>S toxicity but guidelines may be applied to total sulphide measurements.

<sup>4</sup> Aquatic life guidelines for direct application to surface water are found in Environmental Quality Guidelines for Alberta Surface Waters (ESRD, 2014). F1 and F2 subfraction guidelines are used to calculate soil and groundwater guidelines for the protection of aquatic life.

<sup>5</sup> Drinking water guideline calculated from Reference Dose (USEPA 1996b).

<sup>6</sup> Source guidance values for drinking water and aquatic life from CCME (2017).

<sup>7</sup> Source guidance value from health based target presented in HC (2006).

<sup>8</sup> Total means all chemical species.

NGR - no guideline required, calculated value > solubility