

**Background**

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

**Fall Index Netting (FIN)**

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

**How is this information used?**

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is **long-term sustainability**, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information please see Alberta’s FIN and FSI websites,

- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fall-index-netting/default.aspx>
- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fish-sustainability-index/default.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Pike using the standardized Fall Index Net (FIN) method. **Note:** Thresholds align with species management frameworks.

Mature Walleyes / net	Mature Pike / net	Risk to Sustainability
>29.0	>21.8	Very Low
20.3-29.0	15.3-21.8	Low
14.5-20.2	10.9-15.2	Moderate
5.8-14.4	4.4-10.8	High
<5.8	<4.4	Very High

**Results of the 2014 FIN at Gods Lake**

Gods Lake (698 ha) is located 173 km north from the town of Fort McMurray. From September 22-25, 2014, nine gill nets captured 231 Cisco, 27 Lake Whitefish, 49 Northern Pike, 34 Walleye, and 77 Yellow Perch, from Gods Lake.

**Walleye**

The mean catch rate of Walleyes was 3.8/net-night. The catch rates of mature (Figure 1) and immature Walleye were 2.8/net-night and 1.0/net-night, respectively. The corresponding FSI score for the current mature density of Walleye was assessed at **very high risk**.

The length distribution indicates moderate but unstable recruitment, low abundance of 350 to 540 mm Walleyes, and abundant fish larger than 550 mm (Figure 2). The population seems to be supported by a few large size-classes.

The 2014 FIN sample represented approximately 0.5 % of the estimated Walleye population size.




Figure 1 - The FIN catch rate of mature Walleyes from Gods Lake, 2014. Dashed line is the mean catch rate (2.8 fish/net-night), with individual net data as hollow circles (n=9 nets).




Figure 3 - The FIN catch rate of mature Northern Pike from Gods Lake, 2014. Dashed line is the mean catch rate (5.4 fish/net-night), with individual net data as hollow circles (n=9 nets).




Figure 2 – FIN sample of showing size of Walleyes from Gods Lake, 2014. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.




Figure 4 – FIN sample showing size of Northern Pike from Gods Lake, 2014. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

### Northern Pike

The mean catch rate of mature Northern pike was 6.0/net-night (Figure 3). The corresponding FSI score for the mature density of Northern Pike was assessed at **high risk**.

The length distribution indicates very poor recruitment, a very low abundance of 500 to 690 mm Pike, and moderate abundance of fish larger than 730 mm (Figure 4). The population seems to be supported by a few large size-classes (Figure 4).

The 2014 FIN sample represented approximately 0.2% of the estimated Northern Pike population size.

### Summary

Since the FIN assessment in 2008, the corresponding FSI

status of the Walleye population in Gods Lake has decreased from **moderate risk** to **high risk**. Dependant on the management objectives, conservation-based management is required to sustain this Walleye population.

The FSI status of Northern Pike has remained at **very high risk** and strict conservation efforts are therefore necessary, dependant on the management objective.

### Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.