

**Background**

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

**Fall Index Netting (FIN)**

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

**How is this information used?**

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The Fisheries Management Objective for most Alberta fisheries is **long-term sustainability**, shown by the red lines

on the graphs below. Achieving this objective uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information please see Alberta’s FIN and FSI websites,

- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fall-index-netting/default.aspx>
- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fish-sustainability-index/default.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Pike using the standardized Fall Index Net (FIN) method. **Note:** Thresholds align with species management frameworks.

Mature Walleyes / net	Mature Pike / net	Risk to Sustainability
>29.0	>21.8	Very Low
20.3-29.0	15.3-21.8	Low
14.5-20.2	10.9-15.2	Moderate
5.8-14.4	4.4-10.8	High
<5.8	<4.4	Very High

**Results of the 2016 FIN at Fawcett Lake**

Fawcett Lake (3445 ha) is located approximately 240 km north of the city of Edmonton. From September 10-13, 2016, thirteen gill nets captured 112 Cisco, 2 Lake Whitefish, 1 Cisco/Lake Whitefish hybrid, 107 Northern Pike, 6 Spottail Shiners, 266 Walleye, 27 White Suckers, and 186 Yellow Perch, from Fawcett Lake.

**Walleye**

The mean catch rate of Walleyes was 20.5/net-night. The catch rates of mature (Figure 1) and immature Walleye were 12.5/net-night and 7.8/net-night, respectively. The corresponding FSI score for the current mature density of Walleye was assessed at **high risk**.

The length distribution shows stable and somewhat variable recruitment, with moderately abundant Walleye in the 370-500 mm size classes, and above average abundance of fish larger than 500 mm (Figure 2).

The 2016 FIN sample represented approximately 0.7% of the estimated Walleye population size.




Figure 1 - The FIN catch rate of mature Walleyes from Fawcett Lake, 2016. Dashed line is the mean likelihood catch rate (12.5/fish/net-night), with net individual data as hollow circles (n=13 nets).




Figure 3 - The FIN catch rate of Northern Pike from Fawcett Lake, 2016. Dashed line is the mean likelihood catch rate (8.1 fish/net-night), with individual net data as hollow circles (n=13 nets).




Figure 2 – FIN sample of showing size of Walleyes from Fawcett Lake, 2016. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.




Figure 4 – FIN sample showing size of Northern Pike from Fawcett Lake, 2016. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

### Northern Pike

The mean catch rate of mature Northern Pike was 8.1/net-night (Figure 3). The corresponding FSI score for the mature density of Northern Pike was assessed at **high risk**.

The length distribution shows intermittent and unstable recruitment, moderately abundant 480 to 670 mm pike, and very few fish larger than 670 mm (Figure 4).

The 2016 FIN sample represented approximately 0.2% of the estimated Northern Pike population size.

### Summary

From the 2006, 2011 and 2013 FINs, the abundance of walleye has shown incremental improvement however the FSI status of Walleye has remained at **high risk**. The Fawcett Lake Walleye population has improved in stability over 10 years, albeit at low densities.

Since the 2006 FIN, the density of mature Northern Pike has remained at a FSI status of **high risk**. The length distribution indicates a broad size distribution with low and inconsistent recruitment, and moderate to low abundance of medium and large sized pike. Conservation-based management should remain the focus to improve the abundance of pike in order to provide sustainable harvest or potentially quality harvest opportunities.

To ensure the long-term sustainability of both the Walleye and Northern Pike populations and fisheries, conservation-based management remains necessary.

### Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.