

# Lac La Nonne FIN Summary

## 2018

### Background

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

### Fall Index Netting (FIN)

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

### How is this information used?

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is long-term sustainability, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status.

For more information, please see Alberta’s FIN and FSI websites,

- <https://www.alberta.ca/fall-index-netting.aspx>
- <https://www.alberta.ca/fish-sustainability-index.aspx>

**Table 1** – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Northern Pike using the standardized Fall Index Net (FIN) method. Note: Thresholds align with species management frameworks.

| Mature Walleyes/net | Mature Pike/net | Risk to Sustainability |
|---------------------|-----------------|------------------------|
| >29.0               | >21.8           | Very Low               |
| 20.3-29.0           | 15.3-21.8       | Low                    |
| 14.5-20.2           | 10.9-15.2       | Moderate               |
| 5.8-14.4            | 4.4-10.8        | High                   |
| <5.8                | <4.4            | Very High              |

### Results of the 2018 FIN at Lac La Nonne


Lac La Nonne (1292 ha) is located 70 km northwest from the city of Edmonton. From September 24 to 26, 2018, 12 nets captured 9 Lake Whitefish, 151 Northern Pike, 93 Walleyes and 151 Yellow Perch.

### Walleye


The mean catch rate of Walleyes was 7.8/ net-night. The catch rates of mature (Figure 1) and immature Walleyes were 3.1/ net-night and 4.7/ net-night, respectively. The corresponding FSI score for the current mature density of Walleyes was assessed at **very high risk**.

The length distribution indicates unstable yet abundant recruitment, abundant 300 to 390 mm Walleye but very few fish larger than 390 mm (Figure 2).

The 2018 FIN sample represented approximately 1.7% of the estimated mature Walleye population size.



**Figure 1** - The FIN catch rate of mature Walleyes from Lac La Nonne, 2018. Dashed line is the mean catch rate (3.1 fish/ net-night), with individual net data as hollow circles (n=12 nets).



**Figure 2** – FIN sample of showing size of Walleyes from Lac La Nonne, 2018. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.

## Northern Pike


The mean catch rate of mature Northern Pike was 12.6/ net-night (Figure 3). The corresponding FSI score for the current mature density of Northern Pike was assessed at **moderate risk**.

The length distribution indicates weak recruitment, very abundant Northern Pike from 490 to 680 mm and very few fish larger than 680 mm (Figure 4).


The 2018 FIN sample represented approximately 0.4% of the estimated mature Northern Pike population size.

## Summary

In several FIN assessments before 2016 indicated the status of Walleye in Lac La Nonne varied between **moderate** to **low** risk. A severe winterkill in 2015 resulted in a 2016 status of **very high risk**, which continues today. Strict conservation-based management actions are necessary to achieve recovery.



**Figure 3** - The FIN catch rate of mature Northern Pike from Lac La Nonne, 2018. Dashed line is the mean catch rate (12.6 fish/ net-night), with individual net data as hollow circles (n=12 nets).



**Figure 4** – FIN sample showing size of Northern Pike from Lac La Nonne, 2018. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

In the 2008 to 2016 assessments, the corresponding FSI status of Northern Pike remained at **high risk**. The 2015 winter kill reduced the abundance of pike slightly. Given the current status of **moderate risk**, and with careful management, the long-term goal of sustainable harvest can be met achieved.

## Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.