

Background

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

Fall Index Netting (FIN)

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. At Garner Lake in 2013, a half-length variation of the standard index net was used, balancing precision of the catch rates with reduced sampling effort. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

How is this information used?

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is **long-term sustainability**, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information please see Alberta’s FIN and FSI websites,

- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fall-index-netting/default.aspx>
- <http://aep.alberta.ca/fish-wildlife/fisheries-management/fish-sustainability-index/default.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Pike using the standardized Fall Index Net (FIN) method. **Note:** Thresholds align with species management frameworks.

Mature Walleyes / 1/2 net	Mature Pike / 1/2 net	Risk to Sustainability
>14.5	>10.9	Very Low
10.2-14.5	7.7-10.9	Low
7.3-10.1	5.5-7.6	Moderate
2.9-7.2	2.2-5.4	High
<2.9	<2.2	Very High

Results of the 2013 FIN at Garner Lake

Garner Lake (776 ha) is located approximately 170 km northeast of the city of Edmonton. From September 30 to October 2, 2013, eight ½ length nets captured three Northern Pike, 129 Walleye, and 65 Yellow Perch.

Walleye

The mean catch rate of Walleyes was 16.1/ ½ net-night. The catch rates of mature (Figure 1) and immature Walleye were 15.9/ ½ net-night and 0.3/ ½ net-night, respectively. The corresponding FSI score for the current mature density of Walleye was assessed at **very low risk**.

The length distribution shows no recruitment of Walleye to the population fishery, and a high abundance of Walleye larger than 440 mm (Figure 2). Garner Lake was last stocked in 1991 and it is likely no Walleyes are surviving from that stocking.

The 2013 FIN sample represented approximately 1% of the estimated mature Walleye population size.




Figure 1 - The FIN catch rate of mature Walleyes from Garner Lake, 2013. Dashed line is the mean catch rate (15.9 fish/ ½ net-night), with net data as hollow circles (n=8 nets).




Figure 3 - The FIN catch rate of Northern Pike from Garner Lake, 2013. Dashed line is the mean catch rate (0.3 fish/ ½ net-night), with net data as hollow circles (n=8 nets).




Figure 2 – FIN sample of showing size of Walleyes from Garner Lake, 2013. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.




Figure 4 – FIN sample showing size of Northern Pike from Garner Lake, 2013. The red line indicates the average length distribution of Pike from 6 Alberta lakes supporting long-term sustainable harvests of Pike.

Northern Pike

The mean catch rate of mature Northern Pike was 0.3/ ½ net-night (Figure 3). The corresponding FSI score for the current mature density of Northern Pike was assessed at **very high risk**.

The length distribution shows an overexploited population with no recruitment and three Northern Pike from 530 to 560 mm (Figure 4).

The 2013 FIN sample represented approximately 0.3% of the estimated mature Northern Pike population size.

Summary

Since the FINs on Garner Lake in 2005, 2010, and 2013, the density of mature Walleye has been steadily classified at a corresponding FSI status of **very low risk**. The FSI status is

from exceptionally strong size-classes; however, the unusual pulse of Walleyes at Garner Lake may provide angling opportunities, dependant on the management objective.

Since the FIN in 2005, the corresponding FSI status of mature Northern Pike has declined from **moderate risk** to **high risk** in 2010, and then in 2013 to **very high risk**. Dependant on the management objective, stringent conservation-focused management is necessary to recover this Pike stock.

Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.