

Big Island Lake FIN Summary 2020

Background

“How are the fish in my lake doing?” We need this answer to set appropriate fishing regulations, to understand and correct any problems with fish habitat, and to guard against invasive species. A healthy fish population and fish community means we can all enjoy the benefits of sustainable fisheries and healthy ecosystems. A standard method of assessing the status of fish populations is necessary to allow comparisons of fish sustainability across the years at a lake, and to compare to other lakes. In Alberta, we use an accepted standard of index netting for lake fisheries assessment. This method provides the necessary data on fish abundance, biological data (such as age and sex), and species diversity to assess sustainability.

Fall Index Netting (FIN)

Alberta Environment and Parks monitor Walleye and Northern Pike populations using standardized index netting (Morgan, 2002). Fall index netting occurs during late summer and fall when water temperatures are 10-15 °C. Standardized multi-mesh gill nets are set at random locations between 2 and 15 metres deep, set for 21-27 hours (i.e., a net-night), and then reset in new random locations. At Big Island Lake in 2020, a half-length variation of the standard index net was used, balancing precision of the catch rates with reduced sampling effort. Information from Yellow Perch, Lake Whitefish, Burbot, minnow, and sucker species are also collected. The information collected from each fish includes length, weight, age, gender, and maturity. After sampling, if fish are appropriate for human consumption, Alberta biologists provide the fish to local Indigenous peoples or to persons on approved subsistence lists. Typically, a tiny proportion of the lake’s fish population (usually less than 1 or 2%) are killed in this sampling.

How is this information used?

Catch rates (i.e., number of fish captured per net-night) of Walleye and Northern Pike are an index of the populations’ abundance, with higher catch rates meaning there are more fish in the lake. The abundance of adult fish is compared to the standardized thresholds for 5 broad categories of risk to the long-term sustainability of the fish population, with higher densities of fish having lower risk (Table 1). The sizes and age of fish also tell us if problems with overharvest (e.g. too few fish living to old age) or

habitat (e.g., poor spawning success) are a concern. Biologists use this information, as well as a variety of data on water quality, access, development, and habitat threats as part of Alberta’s Fish Sustainability Index (FSI).

The management goal for most Alberta fisheries is long-term sustainability, shown by the red lines on the graphs below. Achieving this goal uses the netting data and the FSI to determine the most appropriate sport fishing regulations for a lake. This landscape-level assessment allows for consistent, broad temporal comparisons of fish sustainability and status. For more information, please see Alberta’s FIN and FSI websites,

- <https://www.alberta.ca/fall-index-netting.aspx>
- <https://www.alberta.ca/fish-sustainability-index-overview.aspx>

Table 1 – Alberta’s Fish Sustainability Index risk thresholds for Walleye and Northern Pike using the standardized Fall Index Net (FIN) method. Note: Thresholds align with species management frameworks.

Mature Walleyes/ ½ net	Mature Pike/ ½ net	Risk to Sustainability
>14.5	>10.9	Very Low
10.2-14.5	7.7-10.9	Low
7.3-10.1	5.5-7.6	Moderate
2.9-7.2	2.2-5.4	High
<2.9	<2.2	Very High

Results of the 2020 FIN at Big Island Lake

Big Island Lake (1582 ha) is located in the Birch Mountains 118 km northwest from the city of Ft. McMurray. From September 1 to 4, 2020, 16 ½-length nets captured 27 Northern Pike, 29 Walleyes, 48 Lake Whitefish, 15 White Suckers and 160 Yellow Perch.

Walleye

The mean catch rate of Walleyes was 1.8/½ net-night. The catch rates of mature (Figure 1) and immature Walleyes were 0.8/½ net-night and 1.0/½ net-night, respectively. The corresponding FSI score for the current mature density of Walleyes was assessed at **very high** risk.




Figure 1 - The FIN catch rate of mature Walleyes from Big Island Lake, 2020. Dashed line is the mean catch rate (1.8 fish/ ½ net-night), with individual net data as shallow circles (n=16 nets).




Figure 3 - The FIN catch rate of mature Northern Pike from Big Island Lake, 2020. Dashed line is the mean catch rate (1.7 fish/ ½ net-night), with individual net data as shallow circles (n=16 nets).




Figure 2 – FIN sample showing size of Walleyes from Big Island Lake, 2020. The red line indicates the average length distribution of Walleye from 5 Alberta lakes supporting long-term sustainable harvests of Walleye.

The length distribution shows moderate but intermittent recruitment with very weak 320 to 610 mm Walleye (Figure 2).

The 2020 FIN sample represented 0.9% of the estimated mature Walleye population size.

Northern Pike

The mean catch rate of mature Northern Pike was 1.2/ ½ net-night (Figure 3). The corresponding FSI score for the current mature density of Northern Pike was assessed at **very high** risk.

The length distribution shows intermittent recruitment of pike, weak abundances of 530 to 670 mm fish, and abundant large fish over 850 mm (Figure 4).

The 2020 FIN sample represented approximately 0.2% of the estimated mature Northern Pike population size.




Figure 4 – FIN sample showing size of Northern Pike from Big Island Lake, 2020. The red line indicates the average length distribution of pike from 6 Alberta lakes supporting long-term sustainable harvests of pike.

Summary

Big Island Lake was assessed in 2014. Since 2014, the status of mature Walleyes has declined from **high** to **very high** risk. Given the Walleye's current risk to sustainability, conservation-based management is required to sustain the population and fishery.

Since the 2014 FIN assessment, the status of mature Northern Pike has remained at **very high** risk. Conservation-focused management will remain the emphasis for this pike population.

Literature

Morgan, G.E. 2002. Manual of Instructions-Fall Walleye Index Netting. Percid Community Synthesis, Diagnostics and Sampling Standards Working Group. Laurentian University, Sudbury Ontario.